

Algal homogeneity in water and sediment of mid-Ethiope river, Nigeria: a response to severe ecological disturbances

Iloba K.I.*


Delta State University, Abraka, Delta State, Nigeria

Received: 08/09/2019, Accepted: 19/12/2020, Available online: 15/01/2021

*to whom all correspondence should be addressed: e-mail: kisiyiloba@gmail.com

<https://doi.org/10.30955/gnj.003214>

Graphical abstract



Abstract

Different aquatic ecological zones exhibit different gradients that present heterogeneous algal composition. The algal community in the water column and sediment with certain environmental factors were screened monthly for twelve months with standard methods, to ascertain disparate in algal distinction. Air and water temperature, conductivity, water depth, dissolved oxygen and nutrients were the primary factors that influenced the algal abundances and diversities. 90% of the algal communities were benthic (33 species) and planktic-benthic (21), with 10% truly planktonic (6) species. Comparatively, the water-column benthic and planktic-benthic surpasses the benthic species associated with the sediment. The PCA on the habitat-type abundance revealed that 99.5% benthic algae in the water column are responsible for the total algal-abundance variance. Species richness and Shannon diversity indices were low. ANCOVA and Levene tests demonstrated significantly equal and homogeneity of algae respectively, between the studied zones. The linear relationships between the algae in the water and sediment zones were non-significant ($p > 0.05$). The Euclidean algorithm confirmed algal community similarity in the water column and sediment. Species diversity limits, a high proportion of benthic algae in the planktonic, and

significant homogenous algae in water and sediment, signatures sediment destruction, breaking and dislodgement and uplifting of associated bottom algae, and the constant mixing of the water column and sediment due to sand dredging in the study area. Machines and human-dredging has exacerbated the abundance of benthic algae in the water column.

Keywords: Dredging, diversity indices, Microflora, univariate analyses, water-river bed, water variables.

1. Introduction

The middle reaches of River Ethiope in the Niger Delta serve as a means of livelihood for the Ethiope East communities living in its watershed for decades. These communities are highly interested in sand mining and dredging as a primary profession, not undermining their trailing problems and other magnetized negative impact anthropogenic activities (Morhit and Mouhir, 2014). The reliance of the communities in and around on this water resource for sand-dredging jobs for sustenance has been on the rise, particularly with the advent of sand dredging machines for small-scale industries (Renuka *et al.*, 2013).

Sand mining in Nigeria is un-regulated despite the stunning policies in place. Sand mining has plagued River Ethiope for decades via the production of very coarse sand of 0.5-1.00mm needed for quality construction (Iloba, 2012). These mechanical events result in water quality degradation, species loss and the destruction aquatic habitats. These environmental-stressor dredgers' employment curates the river bed, impacts on the sediment, algal communities and the water quality via eroding, undercutting and the collapse of river beds and banks (Mugaga and Nabaasa, 2016), notwithstanding its benefits (Grygoruk *et al.*, 2015). Other morphological impacts associated with sand mining include an increase in the river channel, the reconfiguration of the imagery events developed over the years and compromising the aquatic systems' prime condition (Erftemeijer *et al.*, 2012). Algae are heterogeneous autotrophs, distributed heterogeneously in space and time via the creation of distinct habitats, a product of the interactions between the river flow with the sediment (Stancheva and Sheath, 2016). The water column and sediment algal distribution studies

in rivers hypothesized their spatial and temporal distribution in the created distinct habitats (Pla-Rabe's and Catalan, 2018). Therefore, algae distribution in water bodies is naturally temporal and spatial in a relatively calm environment (Hamid *et al.*, 2020). However, the established algal distribution in rivers may be mediated by disturbances such as dredging. Therefore, algal distribution, diversity and richness are tools for detecting disturbance impacts in any aquatic ecosystem; since they have lifestyles associated with habitats and are also highly sensitive to habitat-ecological stressors (El-karim, 2015; Wan Maznah, 2010). Thus, algal species richness, composition, and abundance speak of river system's productivity and trophic dynamics (Jyothi *et al.*, 2016), health (Desianti *et al.*, 2017; Omar, 2010) and specific niche quality and characteristics of the ecological habitat (Estep and Reavie, 2015), regardless of whether pelagic or benthic (Kim *et al.*, 2018) and not underestimated (Colwell, 2012). Evident from works of literature, algal studies in water and sediments of rivers had received considerable attention in the developed world than the underdeveloped nations, particularly in Nigeria (Adeyemo *et al.*, 2008; Tyovenda *et al.*, 2018; Wakoma *et al.*, 2016; Wu *et al.*, 2009). There are no water-sediment algal studies in Ethiope River outside a few studies on heavy metals (Asonye *et al.*, 2007; Kaizer, and Osakwe, 2010), fish (Ikomi *et al.*, 2005; Iloba and Eferakeya, 2019) and macrobenthic invertebrates (Iloba, 2012; Iloba *et al.*, 2019). The limited research and the global use of information on the algal composition, species richness and diversity in the water column and sediments of rivers for development have generated interest and the investigation of this essential economic, natural resource (Desianti *et al.*, 2017). Applications of such information by the developed world in the conservation and management of rivers under intense anthropogenic activities are worth emulation (Apitz, 2012; Ouillon, 2018).

Algae's habitat specificity is first hand in the assessment of dredging impact on algae species composition, and abundance and distribution via the absence or presence of habitat-indicator species, since algae distribution in rivers are not homogenous despite its lotic nature (El-Karm, 2015). Thus the impact of dredging on algae in the sediment and the water column in the mid-Ethiope River was assessed by investigating their composition, abundance and their heterogeneity using different diversity indices. Furthermore, the associations between the algae in both zones with the environmental variables were also evaluated. On this subject, water and sediments samples were collected and scrutinized for algae to evaluate the hypothesized variability in sediment and water column algae due to habitat heterogeneity. Information from this study will establish the River's health as well as guide better water management policies.

2. Materials and methods

2.1. Study area

The steep topography gradient and the morphometry of River Ethiope make the River the final recipient of the terrain-flood and fast-flowing Lowland River. In addition to natural attributes, the River at the middle experiences

several impacts from small-scaled industrial and traditional sand dredging as well as other anthropogenic activities along its course. The River lies within the Niger Delta basin in Delta state, the southern part of Nigeria, with Benin River as its major tributary (Figure 1). The River is a transparent oligoionic freshwater system. Its entire length between is latitude $6^{\circ}31'$ and $6^{\circ}30' N$ and longitude $500-600 E$. The River is tidally influenced partially from Sapele to Aghalokpe, and sometimes it gets at Igun. Its width and depth range from less than 10 m to 90 m and 1m to 20 m, respectively (Iloba, 2017). The river flow is perennial with the highest level and discharges during the flood (July-November).

2.2. Water and algae sampling collection

Water and sediment samples from three sampling stations of 2km apart longitudinally, in the middle reach of the Ethiope River (Figure 1). Firstly, composite water samples from the subsurface, above sediments were collected using a 2-litre Hydro bios sampler and analyzed for dissolved oxygen, temperature, pH, conductivity, alkalinity, nitrate, sulphate and phosphate using APHA (1989). The water and air temperature were determined directly in the field using mercury in the glass thermometer ($0 - 100^{\circ} C$). The use of a calibrated pole measured depth. A pH meter (model 7065) was used to determine the pH.

Alkalinity analyzed by titration method. Dissolved oxygen concentration was by Winkler's method, while Biochemical Oxygen Demand was determined after five days using Winkler's method. Nitrates were determined using the diazotization method. Auto-analyzer II was used to determine the phosphate. Next, composite water column algae samples collected by straining 200L of water with $25 \mu m$ -mesh plankton net, preserved with 4% formaldehyde.




Figure 1. Sampling locations at the middle reaches of Ethiope River at Igun

Sediment samples were collected by pressing a bucket grab into the sediment to a depth of 5cm transferred into a plastic basin. The retrieved sediment samples' temperature and pH were taken and subsequently preserved with 5 % formaldehyde and then taken to the laboratory. In the laboratory, sediment algae were collected by the modified floatation method in a plastic basin with distilled water and sieved through a $25 \mu m$ mesh-plankton net. The algal samples were viewed at 40x magnification using an Olympus binocular microscope.

Taxonomic denominations and identifications were made by consulting (AHPA, 1989; Wehr and Sheath 2003; Wehr *et al.*, 2015; Withford and Schumacher, 1973).

2.3. Statistical analysis

To test statistically significant differences between the water column's monthly physicochemical parameters and sediment, one way ANOVA was performed. One way ANCOVA was used to establish linear regression slopes and their equalities for water and sediment algae for all studied zones. Levene's test was performed to appraise the algae's sameness in the water column and sediment algae tested at 0.05 confidence limits as the impact of dredging. Linear correlation analysis and principal component analysis were performed to buttress similarities between algae's relative abundances in water and sediment among the studied stations. Euclidean distance elucidated the similarities in

habitat-type in the three studied stations. Pearson linear correlation applied on log x+1 transformed abundances, and environmental variables, correct non-uniformity in metric scale and taxa variability and to detect the most influential parameters on algal abundances among studied stations. Species diversity and individual richness were used to predict algae's stability (expected trend) in both water habitats using (Colwell *et al.*, 2012; Hammer *et al.*, 2001).

3. Results

The overview of the outcome of water quality variables in Table 1, identified variations within the different study zones. However, no significant variations were found between the different zones except in nitrate ($p < 0.05$) (Table 1).

Table 1. The summary of mean, coefficient of variation (C.V %) in parenthesis, range, F and P values for the sediments and pelagic zone stations (August to December 2011 and 2012) Mean \pm SE

Stations	Station 1		Station 2		Station 3		F	p
	ST1SD	ST1W	ST2SD	ST2W	ST3SD	ST3W		
Air temperature (°C)	21.7(1.67)	22.0(2.0)	22.8(0.31)	23(1.82)	24.0(1.6)	24.6(1.63)	1.30	0.87
Range	17.4–26.0	15.0–25.0	19.0–26.7	18–27.0	20.2–27.8	20.0–30.0		
Water temp.	25.0(1.77)	27.8(1.39)	28.8(2.29)	30.2(2.24)	29.8(1.76)	31.4(0.98)	1.91	0.69
Range	20.5–29.6	25.0–32.0	23.0–34.7	25.2–38.0	25.3–34.4	30.0–35.0		
Depth (m)	8.6(0.30)	7.5	8.3(0.31)	7.0()	9.2(0.34)	8.0(0.29)	1.19	0.91
Range	7.8–9.4	6.8–8.4	7.5–9.05	6.5–8.1	8.0–10.0	7.0–9.0		
Conductivity	30.8	66.8(2.63)	23.9()	62.2(2.65)	27.4()	61.2(3.31)	1.33	0.85
Range (μ S/cm)	3.22–58	60.0–74.0	5.76–47.8	53–69.0	9.7–39.8	50.0–69.0		
pH	3.7(0.16)	3.7(0.17)	3.8(0.16)	3.8(0.17)	4.0(0.10)	4.04(0.07)	1.5	0.8
Range	3.3–4.2	3.2–4.2	3.4–4.2	3.2–4.2	3.7–4.2	3.8–4.2		
Dissolved oxygen	6.18(0.55)	6.3(0.64)	4.8(0.52)	4.8(0.64)	5.7(0.72)	5.74(0.88)	1.16	0.92
Range	4.8–7.6	4.0–7.5	3.5–6.17	2.5–6.2	3.8–7.5	3.1–8.5		
Biochemical oxygen demand	5.4(0.40)	5.6(0.44)	5.0(0.46)	5.1(0.55)	4.8(0.4)	4.8(0.48)	1.78	0.72
Range	4.3–6.5	4.7–6.8	3.8–6.2	3.5–6.5	3.8–5.8	3.2–6.0		
Phosphate	0.33(0.06)	0.24(0.03)	0.40(0.06)	0.32(0.04)	0.35(0.35)	0.30(0.03)	1.54	0.79
Range	0.15–0.48	0.16–0.32	0.3–0.60	0.19–0.42	0.20–0.45	0.22–0.40		
Nitrate	1.5(0.07)	0.23(0.02)	0.3(0.04)	0.30(0.03)	1.70(0.07)	0.30(0.03)	356.25	0.005*
	1.30–1.70	0.18–0.30	0.20–0.40	0.24–0.38	1.30–1.70	0.18–0.38		

A total of sixty (60) species of phytoplankton were detected in the water and sediment samples during this study, of which 46.67 % was Bacillariophyta (28 species), 31.67% Chlorophyta (19 species), 20 % Cyanophyta (12 species) and 1.67 % Euglenophyta (1 species) (Table 2). 90% of the algal communities were benthic (33 species) and planktic-benthic (21), with 10% truly planktonic (6) species. Total sediment algal density varied from 189 ind ml⁻¹ to 200 ind ml⁻¹ lower than water column algal species which varied from 200 ind ml⁻¹ to 248 ind ml⁻¹. The algal composition in the water column and the sediment were dominated by *Fragilaria javanica*, *Navicula tenelloides*, *Synedra acus* and *Tabellaria fenestrata* (Bacillariophyceae), as shown in Table 2 and Figure 2A. In this study, 66.67% of the algal community had species primarily with 1-5cells/ml except for few species. A few species such as *Fragilaria javanica* (6-70cells/ml), *Navicula tenelloides* (1-45 cells/ml), *Synedra acus* (1-25cells/ml) dominating the group Bacillariophyta, and *Pleurotaenium sp.*, *Scenedesmus incrassatulus*,

Spirogyra insignis had 6-25cells/ml each, dominating for the Chlorophytes (Table 2). *Merismopedia elegans* dominates the group Cyanophyta while others (1-5cells/ml) but one; *Microcystis aureginous*, were widely distributed in the water columns and sediments.

Microcystis aureginous/ viridis registered only at the bottom (sediment). The linear model generated between water and sediment algae compositions (Figure 2B) revealed no marked distinction between water column's algae community and sediment. The co-variance test for equal means for the three significant taxa, Bacillariophyta, Chlorophyta and Cyanophyta, in water, sediment and water-sediment algae revealed no statistically different taxa ($F=3.111$; $p=0.09$). Levene's test for homogeneity of variances for means ($p=0.1214$) and median ($p=0.1243$) equally detected no significant difference. The correction matrix map for the three significant taxa; Bacillariophyta, Chlorophyta and Cyanophyta in water, sediment and

water-sediment are shown in Figure 2C. The environmental variables that influence the major taxa are shown in Table.

Table 2. Species Checklist, distribution, abundance, diversity indices and ecological habitats (in parenthesis **BOLD**) of the algae community in the water-columns and sediments of mid-Ethiopia River.

	Acronym	STN 1		STN 2		STN 3		RA	H	E	
		ST1SD	ST1W	ST2SD	ST2W	ST2SD	ST3W				
Bacillariophyceae											
1	<i>Achnantes minutissima v affinis</i> (B)	AMVA	-	++	-	++	-	+	1.82	1.39	0.67
2	<i>Aulacoseira granulate</i> (P-B)	AUGR	+	+	+	+	+	+	1.5	1.75	0.96
3	<i>Asterionella formosa</i> (P)	ASFO	+	+	+	+	+	+	0.55	1.75	0.96
4	<i>Asterionella sp</i> (P)	ASSP	+	+	+	+	+	+	1.26	1.74	0.95
5	<i>Cyclotella comta</i> (P-B)	CYCO	+	++	+	++	+	++	2.84	1.55	0.73
6	<i>Cymbella gracilis</i> (B)	CYGR	+	+	+	+	+	+	0.79	1.74	0.96
7	<i>Diatom enlongatum</i> (P-B)	DIEN	++	+	++	+	++	+	3.71	1.68	0.89
8	<i>Diatoma hyemale</i> (P-B)	DIHY	-	+	-	+	-	+	0.47	1.67	0.89
9	<i>Diatoma vulgare</i> (P-B)	DIVU	+	+	+	+	+	+	0.71	1.77	0.93
10	<i>Eunotia monodon</i> (B)	EUMO	+	-	+	-	+	-	0.87	1.39	0.80
11	<i>Eunotia sudetica</i> (B)	EUSU	++	-	++	+	++	+	2.61	1.69	0.90
12	<i>Eunotia serra</i> (B)	EUSE	+	+	+	+	+	+	1.19	1.69	0.90
13	<i>Fragilaria javanica</i> (B)	FRJA	++	++++	++	-	++	++++	11.14*	1.32	0.62
14	<i>Mastogloia smithi</i> (B)	MASM	+	+	+	+	+	+	0.87	1.78	0.98
15	<i>Melosira granulata</i> (B)	MEGR	+	+	+	+	+	+	0.71	1.75	0.96
16	<i>Luticola muticola</i> (B)	LUMU	+	+	+	+	+	+	0.79	1.77	0.98
17	<i>Navicula viridula</i> (B)	NAVI	+	+	+	+	+	+	0.63	1.74	0.95
18	<i>Navicula tenelloides</i> (B)	NATE	++++	++	+++	++	+++	++	6.79	1.75	0.96
19	<i>Peronia intermedium</i> (B)	PEIN	+	+	+	+	+	+	0.95	1.76	0.97
20	<i>Pleurosigma obtusatum</i> (B)	PLOB	-	+	+	+	-	+	0.47	1.72	0.93
21	<i>Quadrigula closteriodes</i> (B)	QUCL	+	-	+	-	+	-	0.32	1.52	0.92
22	<i>Rhopalodia gibba</i> (B)	RHGI	+	+	+	+	+	+	0.55	1.75	0.96
23	<i>Stauroneis phoenicentron</i> (B)	STPH	-	-	+	-	+	-	0.16	1.55	0.94
24	<i>Surirella sp</i> (B)	SUSP	-	++	-	++	-	+	1.58	1.37	0.65
25	<i>Synedra acus</i> (P-B)	SYAC	+++	+	+++	++	+++	+	6.32	1.58	0.81
26	<i>Synedra ulna</i>	SYUL	+	-	+	+	+	-	0.71	1.37	0.79
27	<i>Tabellaria fenestrata</i> (P-B)	TAFE	++	+	++	+	++	+	3.08	1.60	0.83
28	<i>Tabellaria flocculosa</i> (B) <i>minina</i> (P-B)	TAFL	++	+	+	+	+	+	1.5	1.70	0.91
Chlorophyceae											
29	<i>Closterium acerosum</i> (P-B)	CLAC	-	+	-	++	-	+	0.87	1.32	0.62
30	<i>Closterium incurvum</i> (P-B)	CLIN	+	+	+	++	+	+	1.58	1.49	0.74
31	<i>Closterium sigmoideum</i> (P-B)	CLSI	+	+	++	+	+	+	1.74	1.75	0.96
32	<i>Closterium granatum</i> (P-B)	CLGR	+	-	+	-	-	-	0.16	1.73	0.94
33	<i>Chlorella Sp</i> (P-B)	CHSP	+	-	+	-	+	-	0.32	1.70	0.91
34	<i>Desmidium asymeticum</i> (P-B)	DEAS	+	++	+	++	-	+	1.74	1.55	0.78
35	<i>Gonatozygon knahani</i> (B)	GOKN	+	+	+	+	+	+	1.26	1.77	0.98
36	<i>Mougeotia parvula</i> (B)	MOPA	-	-	-	++	-	++	1.66	1.15	0.53
37	<i>Mougeotia sp</i> (B)	MOSP	-	-	-	-	-	++	1.01	1.00	0.45
38	<i>Mougeotiosis sp</i> (B)	MOUG	-	-	-	-	-	++	0.71	0.85	0.39
39	<i>Oedogonium grande</i> (B)	OEGR	+	+	+	+	+	+	1.66	1.77	0.98
40	<i>Pleurotaenium sp</i> (P-B)	PLSP	-	+++	-	++	-	-	2.45	1.10	0.50
41	<i>Scenedesmus ecornis v. polymorphus</i> (P)	SEVP	+	+	+	+	+	+	0.63	1.74	0.95
42	<i>Scenedesmus incrassatulus</i> (P)	SCIN	+++	-	+++	-	+++	-	4.19	1.30	0.61
43	<i>Spirogyra africana</i> (B)	SPAF	+	+	+	+	+	+	1.11	1.78	0.99
44	<i>Spirogyra insignis</i> (B)	SPIN	-	++	-	+++	-	++	2.92	1.33	0.63
45	<i>Spirogyra sp</i> (B)	SPSP	-	++	++	-	++	-	3.00	1.34	0.64
46	<i>Ulothrix tenuissima</i> (B)	ULTE	+	+	+	+	+	+	1.34	1.78	0.99
47	<i>Zygorium ericetorum</i> (B)	ZYER	-	+	+	+	+	+	0.71	1.71	0.93
Cyanophyceae											
48	<i>Aphanizomenon sp</i> (P)	APSP	+	-	+	-	+	-	0.79	1.52	0.76
49	<i>Merismopedia elegans</i> (P-B)	MEEL	++	++	++	++	++	++	3.32	1.79	1.00
50	<i>Microcystis aureginous</i> (P-B)	MIAU	++	-	++	-	++	-	1.74	1.42	0.69

	Acronym	STN 1		STN 2		STN 3		RA	H	E	
		ST1SD	ST1W	ST2SD	ST2W	ST2SD	ST3W				
51	<i>Microcystis viridis</i>	MIVI	+	+	+	+	+	+	1.19	1.75	0.96
52	<i>Oscillatoria brevis</i> (P-B)	OSBR	+	+	+	+	+	+	1.34	1.74	0.95
53	<i>Oscillatoria proboscidea</i> (P-B)	OSPR	+	+	+	+	+	+	1.26	1.76	0.97
54	<i>Phormidium ambiguum</i> (B)	PHAM	+	+	+	+	+	+	0.79	1.78	0.98
55	<i>Phormidium ingricum</i> (B)	PHIN	+	+	+	+	+	+	1.19	1.71	0.92
56	<i>Phormidium mucicola</i> (B)	PHMU	+	+	+	+	+	+	1.34	1.78	0.98
57	<i>Phormidium tenius</i> (B)	PHTE	+	+	+	+	+	+	1.34	1.75	0.96
58	<i>Schizothrix sp</i> (B)	SCSP	+	+	+	+	+	+	0.71	1.77	0.98
59	<i>Spirulina major</i> (P-B)	SPMA	+	-	+	-	+	-	0.32	1.70	0.91
Euglenophyceae											
60	<i>Euglena oxyuris</i> (P)	EUOX	+	-	+	-	+	-	0.71	1.06	0.96
Diversity Indices	Taxa		47	46	51	47	47	45			
	Individuals		189	248	193	210	200	200			
	Dominance		0.05	0.01	0.05	0.04	0.05	0.05			
	Shannon		3.33	3.07	3.43	3.57	3.04	3.36			
	Evenness		0.59	0.47	0.60	0.76	0.61	0.61			

Note: (-):0cell/ml; +:1-5cells/ml; ++: 6-15cells/ml; +++: 16-25; ++++:26-45cells/ml; 46-69 :+++++:> 70cells/ml; H: Habitat: (B: Benthic; P: Planktonic; P-B: Planktonic-Benthic) RA: relative abundance; H: Shannon index; E: Evenness.

Table 3. Similarity and distance indices and correlation coefficients in parenthesis between benthic, planktonic, planktonic-benthic algae associations in water column and sediments zones (A) and Pearson correlation and p values of environmental variables and major taxa(B) in mid-Ethiopia River

A	ST1-B	ST1- P	ST1 -P-B	ST2-B	ST2-P	ST2-P-B	ST3-B	ST3-P	ST3-P-B
ST1-B	0	92.5 (0.50)	104.0(0.14)	68.7(0.00)	91.4(0.51)	102.7(0.12)	36.0(0.0)	91.1 (0.50)	99.4 (0.16)
ST1-P	92.5(-0.09)	0	50.0(0.52)	51.8(0.24)	2.7(0.0)	52.3(0.36)	73.3(0.36)	3.9 (0.0)	46.9 (0.45)
ST1-P-B	104.0(-0.20)	50.0(-0.09)	0	65.7(0.02)	50.9(0.50)	18.0(0.0)	86.4(0.04)	50.2 (0.49)	20.4 (0.0)
ST2-B	68.7(0.62*)	51.8(-0.16)	65.7(-0.33)	0	50.7(0.25)	67.0(0.0)	47.0(0.0)	50.2 (0.25)	62.0 (0.03)
ST2-P	91.4(-0.09)	2.7(0.99*)	50.9(-0.09)	50.7(-0.15)	0	52.5(0.36)	72.3(0.38)	4.5 (0.0)	47.4 (0.43)
ST2-P-B	102.7(-0.21)	52.3(-0.12)	18.0(0.91*)	67.0(-0.35)	52.5(-0.12)	0	85.9(0.03)	52.0 (0.35)	22.2 (0.0)
ST3-B	36.0(0.92*)	73.3(-0.12)	86.4(-0.28)	50.0(0.68*)	72.3(-0.12)	85.9(-0.28)	0	72.0 (0.37)	82.1 (0.06)
ST3-P	91.1(-0.09)	3.9(0.99*)	50.2(0.09)	50.2(-0.16)	4.5(0.97*)	52.0(-0.13)	72.0(-0.12)	0	46.9 (0.43)
ST3-P-B	99.4(-0.19)	46.9(-0.10)	20.4(0.88*)	62.0(-0.30)	47.4(-0.10)	22.2(0.86*)	82.1(-0.24)	46.9 (-0.10)	0

B	Air-T	Water-T	Depth	Conductivity	D O	BOD	PO4	NO4
Bacillariophyta	-0.84(0.04)				0.81(0.05)		-0.79(0.059)	
Chlorophyta			-0.91(0.01)	0.88(0.02)				-0.83(0.04)
Cyanophyta						-0.82(0.04)		
Conductivity			-0.82(0.05)				-0.84(0.04)	
NO3			0.81(0.04)					
BOD		-0.83(0.04)						

The species-diversity indices limits at the species level were low (Table 2). Shannon indices and evenness varied from 0.85 to 1.79 and from 0.39 to 1.0 for *Merisimopedia elegans* (Cyanophyceae) and *Mougeotiosis sp* (Chlorophyceae) respectively. Aggregately, the community diversity indices at the different stations and zones revealed very high Shannon indices of $H > 3.0$, with the

correlation coefficient amongst the algae abundances of the studied stations very high ($r > 0.90$). The least correlation coefficient was 0.9769 for between station 1 sediment (ST1SD) and Station3 water column (ST3W), while the maximal correlation coefficient was 0.99915 between Station 2 sediment (ST2SD) and Station 3 sediment (ST3SD). Shannon diversity t-test detected highly statistical differences between algae composition in station

1 bottom(ST1-B) and Station 2 bottom(ST2-B) (t :-4.5165; df :429.41, p : 8.1354E-06; Station 1 bottom (ST1-B) (2.7124) and Station 3 bottom(ST3-B) (3.1523) t :-4.5165, t : 8.1354E-06 and non-statistical differences between Station 2 bottom (ST2-B) and Station 3 bottom (ST3-B), t :1.2617, df : 443.98, p : 0.20778.

The principal components analysis clustered the identified species at the central axis, shown in Figure 3A. The first and the second axes revealed that 82% of the algal composition influence algal variability among the studied stations in the mid-Ethiopian River. The studied zones' position separated the sediments of the three stations, stations 1 and 2 water column and station 3 as three different zones. Figure 3C identified the high strong positive association between benthic algae with the water column. The total number of individuals/species influenced the sediments algae positively while the P and P-B influenced one another positively along the negative axis.

The first axis isolating water column and benthic algae accounted for 99.5% of the total algae variance, separated from the water-sediment and the total algal and planktonic-benthic and benthic. The PCA further confirmed by the Euclidean similarity and distance indices (Table 3A). Although, some similarities (small distances) surprisingly were not correlated (Table 3A). Analyzing the algal community upon habitat/life forms preference, benthic and planktonic-benthic species dominated all stations and zones of mid-Ethiopian River, and real planktonic species were very few (Table 1 and Figure 3B). The affinity between the betwixt the physicochemical parameters and the algae abundances are in Table 3B.

4. Discussion

In the present study, water temperature, air temperature, conductivity, water depth, dissolved oxygen and nutrients ($p < 0.05$) factors influenced algae abundance and diversities in middle River Ethiopia (Table 3B). Thus, biological events in aquatic ecosystems are influenced by these environmental variables. Hou *et al.* (2018), reported that changes in the physicochemical variables modify algal abundance, composition and algae's diversities in rivers.

The geomorphic and inherent fundamentals are primers in structuring and controlling aquatic communities (Hou *et al.*, 2018; Onyema and Popoola, 2013; Wang *et al.*, 2016; Wu *et al.*, 2009). Ethiopian rivers' sediment acidic nature (3.7 to 4.2) and high BOD (4.5 to 6.8mg/l) values inferred higher oxygen consumption zones than in the water column. However, the BOD at both zones was more significant than 3, which infers the impact of released nutrients and bottom-associated vegetation decomposition from the sediment, thereby impairing the river water quality (Pan *et al.*, 2013; Wang *et al.*, 2016). This is also evident in the nutrient-enriched River's sediments noted in this study. Which may be attributable to nutrients released from the sediment during the dredging processes (Hamid *et al.*, 2020). Rich nutrient sediments are crucial in the abundance of the blue-greens in any aquatic ecosystem but not evident in the present study (Adeyemo *et al.*, 2008). This may be attributed to insufficient light due to its absorption

by solids, sand slits in the water column, which is evident from the reduced transparency in this section of the river (Iloba, 2012). The negative association between the Cyanophytes and BOD ($r = -0.8241$; $p = 0.04$), and Chlorophytes versus nitrate ($r = -0.8273$; $p = 0.04$) (Table 3B) was contrary to natural expectations, signature other underlying determining factors in the study area. However, differential responses of algal community to nutrients or organic enrichment have been noted in other water and sediment studies (Desianti *et al.*, 2017). The positive correlation of dissolved oxygen on Bacillariophyta ($r = 0.8137$, $p = 0.0478$) indicates its positive influence on the proliferation of Bacillariophyta in this system. The relatively high amount of dissolved oxygen is due to exchange between the air-water interphase (Hamid *et al.*, 2020).




Figure 2. Taxa distribution(A), linear relationship (B) and the correlation matrix map(C) between water, sediment, water-sediment algae at the various stations (Acronyms in Table 2)

This association is attributable to the dominance of Bacillariophyceae in this study and exertion of the River's flow on algal assemblages and habitat destruction by dredge through dislodgement and nutrients enhancement as demonstrated in this study (Berge *et al.*, 2010). The predominance of diatoms in this river is contrary to the noted dominance of Chlorophytes in rivers. This may be due to the out-competition of the chlorophytes by the benthic. Benthic algae are essential indicators of human disturbances (Gokce, 2016). Abonyi *et al.* (2014) opined that the algae biomasses in the middle reaches of large rivers are exceptionally high in line with the river continuum concept (RCC) (Vannote *et al.*, 1980). A high number of species and abundance is partially valid for mid-River Ethiopia with only 60 species, which is incomparable

between the water column and sediment algae which was further confirmed by the non-violation of equal co-variance of algae in the studied zones ($p > 0.05$). The algal homogeneity is therefore not random, but due to dredging impact. Researchers have demonstrated that the impact of human-disturbance in rivers more significant than the interactions between algae and water (Ishaq and Khan, 2013; X-Gao *et al.*, 2018). Contrary to species diversity, the relatively zonal biological diversities (Shannon (H) > 3) at the different stations portray stable ecological zones (Jyothi *et al.*, 2016). Researchers have reported a reduction in species diversity and richness through habitat destruction, disruption in water and sedimentary environment (X-Gao *et al.*, 2018). The algal composition and abundance in the studied area are unknown to assess the magnitude of damage and alteration of the River's habitat and local biota for comparison but is already threatened evidenced from the low biological indices (Iloba *et al.*, 2019 and Hamid *et al.*, 2020). The non-significant and significant differences demonstrated by the Shannon diversity t-test predict niche overlap among the studied sites. However, species diversity indices are an unparalleled tool to the taxon to taxon analyses, which evidenced stressed condition for the algal species. These findings with Iloba (2017) in microfauna evaluation of Ethiopie River, opined that diversity could be high in a stressful environment.

5. Conclusion

This research evaluated the algal composition and abundance distinctiveness in the water column, and sediment of the middle reaches of the Ethiopie River and observed no heterogeneity. The study identified 60 species of algae belonging to four taxa; Bacillariophyta, Chlorophyta, Cyanophyta and Euglenophyta. The study has substantiated that the algal community in the water column was higher than those of the sediment due to upload and release of algae into the water column in response to the dredging activities; however, the algal communities of both zones were dominantly attached algae (tychoplankton and euplankton). Furthermore, the algal communities at the different zones demonstrated no distinction among the study sites when subjected under different statistical evaluations ($p > 0.05$), thus, revealed no habitat heterogeneity. Thus the destruction of the natural river bed of this River through sand mining and dredging has created a river with significant slits or organic loads. The study envisioned ecological disaster in future if sand mining is not put under control. The boundary disappearance between sediments and water column also indicts the sand dredging activities in the study area. Sand dredging and its associated plumes mixed and impact uniformity between the water column's algal community structure and sediment. Thus, the water column and sediment algal communities in the study are similar. The low species richness and low Shannon diversity indices revealed a disturbed and unstable river indicting observed human interferences; dredging. The algal communities in the water-sediment of the middle reaches of the Ethiopie River have been severely altered and homogenized from

bottom to top. This situation calls for a functional regulatory body to constrain the dredging activities in the area, abate habitat destruction, and avoid socioeconomic disaster. Law enactment will also monitor these activities or put a hold on sand dredging, to allow sediment recovery time before engaging in regulatory and guided dredging.

Acknowledgements

The author appreciates the assistance of the Department of Animal and Environmental Biology laboratory technologists in the analysis of the water samples.

References

- Abonyi A., Leitao M., Stankovic I., Borics G and Varbiro G. (2014), A large river (River Loire, France survey to compare Phytoplankton functional approaches: Do they display river zones in a similar way?, *Ecological Indicators*, **46**, 11–22. <http://dx.doi.org/10.1016/j.ecolind.2014.05.038>.
- Adeyemo O.K., Adedokun O.A. and Adeleye E.A. (2008), Seasonal changes in physico-chemical parameters and nutrient load of river sediments in Ibadan City, Nigeria, *Global NEST Journal*, **10**, (3), 326–336.
- Anderson J.N. (2014), Landscape disturbance and lake response: Temporal and spatial perspectives, *Freshwater Reviews*, **7**(2), 77–120.
- APHA (1989), *American water works Association*. Water pollution control. Federation Standard method for the examination of water and wastewater 17th Edn.
- Apitz S.E. (2012), Conceptualizing the role of sediment in sustaining ecosystem services: Sediment-ecosystem regional assessment (SEcoRA), *Science of the Total Environment*, **415**, 9–30.
- Asonye C.C., Okolie N.P., Okenwa E.E. and Iwuanyanwu U.G. (2007). Some physico-chemical characteristics and heavy metal profiles of Nigerian rivers, streams and waterways, *African Journal of Biotechnology*, **6** (5), 617–624.
- Berge T., Daugbjerg N., Anderson B.B. and Hansen P.J. (2010), Effects of lowered pH on marine phytoplankton growth rate, *Marine Ecology Progress Series*, **416**, 79–91.
- Breuer F., Janz P., Farrelly E. and Ebke K.P. (2017), Environmental and structural factors influencing algal communities in small streams and ditches in central Germany, *Journal of Freshwater Ecology*, **32**(1), 65–83. DOI:10.1080/02705060.2016.1241954.
- Colwell R.K., Chao A., Gotelli N.J., Lin S., Mao C.X., Chazdon R.L., and Longino J.T. (2012), Models and estimators linking individual-based and sample-based rarefaction extrapolation and comparison, *Journal of Plant Ecology*, **5**(1), 3–21.
- Desianti N., Potapova M., Enache M., Belton T.J., Velinsky R., and Mead J. (2017), Sediment diatoms as environmental indicators in New Jersey Coastal Lagoon, *Journal of Coastal Research*, **78**, 127–140.
- El-Karim M.S.A. (2015), Survey to compare phytoplankton functional approaches: How can these approaches assess River Nile water quality in Egypt, *Egyptian Journal of Aquatic Research*, **41**, 247–255.
- Erfteemeijer P.L.A., Riegl B., Hoeksema B.W. and Todd P.A. (2012), Environmental impacts of dredging and other sediment disturbances on corals: A Review, *Marine Pollution Bulletin*, **64**, 1737–1765. DOI:10.1016/j.marpolbul.2012.008-014-0017-7.

- Estep L.R. and Reavie E.D. (2015), The ecological history of Lake Ontario according to phytoplankton, *Journal of Great Lakes Research*, **41**, 669–687. <http://dx.doi.org/10.1016/j.jglr.2015.06.005>.
- Gao X., Lee J.R., Park S.K., Kim N.G. and Choi H.G. (2018), Detrimental effects of sediment on attachment, survival and growth of the brown alga *Sargassum thunbergii* in early life stages, *Phycological Research*, doi: 10.1111/pre.12347.
- Gokce D. (2016). *Algae as an indicator of water quality*. Doi:10.5772/62916.
- Grygoruk M., Frąk M. and Chmielewski A. (2015), Agricultural rivers at risk: Dredging results in a loss of macroinvertebrates. Preliminary observations from the Narew Catchment, Poland. *Water*, **7**, 4511–4522. Doi:10.3390/w7084511.
- Hamid A., Bhat S.U. and Jehangir A. (2020), Local determinants influencing stream water quality. *Allpied Water Science*, **10**, 24, doi.org/10.1007/s13201-019-1043-4.
- Hammer O., Harper D.A.T. and Ryan P.D. (2001), PAST: Paleontological statistics software package for education and data analysis, *Palaeontologia Electronica*, **4**(1), 9pp. http://palaeo-electronica.org/2001_1/past/issue1_01.htm.
- Hassan F.M., Salman J.M., Alkam F.A. and Jawad H.J. (2014), Ecological Observation on Epipelagic Algae in Euphrates River at Hindiya and Manathira, *International Journal of Advanced Research*, **2**(4), 1183–1194.
- Heneash A.M.M., Tadrose H.R.Z., Hussein M.M.A., Hamdona S.K., Abdel-Aziz N. and Gharib S.M. (2015), Potential effects of abiotic factors on the abundance and distribution of the plankton in the Western Harbour, south-eastern Mediterranean Sea, Egypt, *Oceanologia*, **57**, 61–70
- Heneash A.M.M., Tadrose H.R.Z., Hussein M.M.A., Hamdona S.K., Abdel-Aziz N. and Gharib S.M. (2015), Potential effects of abiotic factors on the abundance and distribution of the plankton in the Western Harbour, south-eastern Mediterranean Sea, Egypt. *Oceanologia*, **57**, 61–70
- Hou Z., Jiang Y., Liu Q., Tian Y., He K. and Fu L. (2018), Impacts of environmental variables on a phytoplankton community: A case study of the tributaries of a subtropical river, Southern China, *Water*, **10**, 152. Doi: 10.3390/w10020152.
- Ikomi R.B. and Arimoro F.O. (2014), Effects of Recreational Activities on the Littoral Macroinvertebrates of Ethiope River, Niger Delta, Nigeria, *Journal of Aquatic Sciences*, **29** (1B): 155–170.
- Ikomi R.B., Arimoro F.O. and Abale E.F. (2005), Some aspects of the biology of *Thysia anisorgii* (Boulenger, 1899) (Cichlidae) in River Ethiope, Delta state, Nigeria, *Journal of Tropical Sciences*, **5**(2), 1–7.
- Iloba K.I. (2012), *The Physico-chemical Characteristics and Plankton of Ethiope River, Nigeria*. PhD Thesis, Department of Animal and Environmental Biology, Delta State University Abraka, Delta State.
- Iloba K.I. (2017). Rotifers of River Ethiope, Delta State, Nigeria, *International Journal of Fisheries and Aquatic Studies*, **5**(2), 74–79.
- Iloba K.I. and Eferakeya E. (2019), Length – weight relations and condition factors of three freshwater prawns of Ethiope River, Nigeria, *Journal of Agriculture and Food Environment*, **6**(1), 19–31.
- Iloba K.I., Akawo N. and Anani C. (2019), Sand dredging impact on macrobenthic invertebrates of a hallowed River in Delta State of Nigeria, *Science World Journal*, **14**(1), 171–176.
- Iloba K.I., Akawo O.N. and Nwaefiense F. (2018), Diversity and Community Structure of Macroinvertebrates in Anthropogenically stressed water body in Delta state, Nigeria, *International Journal of Applied Biological Research*, **9**(1), 93–206.
- Ishaq F. and Khan A. (2013), Comparative assessment of physico-chemical conditions and Plankton diversity of River Tons and Asan in Dehradun District of Uttarakhand. *Advances in Applied Science Research*, **4**(2), 342–355.
- Jones R., Fisher R., Stark C., Ridd P. (2015), Temporal patterns in seawater quality from dredging in tropical environments, *PLoS ONE* **10**(10), e0137112. doi:10.1371/journal.pone.0137112.
- Jyothi K., Krishna P.M. and Mohan N.R.G. (2016), Algae in fresh water ecosystem, *Phykos*, **46**(1), 25–31.
- Kaizer A.N and Osakwe S.A. (2010), Physicochemical characteristics and heavy metal levels in water samples from five river systems in Delta State, Nigeria, *Journal of Applied Sciences and Environmental Management*, **14**(1) 83–87.
- Kim J., Lee T. and Seo D. (2018), Algal bloom prediction of the lower Han River, Korea using the EFDC hydrodynamic and water quality model, *Ecological Modelling*, **366**, 27–36.
- Krumins J., Adick van Oevelen., Bezemer T.M., De Deyn G.B., Gera Hol W.H., Vav Donk E., Wietse de Boer De Ruiters P.C. and Middelburg J.J. (2013), Soil and Freshwater and Marine Sediment Food Webs: their Structure and Function, *Bioscience*, **63**(1), 35–42.
- Mabidi A., Bird M.S. and Perissinotto R. (2017), Distribution and diversity of aquatic macroinvertebrate assemblages in a semi-arid region earmarked for shale gas exploration (Eastern Cape Karoo, South Africa), *PLoS ONE*, **12**(6), 1–27. e0178559. <https://doi.org/10.1371/journal.pone.0178559>.
- Moreira Gonzalez A.R., Brustolin M.C., Junior L.L.M. (2020), Composition and abundance of benthic microalgae from the estuarine complex of Paranaguá Bay (Southern Brazil) with special emphasis on toxic Species, *Ocean and Coastal Research*. <http://dx.doi.org/10.1590/52675-28242020068276>.
- Morhit M.E. and Mouhir I. (2014), Study of physico-chemical parameters of water in the Loukkos river estuary (Larache, Morocco), *Environmental Systems Research*, **3**, 17. DOI:10.1186/s40068.
- Mugaga F. and Nabaasa B.B. (2016), The centrality of water resources to the realization of sustainable Development Goals (SDG). A review of potentials and constraints on the African continent, *International Soil and Water Conservation Research*, **4**, 215–223.
- Murnaghan S., Taylor D. and Jennings E. (2015) Reconstructing long-term trophic histories for lakes using two independent approaches: Application of dynamic computer modelling and palaeolimnology to Lough Mask, Ireland, *Biology and Environment: Proceeding of the Royal Irish Academy*, **115**(3), 171–189.
- Omar W.M.W. (2010). Perspectives on the use of algae as biological indicators for monitoring and protecting aquatic environments, with Special Reference to Malaysian Freshwater Ecosystems, *Tropical Life Sciences Research*, **21**(2), 51–67.
- Onyema I.C. and Popoola R.T. (2013), The physicochemical characteristics, chlorophyll a levels and phytoplankton

- dynamics of the East Mole Area of the Lagos Harbour, Lagos, *Journal of Asian Scientific Research*, **3**(10), 99501019.
- Ouillon S. (2018), Why and how do we study sediment transport? Focus on coastal zones and ongoing methods, *Water*, **10**, 390. Doi: 10.3390/w10040390.
- Pan J., Bournaod C.N., Cuadrado D.G., Vitale A. and Piccolo M.C. (2013), Interaction between estuarine microphytobenthos and physical forcings: The role of atmospheric and sedimentary factors. *International Journal of Geosciences*, **4**, 352–361.
- Ray J.G. and Thomas T.B (2012), Ecology and diversity of green-algae of tropical oxic dystrophic soils in relation to different soil parameters and vegetation, *Research Journal of Soil Biology*, **4**(3), 42–68.
- Renuka N., Sood A., Prasanna R. and Ahluwalia A.S. (2013), Influence of seasonal variation in water quality on the microalgal diversity of sewage wastewater, *South African Journal of Botany*, **90**, 137–145. <http://dx.doi.org/10.1016/j.sajb.2013.10.017>.
- Salman J.M., Jawad H.J., Nassar A.J. and Hassan F.M. (2013), A study of Phytoplankton Communities and Related Environmental factors in Euphrates River (between Two Cities: Al Musayyab and Hindiya), Iraq. *Journal of Environmental Protection*, **4**:1071–1079.
- Stancheva R., and Sheath R.G. (2016), Benthic soft-bodied algae as bioindicators of stream water quality. *Knowledge and Management of Aquatic Ecosystems*, **417**, 15, Doi:10.1051/kmae/2016002.
- Turner C.R., Uy K.L. and Everhart R.C. (2015), Fish environmental DNA is more concentrated in aquatic sediments than surface water, *Biological Conservation*, **183**, 93–102.
- Tyovenda A.A., Ikpughul S.I., Ayua T.J., Aligba E.H., Philip I.A., Igbawua T., Aondoakaa S.I. and Korna R.T. (2018), Assessment of radionuclides in water, sediment and algae in river benue at Jimeta-Yola, Adamawa State, Nigeria, *International Journal of Scientific Research*, **7**(11), 84–89.
- Vannote R.L., Minshall G.W., Cummins K.W., Sedell J.R. and Cushing C.E. (1980), The River Continuum Concept, *Canadian Journal of Fisheries and Aquatic Sciences*, **37**, 130–137.
- Viongradov A.K., Bogatova Y.I. and Synegub I.A. (2018), Ecology of marine ports of the black and Azov Sea Basin, *Springer International Publishing*, **417**, 298–309. doi:10.1007/978-3-31963062-5.
- Wan Maznah W.O. (2010), Perspectives on the use of algae as biological indicators for monitoring and protecting aquatic environments, with special reference to Malaysian freshwater ecosystems, *Tropical Life Sciences Research*, **21**(2), 51–67.
- Wang B., Ming X., Xia S., Qinsheng W. and Xuelei Z. (2016), Does reduced sediment load contribute to increased outbreaks of harmful algal blooms off the Changjiang Estuary. *Acta Oceanologica Sinica*, **35**(8), 16–21. Doi:10.1007/s13131-016-0846-5.
- Wehr J.D. and Sheath R.G. (2003), *Freshwater algae of North America. Ecology and classification*. Academic Press, an imprint of Elsevier science. Amsterdam Boston, London: New York Oxford, Paris San Diego San Francisco Singapore Sydney Tokyo.
- Wehr J.D., Sheath R.G. and Kociolek P.J. (2015). *Freshwater Algae of North America, Ecology and Classification*. Academic Press. ISBN: 978-0-12-385876-4.
- Withford L.A and Schumacher G.J. (1973). *A manual of freshwater algae*. Sparks press Raleigh, N.C.
- Wokoma O.A.F., Nwaugo V.O., Chima G.N. (2016), Epipelagic algae species composition and abundance in the brackish water axis of Sombreiro River, Niger Delta, Nigeria, *IIARD International Journal of Geography and Environmental Management*, **2**(1).
- Wu N., Tang T., Qu X. and Cai Q. (2009). Spatial distribution of Benthic algae in the Gangqu River, Shangila, China. *Aquatic Ecology*, **43**, 37–49. Doi: 10.1007/s.10452-007-9153-8.