

Assessment of observed temperature trend patterns of Bhubaneswar city, India with special prominence on future projections using SimCLIM climate model and farmer's perception

Punya Murthy Khristodas^{1*}, K. Palanivelu¹, A. Ramachandran¹ and B. Anjan Kumar Prusty²

¹Centre for Climate Change and Disaster Management, Anna University, Chennai, Tamil Nadu, India


²Department of Environment Studies, Berhampur University, Berhampur-760007, Odisha, India

Received: 11/03/2023, Accepted: 03/10/2023, Available online: 13/01/2024

*to whom all correspondence should be addressed: e-mail: punya.khristodas@gmail.com

<https://doi.org/10.30955/gnj.004859>

Graphical abstract



Abstract

Temperature dynamics is a widely recognized indicator of the global warming phenomenon. Changes in temperature patterns in Bhubaneswar, India, were assessed by examining the monthly minimum and maximum temperature data for 60 years (1956–2015) sourced from the Indian Meteorological Department. SimCLIM climate change risk assessment software was used for projecting the temperature regime for four different representative concentration pathways. Further, a survey of 112 farmers was conducted to understand their perceptions of temperature variations in and around Bhubaneswar city using a multi-stage sampling technique. Mann–Kendall statistics and linear regression were used to analyze the monthly temperature data and trend detection. The study reveals a change of +4%, -4.44%, and +1.09% in the mean monthly maximum, minimum, and annual temperature. The results of the future projection show a temperature change of 0.81°C for RCP 2.6, 1.12°C for RCP 4.5, 1.03°C for RCP 6.0, and 1.54°C for RCP 8.5 for the year 2050. Confirming the analysis findings, most of the interviewed farmers also perceived increasing temperatures and decreasing precipitation in and around the city. The study outcome of temperature trend analysis and future projections will be helpful for farmers and policymakers in formulating adaptation strategies to climate change.

Keywords: Climate change; multistage sampling technique; future projection; Mann–Kendall; SimCLIM; temperature variations

1. Introduction

Anthropogenic activities contributing to recent climate changes have significantly impacted ecosystems, natural resources, and humans. According to the AR6 IPCC report (IPCC, 2021), under very high GHG emissions, the estimated warming will be 2.4°C (2041–2060) and 4.4°C (2081–2100). Furthermore, the very likely range will be 3.3–5.7°C from 2081 to 2100. The average global temperature estimates based on linear interpolations illustrate that the global temperature has risen by 0.85°C from 1880 to 2012 (IPCC, 2014). The degree to which urbanization affects climate at a regional scale has not yet been determined.

Nevertheless, the urban heat islands and their impacts on life support systems have remained a topic of debate in many scientific forums across the globe. Agriculture is the most vulnerable industry to climate change due to its enormous size and susceptibility to weather variables, resulting in significant economic consequences (Mendelsohn, 2009). Over the past few decades, global warming has threatened crop productivity (Hussain *et al.*, 2021). Climate change is an extreme natural hazard faced by humankind, and the changes have become abrupt and devastating in the recent past, aggravated by natural factors and unconstrained anthropogenic activities (Prusty, 2011). Small-scale subsistence farmers are the most affected by climate change and its consequences because of their low levels of adaptive capability and dependence on rain-fed agriculture (Ifejika *et al.*, 2010; Hillel *et al.*, 2011). The Intergovernmental Panel on Climate Change (IPCC), in early 1990, reported that the highest impact of climate change would be reflected in the unseen levels of climate migrants. The rising interest in vulnerability indices is part of a broader effort to measure the degree of vulnerability of individual countries to natural hazards.

In India, notable variations in natural temperature patterns have been observed, and frequent and devastating disasters have become common in recent times. Analysis of temperature data from weather stations across India illustrates a notable rise in temperature over the last century (Dash *et al.*, 2009). Kumar *et al.* (2006) predicted a temperature increase of 3–4°C toward the end of this century, with the pressure projected to be higher in the northern states of India. According to a World Bank report (2013), India faces a warmer climate with repeated spells of hot weather occurring at regular intervals. Furthermore, the report predicts a 4°C rise in temperature. It is projected that some regions of India might shift to high-temperature climatic systems, influencing agriculture.

Climate change affects many sectors of the economy, but agriculture has been the worst. Climate change significantly affects the poor's living situations because they cannot adapt; furthermore, their livelihoods are heavily based on agriculture, which is very susceptible to climate change (Ali *et al.*, 2019). Agriculture contributes to the global economy, creating employment opportunities for billions of people (Allahyari *et al.*, 2016). Agricultural activities in India contribute about 17% to the national GDP and employ over 60% of the population (Arjun, 2013). Thus, climate change-induced disasters such as drought and extreme temperatures are the costliest in India regarding the loss of natural resources and property, leading to overall economic decline. India experienced 41 droughts between 1801 and 2009 (Attri & Tyagi, 2010). Agriculture and its allied sectors are highly vulnerable to climate change; temperature and rainfall directly affect farm production (Thorton *et al.*, 2014). These two phenomena also directly affect soil erosion, decreasing water supply, groundwater level, and irrigation (IPCC, 2001). More significant changes in the rainfall patterns, less diversified agriculture, and increasing frequency of droughts and floods directly affect food production and supply. These issues would seriously affect farmers and other stakeholders. Thus, a better understanding of (i) farmers' perceptions of temperature trends and variation, (ii) their current adaptation measures, and (iii) other allied factors is essential for policymaking and facilitating sustainable agricultural practices for successful mitigation of climate change and to develop adaptation strategies in this sector (Jin *et al.*, 2015).

To predict the climate of the 21st century and beyond, an estimate of future changes in the forcing is essential (Gosse *et al.*, 2010). The global mean surface temperatures are predicted to be 0.3–1.7°C, 1.1–2.6°C, 1.4–3.1°C, and 2.6–4.8°C under four representative concentration pathways (RCP), i.e., RCP 2.6, RCP 4.5, RCP 6.0, and RCP 8.5, respectively, for the period between 2081 and 2100 (IPCC, 2013). Global circulation model (GCM) projections have provided warnings about the occurrence and severity of these temperature variations in the forthcoming years (Amin *et al.*, 2016). Considering GCM projections, better adaptation strategies must be developed to withstand climate hazards such as heat waves, droughts, and cyclones. Many cities in India and beyond in the

subcontinent have been affected by the fury of such risks. Bhubaneswar, a city in the eastern coastal state of Odisha in India, has also experienced its shares of severe climatic hazards, such as the Super Cyclone (1999) and Phailin Cyclone (2013), and Hudhud Cyclone (2014). These extreme events could be assessed by employing integrated models and tools. SimCLIM is an "open-framework" piece of software that can be used to study climate change impacts over time and space (Figure 4). The SimCLIM risk assessment software tools can be used for site-specific climatic data analysis (Bao *et al.*, 2015; Ramachandran *et al.*, 2017).

The United Nations Development Programme (UNDP) 2014 cautioned that climatic variations would profoundly impact small-scale and marginal farmers in developing countries. Bhubaneswar, a second-tier city in India, as classified by the Fifth Central Pay Commission of India in 1997, offers an opportunity to explore the issue along the above mentioned lines. Several researchers have examined temperature change and its variation in Odisha and Bhubaneswar city to understand the severe natural events. Panda and Sahu (2019) reviewed the seasonal rainfall and temperature change in two districts of western Odisha (i.e., Balangir and Kalahandi) and one district of southern Odisha (i.e., Koraput). Their study reveals that annual minimum and maximum temperatures increased during 1980–2017. Gogoi *et al.* (2019) studied the Land Use Cover Change (LULC) impacts on surface temperature and found that the mean temperature has increased significantly in Odisha.

The temperature trend analysis of Bhubaneswar city was studied in some previous works (Majhi & Rath, 2018; Dash, 2016). However, no research has been conducted into future temperature trends and farmer views. What are local farmers' perceptions of climatic trends, how well do these perceptions match meteorological records, and what would be the future temperature trends based on various RCP scenarios? These questions must be answered to understand farmers' perceptions of climate change better and provide empirical evidence for policies aimed at improving farmers' adaptive capacity by improving their ability to perceive climate change correctly. As a result, the present study aims for a better understanding of recent changes and variability in temperature, future temperature trends, and the farmers' perception of climate change in Bhubaneswar city with three distinct objectives, i.e. (i) to investigate the trends of monthly maximum, minimum, and mean temperatures of Bhubaneswar, (ii) to project future temperature trends at the local level under different RCP scenarios of Bhubaneswar city (iii) to identify the farmers' perceptions on climate change and temperature variability. Thus, the present study is the first of its kind from Bhubaneswar.

2. Study area

Bhubaneswar city (20° 14' 0" N, 85° 50' 0" E, Figure 1) is situated on the western side of the "Mahanadi Delta" on the banks of River Kuakhai and at the southwest of Cuttack city in the state of Odisha, India. River Daya, which branches off River Kuakhai, flows along the southeastern

part of the city. It was one of the earliest planned cities of India, after Chandigarh and Jamshedpur, designed by German architect Otto Konigsberger in 1946. Historically, the city dates back to 300 BC and has a geographic area of 135 km² (Figure 1). It is located at an average altitude of 45 m above the mean sea level and is close to the sea; the weather remains hot and humid all year round. UNDP (2014) reveals that the city experienced a maximum temperature of 47°C in May 2013. The maximum atmospheric temperature is more than 38°C during summer. The city received 1406.1–1958.91 mm of rainfall during the study period of 1991–2012, with maximum precipitation occurring in July and August (Dash & Chakraborty, 2016). The relative humidity of the city is very high (around 80%) from July to October. Recently, with the upsurge in economic activities and associated requirements of fossil fuel-based energy sources, a considerable increase in greenhouse gas emissions has been reported in the city (Jena, 2018). The Suspended Particulate Matter (SPM) in the ambient air in the city ranged from 156.7µg/m³ to 280.6 µg/m³ from February 2013 through January 2014 (Mohapatra & Biswal, 2014). The Ministry of Road Transport and Highways (MoRTH, 2016) reveals that the number of vehicles registered in the city is 1.08 Million. Different activities in Bhubaneswar contribute to 0.97 million tons of CO₂ equivalent (TeCO₂) annually, whereas per capita emissions were 0.84 T/year in 2007–08. Bhubaneswar city, with a population of 837,321 per the 2011 Census of India, has a population density of 2131.4/km². The spatial growth of Bhubaneswar was 176.7% from 1961 to 1971 is the highest in India (Naik, 2013). The city has become a socio-economic hub and cultural centre for many people from Odisha and beyond.




Figure 1. Map of Bhubaneswar city

3. Methods

Following Hou *et al.* (2015) and Chombu *et al.* (2020), the present study examined the observed temperature using

statistical techniques and farmers' perceptions of climate change. To develop policy recommendations for future adaptation plans, it is necessary to assess changes in climate characteristics based on past scientific records and the perceptions of the local community (Mubangizi *et al.*, 2017). In addition, the study attempts to forecast future temperature trends. Advances in weather and climate forecasting offer opportunities for proactive risk management at the local level (Fao, 2014).

3.1. Statistical analysis

To examine the historically observed temperature trend, monthly temperature data were obtained from the Indian Meteorological Department (IMD) for 60 years (1956–2015). For ease of understanding and interpretation, the datasets were split into the whole study period (1956–2015) and two phases: 1956–1985 (First Phase) and 1986–2015 (Second Phase). Monthly mean maximum, mean minimum, and mean temperature along with standard deviation (SD) were computed for the whole study period following Jeganathan *et al.* (2013), and the dataset was subjected to the Mann–Kendall Trend (MKT) test to assess the significance in such variability. Most studies have used the non-parametric MKT test to find trends in hydro-meteorological observations that do not follow a normal distribution of data points (Girma *et al.*, 2020). Trend detection and analysis were performed using linear regression (LR) models. Trend detection and analysis were performed through statistical tools LR and MKT test.

3.2. Climate model- SimCLIM

SimCLIM desktop version 4.0 was used with 40 GCMs for temperature scenario generation of Bhubaneswar city for 2050–2100. The "site-specific temperature scenario" tool was used to obtain the local temperature projection. Geographical coordinates of the area of interest given along with the locally observed temperature trend were obtained from IMD, Pune. Further, all the 40 GCMs were organized hierarchically by pattern scaling to perform sensitivity analysis. The central tendency was measured based on the median value of constructed ensembles, i.e., 40 selected GCM patterns finalized for the study area. Analysis of the weather data (1956–2015) record helped generate local-level baseline data compared to SimCLIM. In a changing climate, the SimCLIM modeling system can be used to examine the implications and dangers of climatic extremes (Warrick, 2009; Amin *et al.*, 2016).

3.3. Farmer's perception of climate change

A multi-stage and simple random sampling method following Ndambiri *et al.* (2014) was applied to identify and select the villages around Bhubaneswar city and analyze the farmers' perceptions of climatic change. Finally, a sample of 112 farmers, i.e., 16 from Nuagan, Paikarapur, Madanpur, Durgapur, Padasahi, Sundarapada, and Bainchuan on the periphery of Bhubaneswar city, was selected randomly. The present survey was carried out using a questionnaire customized for the purpose. The survey included a pre-tested structured questionnaire format following Kabir *et al.* (2016) that explored residents' perceptions of temperature change, the frequency and

magnitude of climate-related hazards in the last 30 years and their livelihood impacts. It helped obtain information on the socio-demographic condition of the farmers. It included questions about temperature change perceptions and the related effect on different aspects of the source of their income. Based on the landholding size, the sample households were classified into five categories, viz., (i) no land, (ii) owning ≤ 1 acre, (iii) owning 1–2.5 acres, (iv) owning 2.5–5 acres, and (v) owning 5–10 acres. With the help of XLSTAT statistical software, descriptive statistical tests following Jha and Gupta (2021), which included mean, percentages, and frequency counts, were conducted to summarize the socioeconomic conditions of farmers and their perceptions of temperature change and variability. A frequency table was used for statistical analysis to record the reason behind different climatic constraints. The data analysis included the application of correlation and regression, percent analysis, and a composite index developed in the study.

4. Results and discussions

4.1. Observed temperature trend analysis

A monthly, seasonal, and annual temperature trend significance test was performed using the Mann–Kendall non-parametric statistical test for 60 years (1956–2015) for Bhubaneswar city (Table 1). The change in maximum, minimum, and mean temperatures for the period 1956–2015 were $1.64 \pm 0.21^\circ\text{C}$, $1.10 \pm 0.14^\circ\text{C}$, and $0.85 \pm 0.10^\circ\text{C}$, respectively (Figure 2). According to a previous study by Majhi and Ratha (2018), the mean monthly maximum temperature in Bhubaneswar city increased faster than the average and minimum temperatures from 1970 to 2017. For the monthly maximum temperature, positive trends were observed except for June. Similarly, a positive trend was observed for the whole study period except for May for the monthly minimum temperature. The negative trends in maximum and minimum temperatures for May and June may be attributed to higher atmospheric humidity, as relative humidity is inversely related to air temperature. The average relative humidity in Bhubaneswar is roughly 77%, ranging from 68% in winter (February) to 88% during the monsoon season (July). Moreover, UNDP (2014) indicated that between 1959 and 2009, cyclonic storms and depressions occurred nine times between May and June, which may be one of the reasons for the negative temperature trends in Bhubaneswar City during these months.

Furthermore, the city receives 26 mm of rain in April, 67 mm in May, and 209 mm in June (CSE, 2020). This significant shift in rainfall pattern compared to April may be one of the reasons for the negative trends. The linear temperature trends exhibit a positive change over the whole study period (1956–2015) (Figure 3). The first phase (1956–1985) witnessed an increasing mean temperature ($a=0.009$, $R^2=0.038$). Similarly, the second phase (1986–2015) also experienced an increasing mean annual temperature ($a=0.025$, $R^2=0.353$). The temperature pattern of the city showed positive trends for mean maximum temperature, mean minimum temperature, and

mean annual temperature for the whole study period (Figure 4). The mean maximum temperature pattern ($a=0.017$, $R^2=0.334$) and the mean minimum temperature anomaly ($a=0.007$, $R^2=0.084$) revealed a significantly increasing trend. Similar to the above two variables, the mean annual temperature anomaly also exhibited an increasing trend ($a=0.012$, $R^2=0.26$). A positive trend in the observed temperatures was found in the mean maximum, mean minimum, and mean annual temperatures over the whole study period. The trend lines indicate that the mean annual temperature over Bhubaneswar city has increased by about 0.59°C (60-year mean value).

Table 1. Mann-Kendall Test results

Month	Maximum Temperature	Sen's Slope	Minimum Temperature	Sen's Slope
January	0.27*	0.02	0.06	0.01
February	0.18*	0.02	0.02*	0.00
March	0.22*	0.02	0.19*	0.01
April	0.06	0.01	0.00*	0.00
May	0.04*	0.00	-0.05*	0.00
June	0.06	0.01	0.16*	0.01
July	0.27*	0.02	0.19*	0.01
August	0.36*	0.02	0.21*	0.01
September	0.32*	0.02	0.27*	0.01
October	0.18*	0.01	0.12*	0.01
November	0.24*	0.02	0.18*	0.02
December	0.33*	0.03	0.08	0.01
JF	0.31*	0.02	0.06	0.00
MAM	0.19*	0.01	0.17*	0.01
JJAS	0.26*	0.02	0.26*	0.01
OND	0.23*	0.02	0.13*	0.01
Annual	0.39*	0.02	0.21*	0.01

*significant at 0.1 level




Figure 2. Cluster column chart showing monthly temperature mean and standard error

A previous study by Jeganathan *et al.* (2013) showed that Chennai's mean annual temperature trend increased by about 1.3°C (60-year mean value) from 1951 to 2010. Kolkata, Mumbai, and Chennai had an annual mean temperature of 1°C , 0.6°C , and 0.56°C , respectively Dhorde *et al.* (2009). The highest recorded temperature for the city was 40°C in May 1957, and the lowest minimum temperature was 12.1°C in December 1970. Further, the warmer years on record are in the order 2010 (0.9°C) > 1979 (0.7°C) > 2006 (0.6°C) > 2012 (0.6°C) > 2005 (0.5°C) > 2002 (0.4°C) > 2003 (0.3°C) > 1980 (0.4°C) > 1998 (0.4°C) > (0.3 $^\circ\text{C}$), which depicts the long-term variation in the average temperature for the period 1956–2015. Similarly,

the highest mean minimum temperature was observed to be $32.05^{\circ}\text{C} \pm 0.83$ for May. The study reveals a change of +4%, -4.44%, and +1.09% in the mean monthly maximum, minimum, and annual temperature.

century. The findings also agree with an earlier study by Kumar *et al.* (2011), whose results projected a temperature increase of 1–4° C for the year 2050 and an increase of more than 4.5°C for the year 2100.




Figure 3. Linear temperature trends of Bhubaneswar City 1956-2015

4.2. Projection of mean temperature for 2050 and 2100

Validating the SimCLIM climate model with the observed data obtained from IMD have conferred that the model has shown a reasonable skill in simulating temperature patterns for Bhubaneswar city. The magnitudes and frequency characteristics of the surface temperature variability on annual time scales agreed. Hence, this study attempts to assess various aspects of future projections in Bhubaneswar city using SimCLIM. The mean temperature scenarios of climate change for Bhubaneswar city revealed positive trends for different RCPs for the time scale from 1995 to 2100 (Figure 5). The results obtained from SimCLIM show a temperature change of 0.81°C, 1.12°C, 1.03°C, and 1.54°C for RCP 2.6, RCP 4.5, RCP 6.0, and RCP 8.5 emission scenarios, respectively, under medium projection for the year 2050. Similarly, the results revealed a temperature change of 0.80°C, 1.48°C, 2.08°C, and 3.68°C for RCP 2.6, RCP 4.5, RCP 6.0, and RCP 8.5, respectively, under medium projection for the year 2100. Basha *et al.* (2017) project a temperature increase of 3.2°C in India by the end of the




Figure 4. Maximum, minimum and mean annual temperature anomalies of Bhubaneswar during 1956-2015

Raghavan *et al.* (2020) also reported that under the RCP8.5 scenario, these temperatures are expected to rise by about 4.7°C and 5.5°C by the end of the twenty-first century, respectively, compared to recent temperatures (1976–2005 average). The percentage deviation in mean temperature from base temperature was 2.95% for RCP 2.6, 4.04% for RCP 4.5, 3.75% for RCP 6.0, and 5.56 % for RCP 8.5 emission scenarios for the projection year 2050, and 0.04% for RCP 2.6, 1.22% for RCP 4.5, 3.64% for RCP 6.0, and 7.26% for RCP 8.5 emission scenarios for the projection year 2100.

4.3. Farmer's perceptions of temperature change

Of the interviewees, 50.90% were in the 50+ age group, and 43.63% of the farmers had more than 21 years of farming experience. According to Amadou *et al.* (2015), the older the farmer is, the more experience they have with farming practices. Correct perception of climatic variability positively correlates with age and experience (Rapholo & Makia, 2020). Among them, 55.45% had received an education below the 10th standard. A maximum number of interviewees could be placed in the marginal farming category, and 67.27% practiced rain-fed farming. According to Sathyan *et al.* (2018), millions of rainfed smallholder

farmers would face immediate hardship and hunger due to climate change. They will be less able to make appropriate decisions about when to sow, what to grow, and how to time inputs (NICI, 2009) and have a low ability to respond (Burnham & Ma, 2017). Almost 90% of the farmers work as daily laborers during the off-seasons (Table 2). Among those surveyed, 60% of the farmers have a landholding of 1–2.5 acres, and 90% have an annual income of < 10.8 Lakhs (2056 USD). Almost every interviewee opined that the temperature had increased in the last 20–30 years; 77.71% of respondents strongly agreed and 22.32% agreed that the temperature was rising.

Table 2. Social and economic background of the interviewee in Bhubaneswar City

Variable	Interviewee (%)
<i>Age of farmers (Years)</i>	
Below 30	6.36%
Between 30-40	20.90%
Between 40-50	21.84%
Above 50	50.90%
<i>Farming Experience (Years)</i>	
1-5	1.81%
6-10	10%
11-15	12.72%
16-20	31.84%
21 and above	43.63%
<i>Qualifications of farmers</i>	
Below 10 th	55.45%
10 th	30.92%
+2	4.54%
Graduate	9.09%
<i>Size of landholding under operation (Acres)</i>	
Between 1-2.5	60%
Between 2.5-5	26.36%
Below 1 acre	7.29%
No land	0.90%
Between 5-10	5.45%
<i>Annual income (Lakhs)</i>	
Below 1.5 (1810.01 USD)	90%
1.5-3 (1810.01 USD- 3620.02 USD)	10%
<i>Type of crop</i>	
Rabi	76.36%
Rabi and Kharif	23.64%
<i>Source of water</i>	
Rainfed and Irrigated	32.73%
Rainfed	67.27%

Similarly, around 75% of the farmers strongly agreed on the rainfall pattern, and 25% agreed that the rainfall decreased. About 68.75% of the interviewees strongly agreed to a decrease in the number of cool days, 16.96% agreed, and 14.29% agreed moderately. Further, a majority of the interviewees believed that hot days are increasing; among them, 68.75% of the farmers strongly agreed, and 31.25% only agreed that the number of hot days is increasing during the summer and rainy seasons (Figure 6). Regarding their perception of climate change hazards, 50% of the farmers indicated that climate change is anthropogenic, 6.25% thought it was natural, and 43.75% believed that the temperature change is a result of both

anthropogenic and natural factors (Figure 7). The majority of the farmers (89%) indicated that the Super Cyclone (1999) was the most powerful tropical cyclone in the last two decades that significantly impacted their livelihoods. Atmospheric temperature, water vapor, and ice concentration influence the sea surface temperature (SST) (Carvalho & Wang, 2020). However, cyclonic events such as Phailin (2013), Hudhud (2014), and Fani (2019) have had considerably less impact on their agricultural production. Cyclones have had a physical, economic, and psychological effect on people due to the repeated loss of farms, livelihoods, and assets (Patel, 2018). Around 62.50% of the interviewees opined that urbanization, deforestation, and cyclones are the main reasons for uprooting and damaging trees, leading to temperature increase in the city.




Figure 5. SimCLIM total trend of mean temperature projection at Lat. 20.296/Long. 85.824 for different RCPs




Figure 6. Farmers' perception of climate change and farming losses

In comparison, 18.75% said only tree loss due to the cyclone, 11.46% only deforestation and 7.29% said the city's rapid growth, mainly in the last ten years, was the temperature change. With the growing population and urbanization trend, deforestation has increased, disrupting natural atmospheric and climatic patterns and amplifying the destructive effects of natural disasters (Ali *et al.*, 2014). According to Sahoo *et al.* (2020), in Bhubaneswar, around 5% of trees were damaged entirely, 6% were uprooted,

31% had significant damage, and 57% had partial damage from storm Fani in 2019. Storms and cyclones have exacerbated the loss of current green cover (Sahoo & Bhaskaran, 2016, 2017). The city has experienced the uprooting of millions of trees due to cyclones. Hurricane damage can reduce a forest's ability to absorb carbon dioxide from the atmosphere, contributing to global warming (Tulane University, 2021). Around 66.66% of the interviewees opined that the drought was infrequent, whereas 33.34% indicated it was rare.

According to most interviewees (71.88%), floods happen rarely, and 28.12% stated that floods occur very infrequently. According to Mishra *et al.* (2020), the city witnessed urban flooding in 2008, 2014, 2016, 2018, and 2019. Most interviewees (62.50%) felt that cyclones frequently occurred, whereas 37.50% thought it often.




Figure 7. Farmer's perception of climate change hazards

5. Conclusions

The present study, the first of its kind, focused on Bhubaneswar city, India, and integrates the temperature trend projections using the SimCLIM model and farmers' perceptions. The Mann–Kendall and Sen's Slope estimator test results exhibited positive temperature trends. A notable conclusion from the farmers' perception study is that their responses regarding the temperature variations agree with the meteorological data, which is a testament to their experiential knowledge. These changes in the observed and projected temperature trends reveal that Bhubaneswar city may expose to significant climate change-induced risks of rising temperatures. The present study results will be an essential source of information for environmental managers and policymakers to frame adaptation strategies, particularly for Bhubaneswar city. To some extent, highly educated farmers had additional knowledge of good farming practices and coping and adaptation strategies for climate change-induced hazards like cyclones and floods. The government has to support the farmers in adaptation strategies such as efficient usage of water resources in agriculture since the rain has become more unpredictable and unseasonal. Through policymakers and intellectuals, the government needs to expedite the improvement and dissemination of agricultural technologies such as irrigation schemes and the usage of climate-resilient crops.

Acknowledgments

The financial support as under the Maulana Azad National Fellowship (MANF) scheme 2016–2021 (Reference no.: F1-17.1/2014-15/MANF-2014-1CHR-ORI-42755, February 2017) of the University Grants Commission (UGC), Government of India, awarded to Mr. Punya Murthy Khristodas is gratefully acknowledged. We express our gratitude to our colleagues at the Centre for Climate Change and Disaster Management (CCC&DM), Anna University, Chennai.

References

- Ali A., Riaz S. and Iqbal S. (2014). Deforestation and Its Impacts on Climate Change an Overview of Pakistan, *Papers on Global Change*, **21(1)**, 51–60.
- Ali F M., Ashfaq M., Hasan S. and Ullah R. (2019). Assessing Indigenous Knowledge through Farmers' Perception and Adaptation to Climate Change in Pakistan, *Polish Journal of Environmental Studies*, **29(1)**, 525-532.
- Allahyari M S., Ghavami S., Masuleh Z D., Michailidis A., and Nastis S A. (2016), Understanding Farmers' Perceptions and Adaptations to Precipitation and Temperature Variability: Evidence from Northern Iran, *Climate*, **4(4)**, 58.
- Amadou M L., Villamor V G B., Attua E M. and Traore S B. (2015), Comparing farmers' perception of climate change and variability with historical climate data in the Upper East region of Ghana, *Ghana Journal of Geography*, **7(1)**, 47 –74.
- Amin A., Nasim W., Mubeen M., Sarwar S., Urich P., Ahmad A., Wajid A., Khaliq A., Rasul F., Hammad H M., Rehmani M I A., Mubarak H., Mirza N., Wahid A., Ahamd S., Fahad S., Ullah A., Khan M N., Ameen A A., Shahzad B., Saud S., Alharby H., Ata-ul-karim S T., Adnan M., Islam, F., Ali, Q.S. (2016). Regional climate assessment of precipitation and temperature in Southern Punjab (Pakistan) using SimCLIM climate model for different temporal scales, *Theoretical Applied Climatology*, **131**, 121–131.
- Arjun K M. (2013), Indian Agriculture- Status, Importance, and Role in Indian Economy, *International Journal of Agriculture and Food Science Technology*, **4(4)**, 343-346.
- Attri S D. and Tyagi A. (2010), Climate profile of India, Contribution to the Indian Network of Climate Change Assessment. (National Communication-II), Ministry of Environment and Forests. Climate profile of India, Contribution to the Indian Network of Climate Change Assessment, *Indian Meteorological Department*, pp 1–129.
- Bao Y., Hoogenboom G., McClendon R. and Urich P. (2015), Soybean production in 2025 and 2050 in the southeastern USA based on the SimCLIM and the CSM-CROPGRO-soybean models, *Climate Research*, **63**, 73–89.
- Basha G., Kishore P., Ratnam M V., Jayaraman A., Kouchak A A., Ouarda T B M J., and Velicogna I. (2017), Historical and Projected Surface Temperature over India during the 20th and 21st century, *Scientific Report*, **7**, 2987. <https://doi.org/10.1038/s41598-017-02130-3>
- Burnham M., and Ma Z. (2017), Climate change adaptation: Factors influencing Chinese smallholder farmers' perceived self-efficacy and adaptation intent, *Regional Environmental Change*, **17**, 171–186.
- Carvalho S K., and Wang S. (2020), Sea surface temperature variability in the Arctic Ocean and its marginal seas in a changing climate: Patterns and mechanisms, *Global and Planetary Change*, **193**, 103265.

- Chombo O., Lwasa S., and Tenywa M. (2020), Spatial and Temporal Variation in Climate Trends in the Kyoga Plains of Uganda: Analysis of Meteorological Data and Farmers' Perception, *Journal of Geoscience and Environment Protection*, **8**, 46-71.
- CSE. (2020). Road map for implementation of water-sensitive urban design and planning in Odisha, 56p.
- Dash A. (2016). Analytical Study of Rainfall Probability at Bhubaneswar, *International Journal of Applied and Pure Science and Agriculture*, **2(6)**, 151-155.
- Dash M., and Chakraborty M. (2016). Bio Climatic Analysis of Bhubaneswar-An Investigation to Arrive Human Comfort through Natural Ventilation, *International Journal of Scientific and Research Publications*, **6(5)**, 557-561.
- Dash S K., Kulkarni M A., Mohanty U C. and Prasad K. (2009), Changes in the characteristics of rain events in India, *Journal of Geophysical Research; Atmospheres*, **114**, 1-12.
- Dhorde A., and Gadgil A S. (2009). Long-term temperature trends at the four largest cities of India during the twentieth century, *Journal of Indian Geophysical Union*, **13(2)**, 85-97.
- Emanuel K A. (1986). An air-sea interaction theory of tropical cyclones. Part I: Steady-state maintenance, *Journal of Atmospheric Science*, **43**, 585-604.
- Emanuel K A. (1988), The maximum intensity of hurricanes, *Journal of Atmospheric Science*, **45**, 1143-1155.
- FAO. (2014). Managing climate risks and adapting to climate change in the agriculture sector in Nepal, Food and Agriculture Organization of the United Nations (FAO) Rome, Italy, 143p.
- Girma A., Qin T., Wang H., Yan D., Gedefaw M., Abiyu A., Batsuren D. (2020). Study on Recent Trends of Climate Variability Using Innovative Trend Analysis: The Case of the upper Huai River Basin, *Polish Journal of Environmental Studies*, **29(3)**, 2199-2210.
- Gogoi P P., Vinoj V., Swain D., Roberts G., Dash J., and Tripathy S. (2019). Land use and land cover change effect on surface temperature over Eastern India, *Scientific Reports* <https://doi.org/10.1038/s41598-019-4213-z>
- Goosse H., Barriat P Y., Lefebvre W., Loutre M F., and Zunz V. (2010). Introduction to climate dynamics and climate modeling, (<http://www.climate.be/textbook>).
- Hillel D., and Rosenzweig C., (2011), Handbook of Climate Change and Agro ecosystems: Impacts, Adaptation, and Mitigation, Imperial College Press, London, 437p.
- Holland G J. (1997), The maximum potential intensity of tropical cyclones, *Journal of Atmospheric Science*, **54**, 2519-2541.
- Hou L., Huang J., and Wang J. (2015). Farmers' perceptions of climate change in China: The influence of social networks and farm assets, *Climate Research*, **63(3)**, 191-201.
- Hussain J., Khaliq T., Rahman M.H., Ullah A., Ahmed I., Srivastava A K., Gaiser T., Ahmad A. (2021). Effect of Temperature on Sowing Dates of Wheat under Arid and Semi-Arid Climatic Regions and Impact Quantification of Climate Change through Mechanistic Modeling with Evidence from Field, *Atmosphere*, **12(927)**, 1-15.
- Ifejika S C., Kiteme B., Ambenje P., Wiesmann U., Makali S. (2010), Indigenous knowledge related to climate variability and change: insights from droughts in semi-arid areas of former Makeni District, Kenya, *Climatic Change*, **100(2)**, 295-315.
- IPCC. (2001). Third Assessment Report, *Climate Change*, Cambridge University Press.
- IPCC. (2013). Climate Change: The Physical Science Basis. *Working Group I Contribution to the IPCC Fifth Assessment Report*, Stocker.
- IPCC. (2014) Climate Change: Synthesis Report. Approved Summary for Policymakers. Cambridge, UK, Geneva, Switzerland.
- IPCC. (2021). Summary for Policymakers, In: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., Zhou, B. (eds.) *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge.
- Jeganathan A., and Andimuthu, R. (2013). Temperature trends of Chennai City, India, *Theoretical Applied Climatology*, **111(3-4)**, 417-425.
- Jena P P. (2018). Urban Growth and Climate Change- A Study in the Context of the Bhubaneswar City of Odisha, *Journal of Humanities and Social Science*, **23(4)**, 49-55.
- Jha C. K., and Gupta V. (2021), Farmer's perception and factors determining the adaptation decisions to cope with climate change: An evidence from rural India, *Environmental and Sustainability Indicators*, **10**: 100112. <https://doi.org/10.1016/j.indic.2021.100112>.
- Jin J J., Gao Y W., Wang X M., Nam P K. (2015) Farmers' risk preferences and their climate change adaptation strategies in the Yongqiao District, China, *Land Use Policy*, **47**, 365-372.
- Kabir M I., Rahman M B., Smith W. (2016). Knowledge and perception about climate change and human health: findings from a baseline survey among vulnerable communities in Bangladesh, *BMC Public Health*, **16(1)**, 1-10.
- Kumar K R., Sahai A K., Kumar K K., Patwardhan S K., Mishra P K., Revadkar J V., Kamala K., Pant G B. (2006), High-resolution climate change scenarios for India for the 21st century, *Current Science*, **90(3)**, 334-345.
- Kumar S N., Aggarwal P K., Rani S., Jain S., Saxenas R., Chauhan N. (2011). Impact of climate change on crop productivity in Western Ghats, coastal and northeastern regions of India, *Current Science*, **101(3)**, 332-341.
- Majhi B., and Rath K C. (2018). Changing trends of temperature in Bhubaneswar: A city in eastern India, *International Journal of Geology, Earth & Environmental Sciences*, **8(1)**, 5-14.
- Malkus J S., and Riehl H. (1960). On the dynamics an energy transformation in steady state hurricanes, *Tellus*, **12**, 1-20.
- Mendelsohn R. (2009). The impact of climate change on agriculture in developing countries *Journal of Natural Resources Policy Research*, **1(1)**, 5-19.
- Miller B I. (1958). On the maximum intensity of hurricanes, *Journal of Atmospheric Science*, **15**, 184-195.
- Mishra S P., Kumar C S., and Mohammed, S. (2020), Emerging Threats during Anthropocene as Urban Flooding of Bhubaneswar City, India, *Water and energy international (Water resource section)*, pp 46-58.
- Mohapatra K., and Biswal S K. (2014), Assessment of Ambient Air Quality (AAQ) index in Bhubaneswar City of Odisha, *International Journal of Advanced Research in Science and Engineering*, **3(6)**, 190-196.
- Mubangizi N., Kyazze F B., Mukwaya P I. (2017), Smallholder Farmers' Perception and Adaptation to Rainfall Variability in

- Mt. Elgon Region, Eastern Uganda, *International Journal of Agricultural Extension*, **5**, 103-117.
- Naik M. (2013). Urban Sprawl of Bhubaneswar City using GIS Applications and Entropy. (Master's Thesis, National Institute of Technology, Rourkela, India), Retrieved from <https://www.mobt3ath.com/uplode/book/book-43940.pdf>.
- Ndambiri H K., Ritho C N., Mbogoh S G. (2014), An evaluation of farmer's perception and adaptation to the effects of climate change in Kenya, *International Journal of Food and Agricultural Economics*, **1**, 75-96.
- NICI. (2009). Impact of Climate Change to 2030 -A Commissioned Research Report; National Intelligence Council: New Delhi, India.
- Panda A. and Sahu N. (2019). Trend analysis of seasonal rainfall and temperature pattern in Kalahandi, Bolangir and Koraput districts of Odisha, India, *Atmospheric Science Letter*, **20**, 1–10.
- Patel S K. (2018). Community-level assessment of floods and cyclones in coastal Odisha, India: Impact, resilience, and implications. Research brief. New Delhi: Population Council.
- Prusty B A K. (2011). A book review on "Climate Change and Chemicals: Environmental and Biological Aspects", *Current Science*, **100**,121-122.
- Raghavan K., Sanjay J., Gnanaseelan C., Mujumdar M., Kulkarni A., Chakraborty S. (2020). Assessment of Climate Change over the Indian Region A Report of the Ministry of Earth Sciences (MoES), Government of India, A Report of the Ministry of Earth Sciences (MoES), Government of India, <https://doi.org/10.1007/978-981-15-4327-2>.
- Ramachandran A., Saleem K. A., Palanivelu K., Prasannavenkatesh R., and Jayanthi, N. (2017). Projection of climate change-induced sea-level rise for the coasts of Tamil Nadu and Puducherry, India using SimCLIM: a first step towards planning adaptation policies, *Journal of Coastal Conservation*, **21**, 731–742.
- Rapholo A M T., and Makia L D. (2020). Are smallholder farmers' perceptions of climate variability supported by climatological evidence? Case study of a semi-arid region in South Africa, *International Journal of Climate Change Strategies and Management*, **12**, 571-585.
- Sahoo B., and Bhaskaran P K. (2016). Assessment on historical cyclone tracks in the Bay of Bengal, east coast of India, *International Journal of Climatology*, **36**, 95–109.
- Sahoo B., and Bhaskaran P K A. (2017). comprehensive data set for tropical cyclone storm surge-induced inundation for the east coast of India, *International Journal of Climatology*, **38(6)**, 403–419.
- Sahoo H., Dehury S., Mishra R. (2020). Impact of cyclone Fani on tree damage in Bhubaneswar city, Odisha, India, *e-planet*, **17(2)**,134-138.
- Sathyan A R., Funk C., Aenis T., Breurer L. (2018). Climate vulnerability in rainfed farming: analysis from India watershed, *Sustainability*, **10**, 3357.
- Thornton P K., Ericksen P J., Herrero M., and Challinor A J. (2014). Climate variability and vulnerability to climate change: a review, *Global Change Biology*, **20(11)**, 3313–3328.
- Tulane University. (2021). Tree-Killing Hurricanes Could Contribute to Global Warming, *Science Daily*, Retrieved September 7, 2021 from www.sciencedaily.com/releases/2009/05/090501201353.htm.
- UNDP. (2014). Human Development Report, Sustaining Human Progress: Reducing Vulnerabilities and Building Resilience.
- UNDP. (2014). Multi-Hazard Risk and Vulnerability Analysis for the City of Bhubaneswar, Odisha, 174p.
- Warrick R A. (2009). Using SimCLIM for modeling the impacts of climate extremes in a changing climate: a preliminary case study of household water harvesting in Southeast Queensland. 18thWorld IMACS/MODSIM Congress, Cairns, Australia, pp. 2583–2589.
- WORLD BANK. (2013). Annual Technical Report 2013, Washington DC.

Supplementary File

Questionnaire Survey Form

QUESTIONNAIRE FOR FARMERS IN BHUBANESWAR

A. General Information of the farmers

Name:	Address:	Occupation: Main: Secondary
Gender: <input type="checkbox"/> Male <input type="checkbox"/> Female	Age: <input type="checkbox"/> Below 30 years <input type="checkbox"/> Between 30-40 years <input type="checkbox"/> Between 40-50 years <input type="checkbox"/> Above 50 years	Qualification: <input type="checkbox"/> Below Matric <input type="checkbox"/> Matric <input type="checkbox"/> Graduate <input type="checkbox"/> Post Graduate <input type="checkbox"/> Any other, (please specify)_____
Family Members		
Size of land holding under operation? <input type="checkbox"/> Below 1 acres <input type="checkbox"/> Between 1 - 2.5 acres <input type="checkbox"/> Between 2.5 - 5 acres <input type="checkbox"/> Between 5 - 10 acres <input type="checkbox"/> Above 10 acres		Annual Household Income (Gross in Rs.) Below 1.5 lakhs (1810.01 USD) 1.5 - 3 lakh (1810.01-3620.02 USD) 3 - 5 lakhs (3620.02-6033.36 USD) 5- 10 lakhs (6033.36-12066.73 USD) Above 10 lakhs (12066.73 USD)
Type of Crop farming: <input type="checkbox"/> Rabi <input type="checkbox"/> Kharif <input type="checkbox"/> Other, Specify_____		
Source of water for Agriculture: <input type="checkbox"/> Irrigated <input type="checkbox"/> Rainfed <input type="checkbox"/> Other, Specify_____		

Questions on temperature change perceptions

1. Do you feel any change temperature in past few years?
 Yes No Don't Know Any other _____
 If yes, whether it is Decreasing or Increasing
2. Are the number of hot days has increased?
 Yes No Can't say Any other _____
3. Are the number of cool days decreased?
 Yes No Can't say Any other _____
4. How rainfall quantity in the city has changed?
 Increased Decreased No change Any other _____
5. Erratic rainfall distributions
 Yes No Can't say Any other _____
6. Reasons for temperature change
 Deforestation Urbanization Cyclones Other
7. Do you believe global warming is due to natural or anthropogenic causes?
 Natural causes Anthropogenic causes

- Both natural and anthropogenic There is no global warming
8. Drought has occurred more frequently in recent years?
 Yes No Don't Know Any other _____
9. Temperature variation increases
 Farming losses Reduced rice quality Reduced rice yield
 Cost of water management Reduced rice price due to loss of the quality
 Resulted in farmer's inability to repay loans Crop failure
 Increased the application rate of chemical inputs
 Resulted in the desolation of some farms
10. Which of the climate change impacts you are facing in your area?
 Flood Cyclone Drought Heat Waves Other