

Biowaste compost amendment is a source of microplastic pollution in agricultural soils

Sindujaa A.M., Dhinakaran G.* and Merline Sheela A.


Centre for Environmental Studies, Department of Civil Engineering, Anna University, Chennai – 600 025, Tamil Nadu, India

Received: 26/08/2023, Accepted: 22/09/2023, Available online: 06/10/2023

*to whom all correspondence should be addressed: e-mail: twinsdina@gmail.com

<https://doi.org/10.30955/gnj.005331>

Graphical abstract



Abstract

Microplastics were discovered in soil samples taken from fields in Tamil Nadu, India's Ponneri, Redhills, and Thirumazhisai where the fields had been amended with compost made from biowaste. The soil sample taken from agricultural field in Ponneri had the highest microplastic level, which was determined to be 3.28 ± 0.8 g/kg. The duration of compost addition has a big impact on how microplastics are distributed. When compared to Redhills (5 years' compost amendment) and Thirumazhisai (3 years' compost amendment), the agricultural fields of Ponneri had greater microplastic concentration after 8 years of biowaste compost amendment. The Scanning Electron Microscope (SEM) analysis revealed the presence of multiple forms of microplastics in the study area, including fragments, fibres, and pellets. SEM research confirmed that the surface topography of microplastics exhibited cavities and cracks as a result of weathering activity. Fourier transform infrared (FTIR) spectroscopy analysis confirmed that polypropylene type microplastic was present in the study area. Additionally, the presence of microplastics in the biowaste compost had no impact on the bacterial population or microbial activity in the soil.

Keywords: Agricultural soils, biowaste compost, microplastics distribution, soil microbial activity

1. Introduction

Our daily lives are becoming more and more reliant on plastic. The annual global production of plastic is about 322 million tonnes (Zhang *et al.*, 2020). A whopping 85% of used plastics are not recycled and end up in the

environment (Alabi *et al.*, 2019), where solar radiation and weathering cause them to disintegrate into small bits known as microplastics (Andrady 2022; Zhang *et al.*, 2020). The particle size of the microplastics is 1– 5000 μm (Frias *et al.*, 2019) and are considered as emerging pollutants that have a severe impact on the ecosystem (Sharma *et al.*, 2017; Townsend *et al.*, 2019; Prata *et al.*, 2020). Plastics are introduced into soils through sewage sludge (Corradini *et al.*, 2019; Schell *et al.*, 2022), irrigation, soil runoff (Pérez-Reverón *et al.*, 2022; Henry *et al.*, 2019), and plastic film mulching (Kumar and Sheela, 2022; Khalid *et al.*, 2023). Another source of soil microplastics (Blasing and Amelung 2018; Scopetani *et al.*, 2022; Zhang *et al.*, 2022) is the addition of compost to fields. Compost made from various raw materials, including kitchen waste (Yang *et al.*, 2019), poultry litter (Cao *et al.*, 2022; Sinduja *et al.*, 2021), cow manure (Haroon *et al.*, 2022), garden waste (Langsdorf *et al.*, 2021), and vegetable waste (Ghinea and Leahu 2020), is frequently used in agricultural fields to boost the growth of crops. Municipal solid waste (MSW) compost is currently used to enhance the physicochemical characteristics of soil, including soil structure, water holding capacity, buffering capacity, nutritional status (Kranz *et al.*, 2020), and microbial biomass and activity (Azeem *et al.*, 2022; Mandal *et al.*, 2020). India's per capita generation of solid waste is rising as a result of urbanisation, industrialization, and economic expansion (Kumar *et al.*, 2017).

In India, there are over 133760 tonnes of Municipal Solid Waste (MSW) produced per day (CPCB India 2018). It comprises of home garbage, non-hazardous solid waste thrown by industrial, commercial and institutional institutions, market waste, yard waste and street sweepings (Sharma and Jain 2019; Kranz *et al.*, 2020). MSW is processed to provide energy and compost since it has a high organic matter content (Joseph *et al.*, 2012). The MSW's separated organic part can be composted right away (Kumar and Agarwal 2022). The compost generated from MSW is the source of undesirable inert materials like plastic, glass, and metals that could cause harmful effects to the plants (Balasubramanian 2018). Microplastics are mainly found in the MSW compost due

to food waste and sludge/biosolids (Golwala *et al.*, 2021). Moreover, the compost of household waste has microplastics derived from synthetic fibers from clothing and personal care products (Hernandez *et al.*, 2021). Microplastics of various forms and types can also be found in compost made from rural domestic waste (Gui *et al.*, 2021). Plastic particles were abundant (12 ± 8 – 46 ± 8 particles/kg) in various types of compost tested (Braun *et al.*, 2021). The primary source of plastic particles in the compost may also be the disposal of packaged food waste in the organic waste bin (Katharina *et al.*, 2020). A potential cause of microplastic pollution is the amendment of biowaste compost for agricultural and horticultural crops (Porterfield *et al.*, 2023). Plastics were introduced into arable soils as a result of MSW compost amendment (Blasing and Amelung 2018). Microplastics have been studied in soil amended with MSW compost for ten years (Watteau *et al.*, 2018). However, the vast majority of studies are concentrated on aquatic ecosystems, and there is very little information on microplastic pollution in terrestrial environments. Furthermore, no research has yet been done in India to look into the presence of microplastics in agricultural soil that has been modified using compost made from biowaste. Therefore, the current study was carried out to (1) examine the microplastics in agricultural soils and horticultural soils amended with biowaste compost, (2) investigate the effect of microplastic distribution on soil bacterial population, (3) study the correlation between microplastic content and soil microbial activity.

2. Materials and methods

2.1. Sampling site

The samples were obtained from three different agricultural soils in the Thiruvallur District of Tamil Nadu, where the soils were amended with biowaste compost: Ponneri ($13^{\circ}20'31.6''N$ $80^{\circ}11'54.9''E$), Redhills ($13^{\circ}11'47.0''N$ $80^{\circ}11'11.0''E$) and Tirumazhisai ($13^{\circ}03'17''N$

$80^{\circ}03'12''E$). According to Periyasamy *et al.* (2018), the district receives an average range of 970 to 1100 mm of precipitation each year. Tube wells are the source of irrigation for the chosen study locations. The details of the crop grown in the study area and the duration of biowaste compost amendment are shown in Table 1.

The paddy crop was rotated with groundnut or pulse crops in the Ponneri and Thirumazhisai regions. In Redhills, vegetable crops including brinjal were sown in the field. Biowaste compost was used to amend fields in Ponneri, Redhills, and Thirumazhisai for 8, 5, and 3 years, respectively. The compost was amended @ 20 t/ha obtained from same source. Figure 1 depicts where the sampling fields are located.




Figure 1. Locations of sample collection (a) Ponneri (b) Redhills (c) Thirumazhisai

Table 1. Particulars about the agricultural field where the study was conducted

Location	Crop grown during sampling	Area of land (ha)	Crop rotation followed with	Duration of compost amendment (years)
Ponneri	Paddy (<i>Oryza sativa</i>)	6.47	Groundnut(<i>Arachis hypogaea</i>), Greengram (<i>Vigna radiata</i>), Blackgram (<i>V. mungo</i>)	8
Redhills	Brinjal (<i>Solanum melongena</i>)	16.18	Greengram (<i>V. radiata</i>), Sesame (<i>Sesamum indicum</i>)	5
Thirumazhisai	Paddy (<i>Oryza sativa</i>)	6.07	Blackgram (<i>V. mungo</i>), Greengram (<i>V. radiata</i>)	3

2.2. Sample collection

The soil samples were obtained from the agricultural fields of Ponneri, Redhills and Thirumazhisai and amended with biowaste compost. Each field was divided into four segments. In each segment, 5 m from the edge a 3×3 m² size square plot was marked, and from within the plot, five samples (each 1 kg) were obtained at a depth of 10–30 cm randomly using a spade (Schell *et al.*, 2022) A composite sample was created by combining the collected samples. Similar to the fourth sample, the fifth was also taken from the middle of the field. From each location, a control soil (virgin) sample was collected from 500 m

distal to the biowaste compost amended field as a reference. The samples were then packed in squeaky-clean glass bottles and transported to the laboratory in an ice box. Subsequently, the clumps of samples were broken to obtain a homogeneous sample. The samples were dried in the air before stones and plant debris were manually removed. The powdered air-dried soil was put in glass containers and stored at 4°C for further study.

2.3. Characterization of soil samples

The parameters, such as pH (1:2.5 (soil:water)) using pH metre (Elico L1120), electrical conductivity (EC) using a digital conductivity metre (Labline) (Jackson 1973) and

total organic carbon (TOC) (Walkley and Black 1934), were determined.

2.4. Extraction and identification of microplastics

The soil samples were first dried at 70 °C in an oven before being sieved with a mesh size of 20. Then, 100 mL of saturated NaCl (AR, 99.5%, 8.0% w/v; $\rho=1.2 \text{ g/cm}^3$) was added to the 250-mL conical flasks with the sieved material, and the combination was incubated for 24 hours. The mixture was then treated with hydrogen peroxide (H_2O_2 35%) (Dubai and Liebezeit 2013) to eliminate the organic material before being filtered through a filter assembly (pore size 1–6 μm). Subsequently, the filtrates were dried at 60 °C in a hot air oven (Model BST/HAO-1122; Bionics Scientific Technologies (P) Ltd, India) for 24h. The amount of microplastic in the samples was measured using the initial and final weights of the filter paper.

Microplastics were classified into three categories based on the findings of the visual analysis performed using a stereomicroscope (Accu-Scope; Thermo Scientific Inc.): absence of visible cellular or organic structures in the microplastics, uniformity along the entire length of the fibres, and clear and homogeneous colours (Qiu *et al.*, 2016). Microplastics were divided as fragments, films (thin, soft, and filmy), fibres (elongated), and pellets based on the observations (Jabeen *et al.*, 2017). Furthermore, using SEM-Edx (Carl Zeiss MA15/EVO 18, resolution of 3.0 nm with a 30-kV SE detector), the surface morphology and elemental concentration of the selected microplastics were evaluated. Using double-sided adhesive carbon tabs, the microplastic samples were fixed to the aluminium SEM stubs (Wanget *et al.*, 2017). Using an EmCrafts Virtuoso 1.1 operating at 20 kV at a working distance of 15 nm, the samples were photographed (Tiwari *et al.*, 2017). The polymer type of the microplastics was determined using an IR Affinity-1S FTIR spectrophotometer (Shimadzu, Japan). The FTIR spectra were averaged over 500 scans and measured in the wavelength range of 500–4000 cm^{-1} at a resolution of 4 cm^{-1} (Tiwari *et al.*, 2017). The FTIR absorbance spectra of the microplastics were compared to those in the OMNIC polymer reference library to identify the particles.

2.5. Biological characteristics

The soil microbial population of bacteria was enumerated using soil extract agar medium (Allen 1957). The data are represented as colony forming units (CFU)/g soil (dry weight basis). The biological reduction of triphenyl tetrazolium chloride (TTC) by microorganisms was used to

estimate the dehydrogenase enzyme activity, which is a measure of microbial activity (Casida 1964).

2.6. Quality control/quality assurance

Deionized water was used to thoroughly clean all equipment and containers prior to use. In order to get a composite sample, five samples were taken from each sampling point. The samples were promptly moved to glass vials and closed to avoid on site contamination. The samples were transported in an ice box to preserve the viability of the bacterial species. Analytical-grade chemicals were used for all of the analyses. Deionized water was used to create each and every reagent used in this study. Plastic materials were not used in the analysis of microplastics. The sieve, mesh and mortar and pestle used were made of stainless steel. Also, in the working space, the furnishing and carpeting materials comprised of synthetic fibers were avoided. The work table was cleaned with alcohol prior to the analysis beginning. In order to prevent contamination from synthetic fibres and hairs during conducting the experiments, a white cotton coat, cap, and gloves were worn throughout the extraction of microplastics. The samples were always kept in a closed condition. The spectrophotometer was calibrated using the blank solution (10mL of 0.2 M $\text{Fe}(\text{NO}_3)_3$ diluted to 25 mL with 0.1 M HNO_3) prior to taking readings to determine the dehydrogenase enzyme activity. Furthermore, each analysis was carried out in five replicates to obtain an accurate result.

2.7. Data analysis

The data were statistically analyzed using SPSS 19.0 and the results were presented as mean \pm standard error of the mean.

3. Results and discussion

3.1. Physicochemical characteristics

The data are shown in Table 2 for the parameters such as pH, EC, and TOC that were examined for the soil samples taken from Ponneri, Redhills, and Thirumazhisai. The parameters were pH 7.1–7.6, EC 72.3–81.2 $\mu\text{S/cm}$, and TOC 19.12 \pm 1.2–23.5 \pm 1.4%. As a result, it was found that the characteristics of the samples taken from various study areas varied only little from one another. The outcomes matched those of previous studies (Chandrasekaran and Ravisankar 2015; Kumar and Sheela 2021). The characteristics of the soil are determined by the type of soil and the amount of organic matter in surface soils (Schilatter *et al.*, 2017).

Table 2. Physicochemical characteristics of soil samples

Parameter	Sampling locations					
	Ponneri		Redhills		Thirumazhisai	
	Control	Sample	Control	Sample	Control	Sample
pH	7.3	7.2	7.1	7.0	7.2	7.6
EC ($\mu\text{S/cm}$)	74.1	72.6	75.9	79.8	81.2	72.3
TOC (%)	20.5 \pm (1.6)	20.13 \pm (1.5)	19.2 \pm (1.2)	22.3 \pm (1.8)	23.5 \pm 1.4	22.5 \pm (1.2)

TOC – Total Organic Carbon, Values in parenthesis represent \pm Standard Error ($n=5$)

3.2. Microplastic content in the study area

The microplastic content in the sampling locations, Ponneri, Redhills and Thirumazhisai, was determined and shown in Figure 2(a).

The microplastic content in samples taken from Ponneri, Redhills, and Thirumazhisai ranged from 0.29 ± 0.1 – 3.28 ± 0.8 , 0.42 ± 0.1 – 1.83 ± 0.2 and 0.36 ± 0.2 – 1.82 ± 0.2 g/kg soil, respectively. According to the results, soil samples taken from Ponneri's agricultural areas contain more microplastic than those taken from Redhills and Thirumazhisai. Figure S1 (b) shows the impact of the duration of biowaste compost addition on the presence of microplastic in agricultural soils. We found that the microplastic content increased with the duration of compost amendment. The soils collected from Ponneri, where compost addition was carried out for 8 years, reported the highest microplastic content compared to Redhills and Thirumazhisai.

Microplastics were discovered in soil samples collected from three different locations in the Thiruvallur District of Tamil Nadu, India, for the current study. However, understanding of the presence of microplastics in the terrestrial environment, particularly in agricultural soils, is still inadequate. Only a small number of recent studies have shown the accumulation of plastic debris in agricultural soils (da Costa 2018). The Chen *et al.* study (Chen *et al.*, 2020) confirmed the existence of microplastics in the Chinese agricultural soils. Previous studies confirmed the existence of microplastics in agricultural soils as the outcome of sewage sludge discharge (Corradini 2021), treating effluents for irrigation of crops (Li *et al.*, 2022; Kumar *et al.*, 2020), mulching with plastic film (Huang *et al.*, 2020; Kumar and Sheela 2021), and compost (Watteau *et al.*, 2018).




Figure 2(a) Microplastic content in different locations of the study area (Error bars represent \pm S. E (n = 5)) & 2(b) Effect of duration of biowaste compost amendment on the occurrence of microplastics in agricultural soils

In the present study, neither treated effluent nor sewage sludge were used to modify the soils. Additionally, there was no plastic film mulching in this study area. In addition to sewage sludge and plastic mulches, the study by Huang 2020 *et al.*, found that compost amendment was another source of microplastic pollution in agroecosystems. Composting is applied to agroecosystems all over the world to increase crop growth and yield (Wei *et al.*, 2000). When farmers fertilise their cultivable land with sewage

sludge and compost made from biosolids, they unintentionally add microplastics to the soil (Henseler *et al.*, 2020). As an outcome, the presence of microplastics in the study area is attributable to the biowaste compost's amendment. Solid waste is a substantial source of microplastics in a range of sizes and shapes (Golwala *et al.*, 2021). Biowaste from MSW is often segregated and composted. However, household biowaste is sometimes not properly separated or disposed of in high-density plastic bags (Zumstein *et al.*, 2018; Melgarejo *et al.* 2000). Furthermore, plastic containers used for packaging food are disposed of with spoiled food or leftovers to the biowaste collection bins. It is difficult to separate or remove the plastic particles once they have mixed with the biowaste (Weithmann *et al.*, 2018). In addition, the biodegradable plastic materials are not fully biodegraded in the composting plants because of less process period (Mallikarjuna and Bindu 2016; Pathan *et al.*, 2020). Possible pathways for plastics to enter agricultural soils include compost that has been modified with plastic particles. These particles are further broken down into smaller particles, called microplastics, when exposed to sunshine and UV rays (Shimizu *et al.*, 2017). These then form aggregates with the soil's organic matter particles and stay in the ground for a long time (Peng *et al.*, 2017). In the current study, microplastics have been identified up to a depth of 30 cm. The movement of microplastics into deeper layers is achieved by the colonization of micro arthropods (Maaß *et al.*, 2017), which is a conventional tillage practice (Paustian *et al.*, 2000).

3.3. Type and forms of microplastics

Figure 3 displays the many kinds of microplastics that were investigated in the study. In Ponneri samples, there were 450 ± 25 , 300 ± 25 and 250 ± 10 pellets, pieces, and fibres that were evaluated. The Redhills samples had 320 ± 20 , 200 ± 20 , and 100 ± 12 fragments, pellets, and fibres, respectively. The samples from Thirumazhisai also contained 50, 18, 60, and 38 fragments, pellets, and fibres, respectively. According to the findings, fragments of microplastics were found most frequently in each of the three locations.




Figure 3. Different types of microplastics observed in the study area (Error bars represent \pm Standard Error, n = 5)

In all the locations microplastics in black color was predominant. The other colors examined were white and red. The percent distribution of black, white and red color microplastics in samples of Ponneri was 55%, 25% and

20% respectively. The samples of Redhills harbored 79% (black), 18% (white) and 8% (red) microplastics. The samples collected from Thirumazhisai contained 67% (black), 27% (white) and 6% (red) microplastics (Figure 4).




Figure 4. Percent distribution of different colours of microplastics in the study area (a) Ponneri (b) Redhills (c) Thirumazhisai

The microplastics found in the research area were identified as fibres, pellets, and pieces (Figure 5). A rough surface with eroded cavities was visible in the morphology under SEM (Figure 6). According to the peaks and heights of the peaks, the SEM- Energy Dispersive X-Ray analysis indicated the existence of various secondary elements at various concentrations (Figure 6(a)). Carbon (C), oxygen (O), iron (Fe), aluminium (Al), silicon (Si), and chlorine (Cl) were the elements that were found.

The presence of microplastics of the polypropylene type has been confirmed by the FTIR polymer type analysis of the fragments collected from the study area (Figure 6(b)). The presence of microplastics of the polypropylene type in the studied areas was established by the mapping of the intensity of bands (2905-C-H stretching; 1415-C=O stretching).



Figure 5. Photomicrograph of different forms of microplastics examined in the study area (a) Fragment (b) Fiber (c) Pellet (Examined at 45x magnification. Scale bars represent 200µm)

3.4. Effect of microplastics on biological characteristics

In soil samples taken from Ponneri, the bacterial population was 2×10^5 cfu/g (control), 8×10^5 cfu (field), 6×10^5 cfu/g (control), 9×10^5 cfu/g (field samples), 4×10^6 cfu/g (control), and 12×10^6 cfu/g (field sample) (Figure 7(a)). The dehydrogenase enzyme activity of the soil samples taken from the study area has been determined and the data is displayed in Figure 7b. The soil samples taken from Ponneri had the highest dehydrogenase enzyme activity, 0.94 µg Triphenyl Formazon (TPF)

formed/h/g, while the soil samples taken from Thirumazhisai had the lowest dehydrogenase enzyme activity, 0.81 µg TPF formed/h/g. Furthermore, for the soils in Ponneri, Redhills, and Thirumazhisai, the dehydrogenase activity was 0.89, 0.74, and 0.79 µg TPF formed/h/g, respectively.




Figure 6. (a) SEM-EDX spectra to confirm the presence of individual elements in the plastic particle (b) FTIR spectrum of Polypropylene type microplastic in the study area (Horizontal axis represents the spectral range and vertical axis represents the transmittance %.)




Figure 7. (a) Effect of biowaste compost amendment on bacterial population in the study area (Values represent mean of five replications), (b) Dehydrogenase enzyme activity in biowaste compost amended soil (Values represent mean of five replications)

The bacterial population was higher in soil samples from the Thirumazhisai region than it was in soil samples from Ponneri and Redhills. The results revealed that the microplastic content was less as this region was amended with biowaste compost for 4 years. The microplastic effect on soil and soil organisms depends on the type and concentration of plastics and exposure time (Guo *et al.*, 2020). Hitherto, only a limited number of studies have been carried out to examine the toxicology of microplastics to soil microorganisms (He *et al.*, 2018).

Conversely, some studies explained that soil microorganisms, including bacteria, are affected by exposure to high concentrations of microplastics (Bradney *et al.*, 2019). However, the bacterial population of soil

samples taken from agricultural soils in the present study was not significantly impacted by microplastic pollution. The bacterial population in the soil taken from agricultural fields is comparable to that of the virgin soil collected 500 metres away from compost-amended soil. The cracks and cavities formed on the surfaces due to weathering and altered chemical properties provide a suitable condition for the colonization of various microorganisms (Zhang *et al.*, 2019). Some studies explained that other toxic pollutants in soil are absorbed on the microplastics or within the plastic ecocorona (the bound form of organic material and contaminants) (Galloway *et al.*, 2017), which protect the soil biota from toxic pollutants (Du *et al.*, 2020; Sarker *et al.*, 2020). However, whether the effect of microplastics on soil organisms is positive or negative needs to be investigated further.

Dehydrogenase, one of the most crucial hydrolases in microbial respiration which is present in all living microbial cells (Song *et al.*, 2019). In this study, it was found that soils amended with compost made from biowaste had higher levels of enzyme activity than control soils. According to previous studies, the addition of microplastic enhanced the soil's dehydrogenase enzyme activity (Yi *et al.*, 2021; Kumar and Sheela 2023), which may be attributable to either an improvement in the habitat of the soil (Zhao *et al.*, 2021) or the adaptation of microorganisms to the microplastic amended environment (Yi *et al.*, 2021). Also, the study by (Kumar and Sheela 2021) demonstrated that microbial activity did not get influenced by microplastic addition due to plastic film mulching. As a result, the bacteria in the soil may get nutrients from the compost made from biowaste and grow. The increased dehydrogenase activity in the study area compared to that in the virgin soil may be caused by an increase in the microbial population.

The harmful impacts of microplastics on plants are still not thoroughly studied (Jiang *et al.*, 2019). The different kinds of microplastics have an impact on shoot and root length reduction (van Weert *et al.*, 2019) as well as seed germination (Bosker *et al.*, 2019). Additionally, the interaction of heavy metals and microplastics has an adverse impact on soil biodiversity and plant growth (Wang *et al.*, 2020).

In contrast to primary microplastics, ageing of microplastics does not have an adverse effect (Kokalj *et al.*, 2019). However, in the current study, samples were gathered from agricultural soils amended with biowaste compost. The microplastics in the compost amended soil might have undergone ageing process, thereby effectuating less toxicity to the soil bacterial population. Nonetheless, extensive field monitoring is required to evaluate the potentially toxic effects of microplastics to crop species at various growth stages in the biowaste compost amended soils. The entry of microplastics from the compost to the agroecosystem might be prevented by adopting proper solid waste treatment method. The plastic materials that are adherent to the waste should be properly removed before using it for the composting process so as to prevent microplastic pollution of the agroecosystem.

4. Conclusion

Microplastics in agricultural soils of Ponneri, Redhills, and Thirumazhisai in Thiruvallur district, Tamil Nadu, India, were studied as a result of biowaste compost addition. The microplastic content in the soil samples collected from Ponneri, Redhills, and Thirumazhisai ranged from 0.29 ± 0.1 – 3.28 ± 0.8 , 0.42 ± 0.1 – 1.83 ± 0.2 and 0.36 ± 0.2 – 1.82 ± 0.2 g/kg soil, respectively. The agricultural soils of Ponneri encountered more microplastic content than other samples due to the long duration of biowaste compost amendment. In the study area, different forms of microplastics, including fragment, fibre, and pellet, were studied. In the study area, the three colours of microplastics that were looked at were black, white, and red. Polypropylene was the primary polymer type identified in this study area. The biowaste compost amendment does not have a substantial effect on the soil bacterial population and microbial activity. The presence of less concentration of microplastics in the virgin soils collected near the compost-amended soil might be either due to plastic littering or transported from other places by the wind. However, extensive study is required to examine the sources and effects of microplastic pollution in agricultural soils. Studies are also required to look at how microplastic pollution affects the structure of microbial communities, the availability of nutrients to crops, the germination of seeds, and crop growth. Therefore, better plastic waste management may help to prevent the accumulation of microplastics in terrestrial ecosystems like agricultural soils.

Author Contributions

Sindujaa A.M. (PG Student) conducted the experiment and wrote the original draft. Dhinakaran G (Assistant Professor) and Merline Sheela A. (Assistant Professor) supervised the student and edited the manuscript.

Acknowledgements

The authors are grateful to the Centre for Environmental Studies (CES), Anna University, Chennai, India, for the laboratory facility provided to carry out this study. The authors also acknowledge the arrangements made by A. Ravi, Lab Assistant, to collect and analyse the samples.

References

- Acosta-Coley I., Mendez-Cuadro D.Z., Rodriguez-Cavallo E., de la Rosa J. and Olivero-Verbel J. (2019). Trace elements in microplastics in Cartagena: a hotspot for plastic pollution at the Caribbean, *Marine Pollution Bulletin*, **139**, 402–411.
- Alabi O.A., Ologbonjaye K.I., Awosolu O., Alalade O.E. (2019). Public and Environmental Health Effects of Plastic Wastes Disposal: A Review, *Journal of Toxicol Risk Assess*, **5**, 021.
- Allen O.N. (1957). Experiments in soil bacteriology. 3rd ed. Burgess; Publishing Co, Minneapolis, Minn, p.150.
- Andrady A.L. (2022). Weathering and fragmentation of plastic debris in the ocean environment, *Marine pollution bulletin*, **180**, 113761.
- Azeem I., Adeel M. and Ahmad M.A. et al. (2022). Uptake and Accumulation of Nano/Microplastics in Plants: A Critical Review, *Nanomater*, **11**(11), 2935.

- Balasubramanian M. (2018). Municipal solid waste management in India: Status, problems and challenges, *International Journal of Environment and Waste Management*, **21**(4), 253–268.
- Blasing M. and Amelung W. (2018). Plastics in soil: Analytical methods and possible sources, *Science of the Total Environment*, **612**, 422–435.
- Bosker T., Bouwman L.J., Brun N.R., Behrens P. and Vijver M.G. (2019). Microplastics accumulate on pores in seed capsule and delay germination and root growth of the terrestrial vascular plant *Lepidium sativum*, *Chemosphere*, **226**, 774–781.
- Bradney L., Wijesekara H., Palansooriya K.N., Obadamudalige N., Bolan N.S., Ok Y.S., Rinklebe J. and Kirkam M.B. (2019). Particulate plastics as a vector for toxic trace-element uptake by aquatic and terrestrial organisms and human health risk, *Environment International*, **131**, 104937.
- Braun M., Mail M., Hayse R. and Amelung W. (2021). Plastic in compost: Prevalence and potential input into agricultural and horticultural soils, *Science of the Total Environment*, **760**, 143335.
- Cao Y., Bai M. and Han B. et al. (2020). Enhanced nitrogen retention by lignite during poultry litter composting, *The Journal of Cleaner Production*, **277**, 122422.
- Casida L., Klein D. and Santoro T. (1964). Soil dehydrogenase activity, *Soil Science*, **98**, 371–376.
- Chandrasekaran A. and Ravisankar S. (2015). Spatial distribution of physico-chemical properties and function of heavy metals in soils of Yelahiri hills, Tamil Nadu by Energy Dispersive x-ray fluorescence spectroscopy (EDXRF) by statistical approach. *Spectrochimica Acta, Part A: Molecular and Biomolecular Spectroscopy*, **150**, 586–601.
- Chen B., Fan Y., Huang W., Rayhan A.B.M.S., Cai M. and Chen K. (2020). Observation of microplastics in mariculture water of Longjiao Bay, southeast China: influence by human activities, *Marine pollution bulletin*, **160**, 111655.
- Corradini F., Casado F., Leiva V., Huerta-Lwanga E. and Geissen W. (2021). Microplastics occurrence and frequency in soils under different land uses on a regional scale, *Science of the Total Environment*, **752**, 141917.
- Corradini F., Meza P. and Eguiluz R. et al. (2019). Evidence of microplastic accumulation in agricultural soils from sewage sludge disposal, *Science of the Total Environment*, **671**, 411–420.
- CPCB India 2018. MSW- Generation & Composition CPCB 2018
- da Costa J.P., Paco A., Santos P.S.M., Duarte A.C. and Rocha-Santos T. (2018). Microplastics in soils: assessment, analytics and risks, *Environmental Chemistry*, **16**(1), 18–30.
- Du C., Wu J., Gong J., Liang H., Li Z. (2020). ToF – SIMS characterization of microplastics in soils, *Surface and Interface Analysis*, **52**(5), 29–300.
- Dubaish F. and Liebezeit G. (2013). Suspended Microplastics and Black Carbon Particles in the Jade System, Southern North Sea, *Water Air Soil Pollutant*, **224**, 1352.
- Fotopoulou K.N. and Karapanagioti H.K. Degradation of Various Plastics in the Environment, In: Takada H., Karapanagioti H. (eds) Hazardous Chemicals Associated with Plastics in the Marine Environment, *The Handbook of Environmental Chemistry*, vol 78, Springer, Cham. 2017.
- Fotopoulou K.N. and Karapanagioti K.K. (2012). Surface properties of beached plastic pellets, *Marine Environmental Research*, **81**:70–77
- Frias J.P.G.L. and Nash R. (2019). Microplastics: Finding a consensus on the definition, *Marine pollution bulletin*, **138**, 145–147.
- Galloway T., Cole M., and Lewis C. (2017). Interactions of microplastic debris throughout the marine ecosystem, *Nature Ecology and Evolution*, **1**, 0116.
- Ghinea C. and Leahu A. (2020). Monitoring of Fruit and Vegetable Waste Composting Process: Relationship between Microorganisms and Physico-Chemical Parameters, *Processes*, **8**(3), 302.
- Golwala H., Zhang X. and Md Iskander S. et al. (2021). Solid waste: An overlooked source of microplastics to the environment, *Science of the Total Environment*, **769**, 144581.
- Golwala H., Zhang X., Md Iskander S. and Smith A.L. (2021). Solid waste: An overlooked source of microplastics to the environment, *Science of the Total Environment*, **769**, 144581.
- Gui J., Sun Y., Wang J., Chen X., Zhang S. and Wu D. (2021). Microplastics in composting of rural domestic waste: abundance, characteristics, and release from the surface of macroplastics, *Environment Pollutant*, **274**, 116553.
- Guo J. Huang X., Xiang L., Wang Y., Li Y., Li H., Cai Q., Mo C. and Wong M. (2020). Source, migration and toxicology of microplastics in soil, *Environment International*, **137**, 105263.
- Haroon B., Hassan A., Abbasi A.M. et al. (2020). Effects of co-composted cow manure and poultry litter on the extractability and bioavailability of trace metals from the contaminated soil irrigated with wastewater, *Journal of Water Reuse Desal*, **10**(1), 17–29.
- He D., Luo Y., Lu S., Liu M., Song Y. and Lei L. (2018). Microplastics in soils: analytical methods, pollution characteristics and ecological risks, *TrAC-Trend analytical chemistry*, **109**, 163–172.
- Henry B., Laitala K. and Klepp I.G. (2019). Microfibres from apparel and home textiles: Prospects for including microplastics in environmental sustainability assessment. *Science of the Total Environment*, **652**, 483–494.
- Henseler M., Brandes E. and Kreins P. Microplastics in agricultural soils: A new challenge not only for agro-environmental policy? 172nd seminar, May 28 – 29, 2020, Brussels, Belgium.
- Hernandez E., Nowack B. and Mitrano D.M. (2017). Polyester textiles as a source of microplastics from households: A mechanistic study to understand microfiber release during washing. *Environmental Science & Technology*, **51**(12), 7036–7046.
- Huang Y., Liu Q., Jia W., Yan C., Wang J. (2020). Agricultural plastic mulching as a source of microplastics in the terrestrial environment, *Environmental Pollution*, **260**, 114096.
- Indumathi D. and Sheela A.M. (2023). Occurrence of Coliforms in Microplastic Associated Biofilm in Estuarine Ecosystem. *Pol. Journal of Environmental Science Studies*, **32**(1), 547–557.
- Jabeen K., Su L., Li J., Yang D., Tong C., Mu J. and Shi H. (2017). Microplastics and mesoplastics in fish from coastal and fresh waters of China, *Environmental Pollution*, **221**, 141–149.
- Jackson M.L. Soil Chemical Analysis Prentice Hall of India Pvt. Ltd., New Delhi (1973), 498.

- Jiang X., Chen H., Liao Y., Ye Z., Li M. and Klobučar G. (2019). Ecotoxicity and genotoxicity of polystyrene microplastics on higher plant *Vicia faba*, *Environmental Pollution*, **250**, 831–838.
- Joseph K., Rajendiran S. and Senthilnathan R. et al. (2012). Integrated approach to solid waste management in Chennai: An Indian Metro City, *Journal of Material Cycles and Waste Management*, **14**, 75–84.
- Katharina M., Mona K., and Ines F. (2020). Microplastic in soil—current status in Europe with special focus on method tests with Austrian samples, *AIMS Environmental Science*, **7(2)**, 174–191.
- Khalid N., Aqeel M., Noman Ali N., Rizvi Z.F. (2023), Impact of plastic mulching as a major source of microplastics in agroecosystems, *Journal of Hazardous Materials*, **445**, 130455.
- Kokalj A.J., Kuehnel D., Puntar B., Žgajnar Gotvajn A. and Kalčíkova G. (2019). An exploratory ecotoxicity study of primary microplastics versus aged in natural waters and wastewaters, *Environmental Pollution*, 254(Part A), 112980.
- Kranz C.N., McLaughlin R.A., Johnson A., Miller G., Heitman J.L. (2020), The effects of compost incorporation on soil physical properties in urban soils – A concise review. *Journal of Environmental Management*, **261**, 110209.
- Kumar A. and Agarwal A. (2020). Recent trends in solid waste management status, challenges, and potential for the future Indian cities – A review, *Current Research in Environmental Sustainability*, **2**, 100011.
- Kumar M., Xiong X. He M., Tsang D.C.W., Gupta J., Khan G., Harrad S., Hou D., Ok Y.S. and Bolan N.S. (2020). Microplastics as pollutants in agricultural soils, *Environmental Pollution*, **265**(Part A), 114980.
- Kumar M.V., and Sheela A.M. (2021). Effect of plastic film mulching on the distribution of plastic residues in agricultural fields, *Chemosphere*, **273**, 128590.
- Kumar R.S. and Sheela A.M. (2023), Microplastics Alter Dehydrogenase, Urease, and Cellulase Activities in Soil, *National Academy Science Letters*, **46**, 325–328.
- Kumar S., Smith S.R., Fowler G., Velis C., Jyoti Kumar S., Arya S., Rena Kumar R. and Cheeseman C. (2017). Challenges and opportunities associated with waste management, *Royal Society Open Science*, **4**, 160764.
- Langsdorf A., Volkmar M. and Holtmann D. et al. (2021). Material utilization of green waste: a review on potential valorization methods. *Bioresources and Bioprocessing*, **8**, 19.
- Li X., Chen L., Mei Q., Dong B., Ding G., Dai X. and Zeng E.Y. (2018). Microplastics in sewage sludge from the wastewater treatment plants in China, *Water Research*, **142**, 75–88.
- Maaß S., Daphi D., Lehmann A. and Rilling M.C. (2017). Transport of microplastics by two collembolan species, *Environmental Pollution*, **225**, 456–459.
- Mallikarjuna V. and Bindu M.T. (2016), Soil Stabilization Using Plastic Waste, *International Journal of Research in Engineering and Technology*, **5**, 301–394.
- Mandal A., Dolamani A., Thakur J.K. and Manna M.C. (2020). Municipal solid waste compost and its effect on soil health, *Indian Journal of Waste Management*, **4(1)**, 19–26.
- Melgarejo P., Legua P., Perez-Sarmiento P., Martínez-Font F., Martínez-Nicolás J.J., Giordan E., Tozzi F. and Hernández F. (2000), Effect of a new remediated substrate on bioactive compounds and antioxidant characteristics of pomegranate (*Punica granatum* L.) cultivar ‘Purple Queen’, *Archives of Agronomy and Soil Science*, **64(11)**, 1565–1574.
- Muthukumar T., Aravinthan A., Lakshmi K., Venkatesan R. and Loganathan V. (2011). Fouling and stability of polymers and composites in marine environment, *International Biodeterioration*, **65(2)**, 276–284.
- Pathan S.I., Arifaoli P., Bardelliu T., Ceccherini M.T., Nannipieri P. and Pietramellara G. (2020). Soil Pollution from Micro-and Nanoplastic Debris: A Hidden and Unknown Biohazard, *Sustainability*, **12**, 7255.
- Paustian K., Six J., Elliott E. and Hunt H.W. (2000). Management options for reducing CO₂ emissions from agricultural soils. *Biogeochemistry*, **48**, 147–163.
- Peng J., Wang J. and Cai L. (2017). Current understanding of microplastics in the environment: Occurrence, fate, risks, and what we should do. *Integrated Environmental Assessment and Management*, **13(3)**, 476–482.
- Pérez-Reverón R., González-Sálamo J., Hernández-Sánchez C., González-Pleiter M., Hernández-Borges J., Díaz-Peña F.J. (2022), Recycled wastewater as a potential source of microplastics in irrigated soils from an arid-insular territory (Fuerteventura, Spain), *Science of the Total Environment*, **15**, 152830.
- Pires J.P., Miranda G.M., de Souza G.L., Fraga F., da Silva Ramos A., de Araújo G.E., Ligabue R.A., s Azevedo C.M.N., Lourega R.V. and de Lima J.E.A. (2019), Investigation of degradation of polypropylene in soil using an enzymatic additive, *Iranian Polymer Journal*, **28**, 1045–1055.
- Porterfield K.K., Hobson S.A., Neher D.A., Niles M.T. and Roy E.D. (2023). Microplastics in composts, digestates, and food wastes: A review, *Journal of environmental quality*, **52(2)**, 225–240.
- Prata J.C., Manana M.J., da Costa J.P., Duarte A.C. and Rocha-Santos T. (2020). What Is the Minimum Volume of Sample to Find Small Microplastics: laboratory Experiments and Sampling of Aveiro Lagoon and Vouga River, Portugal? *Water*, **12**, 1219.
- Qiu Q., Tan Z., Wang J., Peng J., Li M. and Zhan Z. (2016), Extraction, enumeration and identification methods for monitoring microplastics in the environment Estuar. *Coast Shelf Science*, **176**, 102–109.
- Sarker A., Deepo D.M., Nandi R., Rana J., Islam S., Rahman S., Hossain M.N., Islam M.S. and Baroi A. (2020). A review of microplastics pollution in the soil and terrestrial ecosystems: a global and Bangladesh perspective, *Science of the Total Environment*, **733**, 1392963.
- Schell T., Martinez-Perez S., Dafouz R., Hurley R., Vighi M. and Rico A. (2022). Effects of Polyester Fibers and Car Tire Particles on Freshwater Invertebrates, *Environmental Toxicology and Chemistry*, **41(6)**:1555–1567.
- Schilatter D.C., Schillinfger W.F., Bary A.J. Sharratt B. and Paulitz T. (2017). Biosolids and conservation tillage: impacts on soil fungal communities in dryland wheat – fallow cropping system, *Soil Biology and Biochemistry*, **115**, 556–557.
- Scopetani C., Chelazzi D., Cincinelli A., Leiniö V. and Pellinen J. (2022). Hazardous contaminants in plastics contained in compost and agricultural soil, *Chemosphere*, **293**, 33645.
- Sharma K.D. and Jain S. (2019), Overview of Municipal Solid Waste Generation, Composition, and Management in India, *Journal of Environmental*, **145(3)**, 04018143.

- Sharma S. and Chatterjee S. (2017), Microplastic pollution, a threat to marine ecosystem and human health: a short review, *Environmental Science and Pollution Research*, **24**, 21530–21547.
- Shimizu K., Sokolov S.V., Katelhon E., Holter J. and Young N.P. (2017), Compton RG, *Electroanalysis*, **29(10)**, 2200–2207.
- Sinduja B., Sheela A.M. and Ilamathi R. (2021), Sunflower seed husk combined with poultry droppings to degrade petroleum hydrocarbons in crude oil-contaminated soil, *Environmental Engineering Research*, **26(5)**, 200361.
- Song T.W., Li S.S., Jin J., Yin Z., Lu Y., Bao M. and Li Y. (2019). Enhanced hydrolyzed polyacrylamide removal from water by an aerobic biofilm reactor-ozone reactor-aerobic biofilm reactor hybrid treatment system: Performance, key enzymes and functional microorganisms, *Bioresource Technology*, **291**, 1–11.
- Tisdall J.M. and Oades J.M. (1982). Organic matter and water-stable aggregates in soils, *European Journal of Soil Science*, **33(2)**:141–163.
- Tiwari S., Gupta N., Malairaman U. and Shankar J. (2017), Anti-aspergillus properties of phytochemicals against aflatoxin producing *Aspergillus flavus* and *Aspergillus parasiticus*, *National Academy Science Letters*, **40(4)**, 267–271.
- Townsend K.R., Lu H.C. and Sharley D.J., et al. (2019), Associations between microplastic pollution and land use in urban wetland sediments, *Environmental Science and Pollution Research*, **26**, 22551–22561.
- van Weert S., Redondo-Hasselerharm P.E., Diepens N.J. and Koelmans A.A. (2019). Effects of nanoplastics and microplastics on the growth of sediment-rooted macrophytes, *Science of the Total Environment*, **654**, 1040–1047.
- Walkley A. and Black I. (1934). An Examination of Degthareff method for Determining soil Organic matter and a proposed modification of the chromic acid titration method, *Soil Science*, **27**, 29–38.
- Wang F, Zhang X, Zhang S, Zhang S, Sun Y. Interactions of microplastics and cadmium on plant growth and arbuscular mycorrhizal fungal communities in an agricultural soil. *Chemosphere*.2020;254:126791.
- Wang W., Ndungu A.W., Li Z. and Wang J. (2017). Microplastics pollution in inland freshwaters of China: a case study in urban surface waters of Wuhan, China. *Science of the Total Environment*, **575**, 1369–1374.
- Watteau F., Dignac M., Bouchard A., Revallier A. and Houot S. (2018). Microplastic Detection in Soil Amended with Municipal Solid Waste Composts as Revealed by Transmission Electronic Microscopy and Pyrolysis/GC/MS, *Frontiers in sustainable food systems*, **2**, 81.
- Wei Y., Fan Y.B. and Wang M. Wang J. (2000). Composting and Compost Application in China. *Resources, Conservation and Recycling*, **30(4)**, 277–300.
- Weinstein J.E., Dekle J.L., Leads R.R. and Hunter R.A. (2020), Degradation of bio-based and biodegradable plastics in a salt marsh habitat: Another potential source of microplastics in coastal waters, *Marine Pollutant Bulletin*, **160**, 111518.
- Weithmann N., Moller J.N., Loder M.G.J., Laforsch C. and Freitag R. (2018). Organic fertilizer as a vehicle for the entry of microplastic into the environment, *Science advances*, **4(4)**, eaap8060.
- Yang F., Li Y. and Han Y., et al. (2019). Performance of mature compost to control gaseous emissions in kitchen waste composting, *Science of the Total Environment*, **657**, 262–269.
- Yang J., Li L., Xu L., Shen Y., Li S., Tu C., Wu L., Christie P., Luo Y. (2021). Microplastics in an agricultural soil following repeated application of three types of sewage sludge: A field study, *Environmental Pollution*, **289**, 117943.
- Yi M., Zhou S, Zhang L. and Ding S. (2021), The effects of three different microplastics on enzyme activities and microbial communities in soil, *Water Environment Research*, **93**, 24–32.
- Zhang S., Wang J., Liu X., Qu F., Wang X., Wang X., Li Y. and Sun Y. (2019). Microplastics in the environment: A review of analytical methods, distribution, and biological effects, *TrAC-Trends Analytical Chemistry*, **111**, 62–72.
- Zhang S., Zhang W., Ju M., Qu L., Chu X., Huo C., Wang J. (2022), Distribution characteristics of microplastics in surface and subsurface Antarctic seawater, *Science of the Total Environment*, **838**, Part 3, 56051.
- Zhang Y., Kang S., Allen S., Allen D., Gao T and Sillanpää M. (2020). Atmospheric microplastics: a review on current status and perspectives, *Earth-Science Reviews*, **203**, 103118.
- Zhao T., Lozano Y.M. and Rillig M.C. (2021), Microplastics Increase Soil pH and Decrease Microbial Activities as a Function of Microplastic Shape, Polymer Type, and Exposure Time, *Frontiers in Environmental Science*, **9**, 675803.
- Zhou B., Wang J., Zhang H., Shi H., Fei Y., Huang S., Tong Y., Wen D., Luo Y. and Barceló D. (2020). Microplastics in agricultural soils on the coastal plain of Hangzhou Bay, east China: Multiple sources other than plastic mulching film, *Journal of Hazardous Materials*, **388**, 121814.
- Zumstein M., Schintlmeister A., Nelson T.F., Baumgartner R., Woebken D., Wagner M., Kohler H.E., McNeill K. and Sander M. (2018), Biodegradation of synthetic polymers in soils: Tracking carbon into CO₂ and microbial biomass. *Science advances*, **4(7)**, eaas9024.