

Current scenario of ozone pollution and its influence on population health in China

Yongsheng Wang^{1, #}, Xingyu Peng^{2, #}, Haiping Tan³, Lei Zhang⁴, Shu Guo^{3, *} and Ming Zhu^{3, *}

¹College of Environmental and Chemical Engineering, Shanghai University of Electric Power, Shanghai 200090, China

²Loudi Center for Disease Control and Prevention, Loudi, Hunan 417000, China

³State Environmental Protection Key Laboratory of Environmental Pollution Health Risk Assessment, South China Institute of Environmental Sciences, Ministry of Ecology and Environment, Guangzhou 510655, China

⁴Dezhou Center for Disease Control and Prevention, Dezhou, Shandong 253000, China


[#]These authors contributed equally to this work and should be considered co-first authors

Received: 05/09/2023, Accepted: 28/03/2024, Available online: 07/04/2024

*to whom all correspondence should be addressed: e-mail: zhuming@scies.org; guoshu@scies.org

<https://doi.org/10.30955/gnj.005358>

Graphical abstract



Abstract

Ozone (O₃) is a secondary pollutant formed by photochemical reactions in the atmosphere, significantly contributing to air pollution, particularly in global cities and economically developed regions. The escalating O₃ concentration has emerged as a critical air pollution concern in China. When near-surface O₃ surpasses natural levels, it adversely impacts human health. Our understanding of O₃ pollution remains limited, partly due to the delayed implementation of atmospheric O₃ and its precursors' monitoring. Accordingly, utilizing observed data, this paper assesses the O₃ levels in China over the past decade. We found that surface O₃ concentrations had consistently risen since 2013, with the only decline noted in 2018. O₃ pollution is particularly severe in economically developed areas such as the Beijing-Tianjin-Hebei region, the Yangtze River Delta, the Pearl River Delta, and the Weihe Plain. Chronic exposure to O₃ can negatively impact respiratory and cardiovascular systems. By introducing research findings related to O₃ exposure and human health, we offer suggestions for future research on human health implications of surface O₃ exposure. These findings underscore the importance of O₃ as a focal point in China's future air quality policy and highlight the urgent necessity for stricter control of precursor emissions.

Keywords: Ozone; China; pollution trends; ozone exposure; human health

1. Introduction

Air pollution is an important global public health concern. Ozone (O₃) is a common component of air pollutants and is produced by the photochemical reactions of volatile organic compounds (VOCs) and nitrogen oxides (NO_x) emitted by natural and anthropogenic sources under high-temperature ultraviolet irradiation (Sillman 1999). A number of air pollution incidents have harmed human health in the past, such as the smog incidents in Los Angeles in the United States and in London in the United Kingdom (Guan *et al.* 2016). As the largest developing country and the second-largest economy in the world, China has made great economic achievements since its reform and opening up. With the rapid development of China's economy, however, there has been an increase in the consumption of chemical fuels and O₃ pollution is becoming increasingly serious (Liu *et al.* 2015). Previous studies indicated that the distribution of O₃ had strong seasonality and generally showed an inverted V-shaped trend (Lei *et al.* 2019; Li *et al.* 2019; Wang *et al.* 2020). In recent years, particulate matter pollution has significantly decreased, but the pollution of O₃ has worsened. According to the China Environment Report 2019 (available at <https://www.mee.gov.cn>, in Chinese), 30% of the 338 cities in China have O₃ concentrations exceeding the secondary limit of Environmental Air Quality Standards (GB3095-2012) (160 µg/m³ [101.325 kPa, 20 °C]), especially in the Beijing-Tianjin-Hebei and Yangtze River Delta regions (Wei *et al.* 2017; Zhang *et al.* 2015).

With rapid industrialization and urbanization, O₃ has become a major air pollutant in China (Al-Jassim *et al.* 2018; Lu *et al.* 2018). Due to its low water solubility, it easily enters the respiratory tract; therefore, it has a strong stimulating effect and undergoes strong oxidation (Nuvolone *et al.* 2017; Zhang *et al.* 2019b). The respiratory

system is the first part of the human body to receive O₃ following inhalation and, therefore, suffers the most obvious effects of O₃ exposure. Epidemiological studies have shown a significant positive correlation between acute O₃ exposure and decreased lung function in both average and susceptible individuals, especially in seasons with high O₃ concentrations (Adams & William 2002, 2003, 2006). According to several epidemiological studies, Long- and short-term exposure to high concentrations of O₃ can lead to respiratory diseases such as infections and asthma, cardiovascular diseases such as stroke and arrhythmias, and neurological diseases such as autism and Alzheimer's disease in children (Bell *et al.* 2005; Dominici *et al.* 2006; Pope & Dockery, 2006). O₃ exposure is an important factor that can lead to premature death in humans. It can activate a large number of inflammatory mediators in the respiratory system, leading to the accumulation of toxic lipid oxidation products and eventually chronic inflammation (Canella *et al.* 2016). In addition, it can produce strongly oxidizing free radicals in the human body, disrupt cell metabolism, accelerate senescence, induce chromosomal lesions in lymphocytes, and damage the immune system (Rider & Carlsten, 2019). The exposure of certain groups to high concentrations of O₃, such as pregnant women, infants, and young children, may cause serious health threats and even increase the risk of mortality (Silva *et al.* 2013). A review that O₃ pollution trends and health impacts in China is therefore needed to aid in formulating a mitigation policy and to guide future research.

Specifically, this study aims to evaluate the current state of O₃ pollution in mainland Chinese cities on an annual basis from 2013 to 2021 and review O₃ pollution's impact on population health. The significance of this study is twofold. Firstly, it elucidates the prevailing situation and trends of O₃ pollution in mainland China over the past decade, with the aim of drawing attention to and prompting further research on this issue. Secondly, the study can aid in understanding the health effects of O₃ and support the formulation of O₃ standards in China.

2. Current scenario of O₃ pollution in China

Since the 1990s, with the increase in the number of motor vehicles, there has been a parallel increase in coal, oil, and other forms of energy consumption, such that compound

Table 1. Mean value of O₃-8h concentration and 90th percentile range in some Chinese cities from 2013 to 2021

Year	Number of cities (number)	Average concentration of O ₃ -8h (µg/m ³)	Range of 90 th percentile of O ₃ -8h concentration (µg/m ³)	Over-standard rate (%)
2013	74	139	72–190	–
2014	161	140	69–210	6.1
2015	338	134	62–203	4.6
2016	338	138	73–200	5.2
2017	338	149	78–218	7.6
2018	338	151	76–217	8.4
2019	338	148	—	7.6
2020	338	138	—	4.9
2021	338	137	—	4.4

Note: —Statistical analysis of the 74 cities should refer to the year 2013. (The data come from the Ministry of Ecology and Environment of the People's Republic of China, available at <https://www.mee.gov.cn>).

air pollution has replaced the typical soot-type pollution in China. With the gradual implementation of the government's environmental optimization policy, particulate matter pollution has significantly reduced, although O₃ pollution is becoming increasingly serious. In recent years, O₃ concentrations exceeding the Chinese standard have become increasingly common.

According to the bulletin on the state of China's environment issued by the Ministry of Ecology and Environment from 2013 to 2021, China has set up national air monitoring stations in 74 cities in the regions of Beijing, Tianjin, Hebei, Yangtze River Delta, Pearl River Delta, and in municipalities directly under the Central Government, as well as in provincial capitals and cities separately listed on the State plan, to carry out air monitoring in accordance with the new Environmental Air Quality Standards (GB3095-2012). The 2013–2021 average concentration of O₃-8h (Daily maximum 8h average) and the 90th percentile range in Chinese cities are shown in Table 1, and the changing trends are shown in Figure. 1.




Figure 1. The changing trend of O₃ concentration in the key regions of China (Beijing-Tianjin-Hebei and Yangtze River Delta) and the entire country from 2013 to 2021

Overall, the O₃ concentrations in the Beijing-Tianjin-Hebei region, Yangtze River Delta, Pearl River Delta, and Weihe Plain were higher than those in other areas, as shown in Table 2. The concentration of O₃ in the Beijing-Tianjin-Hebei region was the highest, with an average annual

concentration of 176.67 $\mu\text{g}/\text{m}^3$, and the proportion of over- Environmental Air Quality Standard (GB3095-2012) for O₃ in the Pearl River Delta was the largest, with was 57.33%.

Table 2 O₃ pollution in the key regions since 2013

Year	Index	BTH	YRD	PRD	WP
2013	O ₃ mean concentration ($\mu\text{g}/\text{m}^3$)	155	144	155	—
	O ₃ exceedance ratio (%)	7.6	13.9	31.9	—
2014	O ₃ mean concentration ($\mu\text{g}/\text{m}^3$)	162	154	156	—
	O ₃ exceedance ratio (%)	—	—	—	—
2015	O ₃ mean concentration ($\mu\text{g}/\text{m}^3$)	162	163	145	—
	O ₃ exceedance ratio (%)	17.2	37.2	56.5	—
2016	O ₃ mean concentration ($\mu\text{g}/\text{m}^3$)	172	159	151	—
	O ₃ exceedance ratio (%)	26.3	39.8	70.3	—
2017	O ₃ mean concentration ($\mu\text{g}/\text{m}^3$)	193	170	165	—
	O ₃ exceedance ratio (%)	41.0	50.4	70.6	—
2018	O ₃ mean concentration ($\mu\text{g}/\text{m}^3$)	199	167	—	180
	O ₃ exceedance ratio (%)	46.0	49.3	—	36.4
2019	O ₃ mean concentration ($\mu\text{g}/\text{m}^3$)	196	164	—	171
	O ₃ exceedance ratio (%)	48.2	49.5	—	37.6
2020	O ₃ mean concentration ($\mu\text{g}/\text{m}^3$)	180	152	—	161
	O ₃ exceedance ratio (%)	46.6	50.7	—	36.1
2021	O ₃ mean concentration ($\mu\text{g}/\text{m}^3$)	171	151	—	165
	O ₃ exceedance ratio (%)	41.8	55.4	—	39.3

Note: BTH, Beijing-Tianjin-Hebei; YRD, Yangtze River Delta; PRD, Pearl River Delta; WP, Weihe Plain.

O₃ is primarily formed through the photochemical reactions of NO_x and VOCs (Sillman, 1999), which mainly stem from biomass fuel combustion, urban construction, automobile exhaust, and natural sources (Xu *et al.* 2019; Zong *et al.* 2018). The concentration of O₃ in the economically developed eastern part of China is higher than in other regions (Peng *et al.* 2017). Kamal *et al.* (2019) reported that O₃ concentrations in China range from 74 to 201 $\mu\text{g}/\text{m}^3$. Approximately 30% of the population experiences O₃ levels exceeding 160 $\mu\text{g}/\text{m}^3$, and approximately 67.2% of the population is exposed to an O₃ environment greater than 100 $\mu\text{g}/\text{m}^3$ (Kamal *et al.* 2019). The comprehensive control and monitoring of O₃ in China began relatively late. Since 2012, the Ministry of Ecology and Environment of the People's Republic of China has been listing O₃ as an environmental air pollutant. In 2013, large-scale air monitoring stations were built; in recent years, air monitoring stations have been gradually installed in the entire country. As shown in Table 1 and Figure. 2, before 2018, the over-standard rate of O₃ increased annually; after 2018, the over-standard rate of O₃ also improved significantly as the government issued a series of targeted control measures on atmospheric O₃ pollution.

3. An epidemiological study of the effect of O₃ on human health

Since the smog events in London and Los Angeles in the last century, increasing attention has been paid to air pollution, and researchers from various countries have begun to study the effects of O₃ exposure on human health. These effects include an increase in mortality and

morbidity, and a decrease in lung function (Huangfu & Atkinson, 2020; Li *et al.* 2020). A Chinese multi-city study found that when the concentration of O₃-8h increased by 10 $\mu\text{g}/\text{m}^3$, the total risks of mortality, cardiovascular disease, hypertension, coronary artery disease, and stroke mortality increased by 0.23%, 0.27%, 0.60%, 0.24%, and 0.29%, respectively (Peng *et al.* 2017). The main effects of O₃ exposure on the respiratory and cardiovascular systems and potential mechanisms of its impact on human health are detailed in the following paragraphs.




Figure 2. Ozone levels exceeding the standard rate in some Chinese cities, from 2013 to 2021 (Note in the data come from the Ministry of Ecology and Environment of the People's Republic of China, available at <https://www.mee.gov.cn>)

3.1. Effects of O₃ on the respiratory system

The respiratory tract is the primary route through which air pollutants enter the human body. Most respiratory diseases involve airway lesions, including airway inflammation, remodeling, changes in responsiveness, and

decreased host resistance to infection (Thurston *et al.* 2016). The forced expiratory volume in 1 sec (FEV1) is an important index for studying the effect of O₃ exposure on the respiratory system. O₃ concentration is usually controlled at 40–600 ppb in a study population of healthy non-smoking adults (McDonnell *et al.* 2012). A study of 30 healthy young people reported that FEV1 decreased significantly with O₃ exposure at 0.08 ppm, compared with 0.06 ppm. In studies conducted under similar exposure conditions (0.06 ppm exposure for 6.6 h), Brown *et al.* (2008) discovered that O₃ exposure led to an average 2.85% decrease in FEV1 (Brown *et al.* 2008). Concurrently, Chong *et al.* (2011) reported a 1.71% decrease in FEV1 expression (Chong *et al.* 2011). Furthermore, in a separate study, FEV1 decreased by 2.72% with 0.060 ppm of O₃, 5.34% with 0.070 ppm, 7.02% with 0.080 ppm, and as much as 11.42% with 0.087 ppm (Schelegle *et al.* 2009).

In summary, there was a significant negative correlation between O₃ concentration and pulmonary function and a positive correlation between O₃ concentration and number of hospitalizations and visits to the emergency department, especially in the hot season (Darrow *et al.* 2011; Stieb *et al.* 2009; Strickland *et al.* 2010). Studies at home and abroad have found a positive correlation between O₃ exposure and hospital admission for respiratory diseases (Cakmak *et al.* 2006; Dales *et al.* 2006; Mercedes *et al.* 2006; Silverman & Ito, 2010; Wong *et al.* 2010; Yang *et al.* 2005).

Recent studies have focused on the relationship between O₃ exposure and respiratory mortality. Single- and multi-city studies have found that there is a significant positive correlation between O₃ exposure and respiratory mortality (Bell & Michelle, 2004; Lei *et al.* 2019; Li *et al.* 2020; Peng *et al.* 2017; Wu *et al.* 2019). These studies also found that the effect of O₃ on the respiratory system exhibited lag and seasonal effects. Shang *et al.* (2013) conducted a meta-analysis, and the results showed that when the O₃ concentration increased by 10 µg/m³, the mortality rate of respiratory diseases increased by 0.73% (95% CI: 0.49–0.97%) (Shang *et al.* 2013).

3.2. Effects of O₃ on the cardiovascular system

Cardiovascular disease poses one of the most important public health problems in our country. O₃ exposure can lead to changes in heart rate and a decrease in heart rate variability has been identified as a predictor of increased cardiovascular morbidity and mortality. Studies by Liao *et al.* (2004) and others demonstrated a relationship between O₃ exposure and cardiovascular diseases (Liao *et al.* 2004; Park *et al.* 2005; Ruidavets & J.-B., 2005). It has also been reported that short-term O₃ exposure has no effect on the vascular function, blood pressure, or heart rate (Barath *et al.* 2013; Hoffmann *et al.* 2012). In a study of veterans, it was found that for every 2.6 µg/m³ increase in O₃ concentration, heart rate variability decreased by 11.5% (95% CI: 0.4–21.3%), and this effect was more evident in men with hypertension and ischemic heart disease than in men with good heart health (Park *et al.* 2005).

Some studies have shown that O₃ exposure has a greater impact on cardiovascular mortality in aging populations as compared to young ones, and the effect of O₃ is stronger in summer (Samoli *et al.* 2009). Studies carried out in the United States and China found different impacts of O₃ on the health of populations.

A study of 48 cities in the United States found that each 10 ppb increase in the concentration of O₃ increased cardiovascular mortality by 0.5% (95% CI: 0.30–0.60%) (Zanobetti *et al.* 2008), while another study of 96 US cities showed an increase in cardiovascular mortality by 1.09% (95% CI: 0.30–2.25%) (Jerrett *et al.* 2009). Li *et al.*'s research in Guangzhou showed that every 10 µg/m³ increase in O₃ concentration increased the mortality rate of cardiovascular disease by 0.59% (95% CI: 0.30–0.88%) (Li *et al.* 2021), whereas Zhang *et al.*'s study in Nanjing demonstrated that the mortality rate of cardiovascular disease increased by 0.98% (95% CI: 0.59–1.38%) (Zhang *et al.* 2019b). Therefore, overall, O₃ exposure has a greater impact on the health of Chinese residents compared with that of United States residents.

3.3. Mechanisms underlying the effects of O₃ on human health

O₃ is ubiquitous and has low water solubility, allowing it to easily enter deep into the respiratory tract, where it has a strong stimulating effect and strong oxidation. More than 80% of the O₃ absorbed by humans is inhaled through respiratory tract, where it can cause injury. Studies have shown that O₃ in certain concentrations reacts with cells in the respiratory tract and leads to increase in airway inflammation (Wu *et al.* 2011). Related studies have shown that O₃ can cause epithelial cells to release reactive oxygen species (ROS) and prostaglandin E₂ (PGE₂), activate the NF-κB signaling pathway, initiate the transcription of pro-inflammatory factors such as interleukin-8 (IL-8), and cause neutrophils to gather in the bronchi, leading to bronchial inflammation and injury. The effect of O₃ is aggravated when combined with the synergistic action of particulate fine particulate matter (Damera *et al.* 2009; Li *et al.* 2013; Sunil *et al.* 2013; Wu *et al.* 2011). Researchers have found that O₃ can also activate tyrosine kinases, promote the phosphorylation of epithelial cells, and upregulate the expression of IL-8 in the bronchi (Weidong *et al.* 2015).

O₃ exposure can cause oxidative decomposition, peroxidative modification, and peroxidation of lipids in the body, resulting in potential lipid peroxidation. Peroxidation increases aging-related risks and cancer, while ROS stimulate the rapid division of tumor cells (Wang *et al.*). In vitro studies have demonstrated that O₃ can stimulate alveolar cells to release ROS and induce cell membrane lipid peroxidation, leading to a mixture of ROS and lipid ozonation products (Kadiiska *et al.* 2013). This can result in damage to the lungs and other organs. Some researchers have found that O₃ exposure can also lead to the destruction of spirochete DNA structures, induce DNA mutations, and inhibit DNA replication. Additionally, O₃ can induce systemic effects by regulating the activation of the neurohormonal stress response pathway.

3.4. Comparison of the effects of O₃ exposure on human health among countries worldwide

Globally, O₃ exposure is believed to have contributed to 250,000 premature deaths in 2015 (Lin *et al.* 2018). A study that examined O₃ levels and their health consequences in 50 cities in the eastern United States determined that higher O₃ levels were associated with an increase in total daily mortality of approximately 0.11–0.27% (Bell *et al.* 2007). Moreover, studies have indicated that in the United States, elevated O₃ levels due to climate change may result in an additional 50 premature deaths per year nationwide (Stowell *et al.* 2017). Future O₃ levels in Fennoscandia and the northern United Kingdom are predicted to exceed the World Health Organization Global Air Quality Guidelines level of 50 ppb. Consequently, by 2050, it is anticipated that acute effects of O₃ on human health will diminish in the majority of Fennoscandia, but chronic effects are likely to persist or perhaps worsen (Karlsson *et al.* 2017). In Athens, Greece, long-term exposure to O₃ was identified as the main cause of reduced life expectancy, while short-term exposure to O₃ was not found to have the same impact; the relevant characterization factors for human health damage suggested that the impact of short-term exposure to O₃ ranged between 1.58×10^{-7} and 4.71×10^{-7} years of life lost (Kassomenos *et al.* 2013). Surface O₃ levels in Delhi, India have been measured at concentrations that are

considerably higher than hazardous levels, exceed the exposure threshold for human health for up to 45 days annually, and far exceed the European Union directive's maximum of 25 days annually (Ghude *et al.* 2009). There is strong evidence that there is a positive correlation between daily O₃ concentrations that exceed existing regulatory requirements and daily non-accidental death (Bell *et al.* 2007; Karlsson *et al.* 2017; Nuvolone *et al.* 2017; Stowell *et al.* 2017; Zhang *et al.* 2019).

4. Ozone exposure assessment model

Typical studies of the atmospheric O₃ take the concentration exposure levels from the monitoring station to be those of the population, which could lead to serious measurement errors.

With the continued development of computer technology and geographic information, global positioning, and remote sensing systems, air pollutant exposure assessment models have diversified. At present, the exposure assessment models used by researchers mainly include proximity, spatial interpolation, land-use regression, atmospheric diffusion, and Bayesian spatiotemporal models. These statistical models provide a more precise assessment of air pollutants.

The parameters used by each exposure assessment model are different, with varying advantages and disadvantages, as shown in Table 3 below.

Table 3. Principles, advantages, and disadvantages of various exposure assessment models

Exposure model	Principle	Advantages	Disadvantages
Proximity model (Andersson <i>et al.</i> 2011)	Assign the parameters needed by the research object to the coordinates of the geographic information system (GIS), combined with the environmental variables in the study area, compare the distance between the research site and the air pollution source, and evaluate the impact of the pollution source on the health of the research object	Easy to operate and less expensive, and can also use geographic information systems	Only qualitative analyses can be carried out. The results are subject to the influence of environmental contaminants' sources and concentration distributions, and there is a certain degree of report deviation.
Spatial interpolation model (Jin & Heap, 2011)	Some geographic information statistical techniques are used to infer the concentration of pollutants in the study area by obtaining pollution data from monitoring points around the area	This model uses the monitoring points to obtain the temporal and spatial changes of pollutants	Because of the large consumption of funds, professional and technical personnel are needed, and the monitoring data have errors and great uncertainty.
Land use regression model (Alexeeff <i>et al.</i> 2015; Morley & Gulliver, 2018)	The concentration of pollutants measured by the air quality monitoring station is used as the dependent variable, and geographical variables such as land use, traffic roads, topography, and population distribution around the monitoring station are predictive variables to establish a regression model. Thus, the regression model is used to predict the concentration of pollutants at any spatial site in the study area	Compared with the interpolation model, this model has a lower economic cost and more accurate prediction of atmospheric pollutant concentrations	Cannot effectively distinguish the influence of major pollutants, struggles to predict the subtle changes of air pollutants, and is easily affected by confounding factors
Atmospheric diffusion model (Sørensen <i>et al.</i> 2012)	According to the Gaussian equation, the temporal and spatial exposure assessment of pollutant concentration is carried out by using pollutant emission, meteorological, and topographic data	It can combine the changes of air pollution in time and space without intensive monitoring networks, and has higher resolution	Hardware and software equipment is expensive and professional operators need to be trained; monitoring data need to be cross-checked with each other; temporary mismatch or misclassification of data may cause estimation offset

Bayesian space-time mode (Szpiro <i>et al.</i> 2010)	Based on local geographical, meteorological, and station monitoring data, pollutant concentration in a certain area is divided into several time-space domains. When fully considering various possible uncertain factors, the indoor pollutant monitoring data of the outdoor concentration of each study object are obtained by simulation. Then, combined with the characteristics of the house and the permeability coefficient, the indoor concentration of the house is simulated, and according to the time ratio of indoor and outdoor activities, the exposure concentration of each object is measured by weight	Bayesian models have higher inference accuracy, acceptable spatiotemporal interaction, and over-discretization in small sample data, which overcomes the limitations of classical statistical methods, such as the needs of randomness, independence, uniform distribution, linearity, and isotropy	Excessive smooth processing, difficultly exploring complex spatiotemporal data information
--	--	---	--

5. Summary and recommendations

This paper evaluates ground-level O₃ concentrations in major urban areas across China and reviews research findings on the health effects of O₃ exposure. This includes discussions on the pathogenic mechanisms of O₃ and the evaluation of O₃ exposure models. We propose the following conclusions and recommendations:

(1) Following the implementation of regulations aimed at curbing particulate matter and other forms of air pollution in 2013, China has seen a consistent increase in surface O₃ levels, which only began to decelerate in 2018. The available data unequivocally indicate that O₃ pollution is a serious issue in China's major cities, with concentrations in the Beijing-Tianjin-Hebei region, Yangtze River Delta, Pearl River Delta, and Weihe Plain surpassing those in other areas. China has initiated a monitoring system for O₃ and its precursors that can assist with O₃ pollution prevention and control measures. This system provides a comprehensive understanding of O₃ pollution across China. However, the quality control and assurance of O₃ and its precursors' observations need further enhancement to furnish robust scientific and technological support for O₃ pollution prevention and control efforts.

(2) O₃ pollution is a global health concern. As previously mentioned, O₃ concentrations are projected to rise in many parts of the world, potentially increasing ozone-related mortality and morbidity rates. Due to the current uncertainty surrounding the cardiovascular effects of O₃, the overall impact could be even more substantial when considering other potential effects of O₃ exposure. At present, there are many studies on the relationship between O₃ and human health in the developed areas of China, but there are few studies focusing on the central and western regions, and the spatial representation is insufficient. Most of the O₃ exposure-response curves are complex linear, which makes the study of O₃ health effects very challenging. Moreover, studies often focus on populations, so it is difficult to accurately evaluate the level of individual O₃ exposure. There are variations in models and parameters used among studies, such that the results differ; therefore, choosing the correct exposure coefficient is difficult. Future research should focus on the following aspects:

- Carrying out additional cohort studies in a large population
- Studying the effects of combined exposure of two or more pollutants on population health
- Carrying out research in different geographic areas to obtain sufficient spatial representation
- Determining the minimum human threshold for O₃ pollution, evaluating its effects on the human body, and further exploring its underlying mechanisms.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

All authors consent for the publication of the manuscript and the materials incorporated.

Competing interests

The author declares no competing interests.

References

- Adams and William C. (2002). Comparison of chamber and face-mask 6.6-hour exposures to ozone on pulmonary function and symptoms responses. *Inhalation Toxicology*, **14**(7), 745–764.
- Adams and William C. (2003). Comparison of Chamber and Face Mask 6.6-Hour Exposure to 0.08 ppm Ozone Via Square-Wave and Triangular Profiles on Pulmonary Responses. *Inhalation Toxicology*, **15**(3), 265–281.
- Adams and William C. (2006). Comparison of Chamber 6.6-h Exposures to 0.04–0.08 PPM Ozone via Square-wave and Triangular Profiles on Pulmonary Responses. *Inhalation Toxicology*, **18**(2), 127–136.
- Alexeeff S.E., Schwartz J., Kloog I., Chudnovsky A., Koutrakis P. and Coull B.A. (2015). Consequences of kriging and land use regression for PM_{2.5} predictions in epidemiologic analyses: insights into spatial variability using high-resolution satellite data. *Journal of Exposure Science and Environmental Epidemiology*, **25**(2), 138–144.
- Andersson M., Modig L., Hedman L. and Rönmark B.F. (2011). Heavy vehicle traffic is related to wheeze among schoolchildren: a population-based study in an area with low traffic flows. *Environmental Health*, **10**, 91.

- Al-Jassim N., Mantilla-Calderon D., Scarascia D and P.-Y. (2018). *Hong Environmental Science & Technology*, **52**(24), 14331–14341
- Barath S., Langrish J.P., Lundback M., Bosson J.A., Goudie C., Newby D.E., Sandstrom T., Mills N.L. and Blomberg A. (2013). Short-Term Exposure to Ozone Does Not Impair Vascular Function or Affect Heart Rate Variability in Healthy Young Men. *Toxicological Sciences*, **135**(2), 292–299.
- Bell and Michelle L. (2004). Ozone and short-term mortality in 95 US urban communities, 1987–2000. *Jama*, **292**(19), 2372–2378.
- Bell M.L., Dominici F. and Samet J.M. (2005). A meta-analysis of time-series studies of ozone and mortality with comparison to the national morbidity, mortality, and air pollution study *Epidemiology*, **16**(4), 436–445.
- Bell M.L., Goldberg R., Hogrefe C., Kinney P.L., Knowlton K., Lynn B., Rosenthal J., Rosenzweig C. and Patz J.A. (2007). Climate change, ambient ozone, and health in 50 US cities. *Climatic Change*, **82**(1–2), 61–76.
- Brown J.S., Bateson T.F. and McDonnell W.F. (2008). Effects of Exposure to 0.06 ppm Ozone on FEV1 in Humans: A Secondary Analysis of Existing Data. *Environmental Health Perspectives*, **116**(8), 1023–1026.
- Cakmak S., Dales R.E. and Judek S. (2006). Respiratory Health Effects of Air Pollution Gases: Modification by Education and Income. *Archives of Environmental & Occupational Health*, **61**(1), 510.
- Canella R., Borriello R., Cavicchio C., Cervellati F., Martini M., Muresan X. and Valacchi G. (2016). Tropospheric ozone effects on chlorine current in lung epithelial cells: an electrophysiological approach. *Free Radical Biology and Medicine*, **96**, S58–S59.
- Chong S.K., Alexis N.E., Rappold A.G., Kehrl H., Hazucha M.J., Lay J.C., Schmitt M.T., Case M., Devlin R.B. and Peden D.B. (2011). Lung Function and Inflammatory Responses in Healthy Young Adults Exposed to 0.06 ppm Ozone for 6.6 Hours. *American Journal of Respiratory & Critical Care Medicine*, **183**(9), 1215.
- Dales R.E., Cakmak S. and Doiron M.S. (2006). Gaseous Air Pollutants and Hospitalization for Respiratory Disease in the Neonatal Period. *Environmental Health Perspectives*, **114**(11), 1751–1754.
- Damera G., Zhao H., Wang M., Smith M., Kirby C., Jester W.F., Lawson J.A. and Panettieri R.A. (2009). Ozone modulates IL-6 secretion in human airway epithelial and smooth muscle cells. *American Journal of Physiology-Lung Cellular and Molecular Physiology*, **296**(4).
- Darrow L.A., Klein M., Sarnat J.A., Mulholland J.A. and Tolbert P.E. (2011). The use of alternative pollutant metrics in time-series studies of ambient air pollution and respiratory emergency department visits. *Journal of Exposure Science & Environmental Epidemiology*, **21**(1), 10–19.
- Dominici Francesca, Peng Roger D., Bell Michelle L., Pham and Luu. (2006). Fine Particulate Air Pollution and Hospital Admission for Cardiovascular and Respiratory Diseases. *JAMA: Journal of the American Medical Association*, **295**(10), 1127–1134.
- Ghude S.D., Jain S.L., Arya B.C., Beig G., Ahammed Y.N., Kumar A. and Tyagi B. (2009). Ozone in ambient air at a tropical megacity, Delhi: characteristics, trends and cumulative ozone exposure indices. *Journal of Atmospheric Chemistry*, **60**(3), 237–252.
- Guan W.J., Zheng X.Y., Chung K.F. and Zhong N.S. (2016). Impact of air pollution on the burden of chronic respiratory diseases in China: time for urgent action. *The Lancet*, **388**, 1939–1951.
- Hoffmann B., Luttmann-Gibson H., Cohen A., Zanobetti A., Souza C.D., Foley C., Suh H.H., Coull B.A., Schwartz J. and Mittleman M. (2012). Opposing Effects of Particle Pollution, Ozone, and Ambient Temperature on Arterial Blood Pressure. *Environmental Health Perspectives*, **120**(2), 241–246.
- Huangfu P. and Atkinson R. (2020). Long-term exposure to NO₂ and O₃ and all-cause and respiratory mortality: A systematic review and meta-analysis. *Environment International*, **144**(2).
- Jerrett M., Burnett R.T., Pope C.A., Ito K., Thurston G., Krewski D., Shi Y., alle E.C. and Thun M. (2009). Long-term ozone exposure and mortality. *New England Journal of Medicine*, **360**(11), 1085–1095.
- Jin L. and Heap A.D. (2011). A review of comparative studies of spatial interpolation methods in environmental sciences: Performance and impact factors. *Ecological Informatics*, **6**(3–4), 228–241.
- Kadiiska M.B., Basu S., Brot N., Cooper C., Csallany A.S., Davies M.J., George M.M., Murray D.M., Roberts L.J. and Shigenaga M.K. (2013). Biomarkers of oxidative stress study V: Ozone exposure of rats and its effect on lipids, proteins, and DNA in plasma and urine. *Free Radical Biology & Medicine*, **61**(8), 408–415.
- Karlsson P.E., Klingberg J., Engardt M., Andersson C., Langner J., Karlsson G.P. and Pleijel H. (2017). Past, present and future concentrations of ground-level ozone and potential impacts on ecosystems and human health in northern Europe. *Science of The Total Environment*, **576**, 22–35.
- Kassomenos P.A., Dimitriou K. and Paschalidou A.K. (2013). Human health damage caused by particulate matter PM₁₀ and ozone in urban environments: the case of Athens, Greece. *Environmental Monitoring and Assessment*, **185**(8), 6933–6942.
- Kamal, Jyoti, Maji, Wei-Feng, Ye Mohit, Arora and Shiva. (2019). Ozone pollution in Chinese cities: Assessment of seasonal variation, health effects and economic burden. *Environmental pollution*, **247**, 792–801.
- Lei R., Zhu F., Cheng H., Liu J., Shen C., Zhang C., Xu Y., Xiao C., Li X., Zhang J., Ding R. and Cao J. (2019). Short-term effect of PM_{2.5}/O₃ on non-accidental and respiratory deaths in highly polluted area of China. *Atmospheric Pollution Research*, **10**(5), 1412–1419.
- Li F., Wiegman C., Seiffert J.M., Zhu J., Clarke C., Chang Y., Bhavsar P., Adcock I., Zhang J. and Xin Z. (2013). Effects of N-Acetylcysteine in Ozone-Induced Chronic Obstructive Pulmonary Disease Model. *Plos One*, **8**(11), e80782.
- Li K., Jacob D.J., Liao H., Shen L., Zhang Q. and Bates K.H. (2019). Anthropogenic drivers of 2013–2017 trends in summer surface ozone in China. *Proceedings of the National Academy of Sciences*, **116**(2), 422–427.
- Li M., Dong H., Wang B., Zhao W., Sakhvidi M.J.Z., Li L., Lin G. and Yang J. (2021). Association between ambient ozone pollution and mortality from a spectrum of causes in Guangzhou, China. *Science of the Total Environment*, **754**, Article 142110.
- Li T., Chen R., Zhang Y., Fang J., Zhao F., Chen C., Wang J., Du P., Wang Q., Shi W., Han J., Hu X., Kan H. and Shi X. (2020). Cohort profile: Sub-clinical outcomes of polluted air in China

- (SCOPA-China cohort). *Environment International*, **134**, Article 105221.
- Liao D., Duan Y., Whitsel E., Zhi-Jie Z., Gerardo H., Chinchilli V.M. and Lin H.M. (2004). Association of Higher Levels of Ambient Criteria Pollutants with Impaired Cardiac Autonomic Control: A Population-based Study. *American Journal of Epidemiology*, **(8)**, 768–777.
- Liu Z., Guan D., Wei W., Davis S.J., Ciais P., Bai J., Peng S., Zhang Q., Hubacek K., Marland G., Andres R.J., Crawford-Brown D., Lin J., Zhao H., Hong C., Boden T.A., Feng K., Peters G.P., Xi F. and He K. (2015). Reduced carbon emission estimates from fossil fuel combustion and cement production in China. *Nature*, **524**(7565), 335–338.
- Lin Y., Jiang F., Zhao J., Zhu G., He X., Ma X., Li S., Sabel C.E. and Wang H. (2018). Impacts of O₃ on premature mortality and crop yield loss across China. *Atmospheric Environment* **194**, 41–47.
- Lu X., Hong J., Zhang L., Cooper O.R., Schultz M.G., Xu X., Wang T., Gao M., Zhao Y. and Zhang Y. (2018). Severe Surface Ozone Pollution in China: A Global Perspective. *Environmental Science & Technology*, 487–494.
- Mcdonnell W.F., Stewart P.W., Smith M.V., Kim C.S. and Schelegle E.S. (2012). Prediction of lung function response for populations exposed to a wide range of ozone conditions. *Inhalation Toxicology*, **24**(10), 619–633.
- Mercedes, M.R., Antonella Z. and Joel S. (2006). The Effect of Ozone and PM₁₀ on Hospital Admissions for Pneumonia and Chronic Obstructive Pulmonary Disease: A National Multicity Study. *American Journal of Epidemiology*, **(6)**, 579–588. <https://doi.org/10.1093/aje/kwj078>
- Morley D.W. and Gulliver J. (2018). A land use regression variable generation, modelling and prediction tool for air pollution exposure assessment. *Environmental Modelling & Software*, **105**, 17–23.
- Nuvolone D., Petri D. and Voller F. (2017). The effects of ozone on human health. *Environmental Science and Pollution Research*, **25**(9), 8074–8088.
- Park S.K., O'Neill M.S., Vokonas P.S., Sparrow D. and Schwartz J. (2005). Effects of Air Pollution on Heart Rate Variability: The VA Normative Aging Study. *Environmental Health Perspectives*, **113**.
- Peng Y., Renjie C., Lijun W., Xia M., Cong L., Yue N., Zhijing L., Yunning L., Jiangmei L., Jinlei Q., Jinling Y., Maigeng Z. and Haidong K. (2017). Ambient Ozone Pollution and Daily Mortality: A Nationwide Study in 272 Chinese Cities. *Environmental Health Perspectives*, **125**(11).
- Pope C. and Dockery D. (2006). Health effects of fine particulate air pollution: lines that connect. *Journal of the Air & Waste Management Association*, **56**(6), 707–708.
- Rider C.F. and Carlsten C. (2019). Air pollution and DNA methylation: effects of exposure in humans. *Clin Epigenetics*, **11**(1), 131.
- Ruidavets and J.-B. (2005). Ozone Air Pollution Is Associated with Acute Myocardial Infarction. *Circulation*, **111**(5), 563–569.
- Samolli E., Zanobetti A., Schwartz J., Atkinson R., Letertre A., Schindler C., Perez L., Cadum E., Pekkanen J. and Paldy A. (2009). The temporal pattern of mortality responses to ambient ozone in the APHEA project. *Journal Epidemiol Community Health*, **63**(12), 960–966.
- Schelegle E.S., Morales C.A., Walby W.F., Marion S. and Allen R.P. (2009). 6.6-hour inhalation of ozone concentrations from 60 to 87 parts per billion in healthy humans. *American Journal of Respiratory & Critical Care Medicine*, **180**(3), 265–272.
- Shang Y., Sun Z., Cao J., Wang X., Zhong L., Bi X., Li H., Liu W., Zhu T. and Huang W. (2013). Systematic review of Chinese studies of short-term exposure to air pollution and daily mortality. *Environment International*, **54**, 100–111.
- Sillman S. (1999). The relation between ozone, NO_x and hydrocarbons in urban and polluted rural environments. *Atmospheric Environment*, **33**(12), 1821–1845
- Silva R.A., West J.J., Zhang Y., Anenberg S.C., Lamarque J.F.O., Shindell D.T., Collins W.J., Dalsoren S., Faluvegi G. and Folberth G. (2013). Global premature mortality due to anthropogenic outdoor air pollution and the contribution of past climate change. *Environmental Research Letters*, **8**(3), 034005.
- Silverman R.A. and Ito K. (2010). Age-related association of fine particles and ozone with severe acute asthma in New York City. *Journal of Allergy & Clinical Immunology*, **125**(2), 367–373.
- Sørensen M., Hoffmann B., Hvidberg M., Ketzel M., Jensen S. S., Andersen Z.J., Tjønneland A., Overvad K. and Raaschou-Nielsen O. (2012). Long-Term Exposure to Traffic-Related Air Pollution Associated with Blood Pressure and Self-Reported Hypertension in a Danish Cohort. *Environmental Health Perspectives*, **120**, 3.
- Stieb D.M., Szyszkowicz M., Rowe B.H. and Leech J.A. (2009). Air pollution and emergency department visits for cardiac and respiratory conditions: a multi-city time-series analysis. *Environmental Health*, **8**(1), 25.
- Stowell J.D., Kim Y.-m., Gao Y., Fu J.S., Chang H.H. and Liu Y. (2017). The impact of climate change and emissions control on future ozone levels: Implications for human health. *Environment International*, **108**, 41–50.
- Strickland M.J., Darrow L.A., Klein M., Flanders W.D., Sarnat J.A., Waller L.A., Sarnat S.E., Mulholland J.A. and Tolbert P.E. (2010). Short-term associations between ambient air pollutants and pediatric asthma emergency department visits. *American Journal of Respiratory & Critical Care Medicine*, **182**(3), 307.
- Sunil Vasanthi R., Vayas Kinal N., Massa Christopher B. and Gow. (2013). Ozone-Induced Injury and Oxidative Stress in Bronchiolar Epithelium Are Associated with Altered Pulmonary Mechanics. *Toxicological Sciences*, **133**(2), 309–319.
- Szpiro A.A., Sampson P.D., Sheppard L., Lumley T., Adar S.D. and Kaufman J.D. (2010). Predicting intra-urban variation in air pollution concentrations with complex spatio-temporal dependencies. *Environmetrics*, **21**(6), 606–631.
- Thurston G.D., Kipen H., Annesi-Maesano I., Balmes J. and Brunekreef B. (2016). A joint ERS/ATS policy statement: what constitutes an adverse health effect of air pollution? An analytical framework. *European Respiratory Journal*, **49**(1), 1600419.
- Wang L., Wise J., Zhang Z. and Shi X. Progress and prospects of reactive oxygen species in metal carcinogenesis. *Current pharmacology reports*, **2**(4), 178–186.
- Wang Y., Gao W., Wang S., Song T., Gong Z., Ji D., Wang L., Liu Z., Tang G. and Huo Y. (2020). Contrasting trends of PM_{2.5} and surface ozone concentrations in China from 2013 to 2017. *National Science Review*, **7**(8), 1331–1339.

- Wei G., Tie X., Xu J., Huang R., Mao X., Zhou G. and Chang L. (2017). Long-term trend of O₃ in a mega City (Shanghai), China: Characteristics, causes, and interactions with precursors. *Science of the Total Environment*, 603–604(dec.15), 425–433.
- Weidong Wu, Phillip Wages, Robert, Devlin, David and Diaz-Sanchez. (2015). SRC-mediated EGF receptor activation regulates ozone-induced interleukin 8 expression in human bronchial epithelial cells. *Environmental Health Perspectives*, **123**, 3.
- Wong C., Vichitvadakan N., Vajanapoom N., Ostro B., Thach T., Chau P., Chan E., Chung R., Ou C. and Yang, L. (2010). Part 5. Public health and air pollution in Asia (PAPA): a combined analysis of four studies of air pollution and mortality. *Res Rep Health Eff Inst*, **23**(154), 377–418.
- Wu R., Song X., Chen D., Zhong L., Huang X., Bai Y., Hu W., Ye S., Xu H., Feng B., Wang T., Zhu Y., Fang J., Liu S., Chen J., Wang X., Zhang Y. and Huang W. (2019). Health benefit of air quality improvement in Guangzhou, China: Results from a long time-series analysis (2006–2016). *Environment International*, **126**, 552–559.
- Wu W., Doreswamy V., Diaz-Sanchez D., Samet J.M., Kesic M., Dailey L., Zhang W., Jaspers I. and Peden D.B. (2011). GSTM1 modulation of IL-8 expression in human bronchial epithelial cells exposed to ozone. *Free Radical Biology & Medicine*, **51**(2), 522–529.
- Xu H., Xiao Z., Chen K., Tang M., Zheng N., Li P., Yang N., Yang W. and Deng X. (2019). Spatial and temporal distribution, chemical characteristics, and sources of ambient particulate matter in the Beijing-Tianjin-Hebei region. *Science of the Total Environment*, **658**, 280–293.
- Yang Q., Chen Y., Krewski D., Burnett R.T., Shi Y. and Mcgrail K.M. (2005). Effect of short-term exposure to low levels of gaseous pollutants on chronic obstructive pulmonary disease hospitalizations. *Environmental Research*, **99**(1), 99–105.
- Zanobetti Antonella, Schwartz and Joel. (2008). Mortality Displacement in the Association of Ozone with Mortality. *American Journal of Respiratory and Critical Care Medicine*, **177**(2), 184–189.
- Zhang J., Chen Q., Wang Q., Ding Z., Sun H. and Xu Y. (2019a). The acute health effects of ozone and PM_{2.5} on daily cardiovascular disease mortality: A multi-center time series study in China. *Ecotoxicology and Environmental Safety*, **174**, 218–223.
- Zhang J., Wei Y. and Fang Z. (2019b). Ozone Pollution: A Major Health Hazard Worldwide. *Frontiers in Immunology* **10**.
- Zhang Z., Zhang X., Gong D., Quan W., Zhao X., Ma Z. and Kim S.J. (2015). Evolution of surface O₃ and PM_{2.5} concentrations and their relationships with meteorological conditions over the last decade in Beijing. *Atmospheric Environment*, **108**, 67–75.
- Zong R., Xue Y., Liang W., Xu C., Zhu Y., Chen T., Lan Y., Wang L., Zhang J. and Yang L. (2018). Strong ozone production at a rural site in the North China Plain: Mixed effects of urban plumes and biogenic emissions. *Journal of Environmental Sciences*, **71**(9), 261–270.