

Exploring the effect of salinity as a primary cause of teratology in freshwater diatoms

Faïza Nouné^{1,2,*}, Nadjla Chaib^{3,4}, Sophia Metallaoui^{5,6}, Hadjer Kaddeche^{1,2}, Sabrina Dzizi^{2,3} and Saül Blanco^{7,8}

¹Department of natural and life sciences/Faculty of Sciences/University of 20 august 1955, Skikda, Algeria

²Laboratoire de recherche sur la physico-chimie des surfaces et interfaces (LRPCSI)–University of 20 august 1955, Skikda, Algeria

³Department of Process Engineering/Faculty of Technology/University of 20 august 1955, Skikda, Algeria

⁴Laboratory of Catalysis, Bioprocess and Environment–LCBE, University of 20 august 1955, Skikda, Algeria

⁵Department of Ecology/Faculty of Sciences/University of 20 august 1955, Skikda, Algeria

⁶Laboratoire de recherche des Interactions, Biodiversité, Ecosystèmes et Biotechnologie (LRIBEB), University of 20 august 1955, Skikda, Algeria

⁷Departamento de Biodiversidad y Gestión Ambiental/Facultad de Ciencias Biológicas y Ambientales/Universidad de León, Campus de Vegazana s/n, 24071, León, España


⁸Laboratorio de diatología. La Serna 58, 24007, León, España

Received: 23/11/2023, Accepted: 12/01/2024, Available online: 28/01/2024

*to whom all correspondence should be addressed: e-mail: f.nouné@univ-skikda.dz ; nounefaiza@yahoo.fr

<https://doi.org/10.30955/gnj.005586>

Graphical abstract



Abstract

Increased water conductivity levels have been proposed as a key factor to explain the occurrence of teratological forms in freshwater diatom assemblages. The current study aimed to carry out an experiment on a laboratory scale to assess the response of periphytic diatoms to artificially increased salinity levels. The obtained results show that: a) the proportion of aberrant cells increased in high conductivity treatments, b) developed teratologies were preserved in diatom populations even after being translocated to normal conditions, and c) the degree of valve outline deformation in the dominant species was proportional to the induced water conductivity. All these data support previous field observations that linked high electrolyte content and the abundance of aberrant cells among microalgal communities in continental aquatic ecosystems.

Keywords: aberrant cells, water conductivity, salt stress, diatom assemblages, phototrophic biofilm

1. Introduction

Diatoms are a type of single-celled algae with an important ecological role in the functioning of freshwater ecosystems (Morin *et al.* 2016). They have been widely used as bioindicators of various environmental conditions, particularly salinity, pH, and nutrients, due to their high diversity, short generation time, sensitivity, and quick response to changes in water quality (Kelly 2003; Smol and Stoermer, 2010; Stevenson *et al.* 1999). They also play a crucial role in aquatic food webs (Ragueneau *et al.* 2006; Weitere *et al.* 2018). Diatoms are also known to be sensitive to toxic substances (McCormick and Cairns 1994; Stevenson 2014). In their natural habitat, diatom communities are exposed to multiple anthropogenic inputs that affect their life cycle, so that the resulting valve morphologies developed under unfavorable environmental conditions differ from those inhabiting undisturbed environments. Valves that deviate that differ from normality in a population for a given species, show abnormalities in terms of symmetry, striation pattern, raphe course, and structure, are called teratological forms (Dziengo-Czaja *et al.* 2008; Falasco *et al.* 2009a; Gonçalves *et al.* 2019; Riato *et al.* 2018; Smol and Stoermer 2010).

Deformation in valve outline, loss of areolae, changes in striation patterns, and disruption of the raphe formation, are the main teratological forms affecting diatom valves, often correlated with physiological and metabolic impairment in the diatom cell (Falasco *et al.* 2021; Falasco *et al.* 2009a).

Teratology does not appear to weaken the reproductive capacity or viability of the affected cells (Falasco *et al.* 2021). However, some forms of teratologies are suspected of being lethal (Arini *et al.* 2013).

Recovering normal morphology after sexual reproduction in deformed diatoms from long-term cultures indicates that teratologies do not arise from genetic drift (Granetti, 1968). In this regard, the work of Arini *et al.* (2013) showed that the cadmium-induced teratologies in *Planothidium frequentissimum* (Lange-Bertalot) Lange-Bertalot decreased with decontamination, evidencing that teratology in diatoms appears in altered environments but progressively disappears along with the return to normal conditions.

The relationship between teratological forms and unhealthy conditions has been investigated in laboratory cultures (Duong *et al.* 2010; Windler *et al.* 2014) as well as in the field (Cantonati *et al.* 2014; Falasco *et al.* 2009a; Muhr 2014). In the latter case, natural teratology is rare and typically recorded at relative abundances not exceeding 1% (Morin *et al.* 2012a), so an increase in abundance can be linked to an increase in pollutants or other stressors (Cattaneo *et al.* 1998; Dziengo-Czaja *et al.* 2008; Lavoie *et al.* 2017; Sládeček 1986; Stevenson *et al.* 1999).

Factors known to be teratogenic for diatoms include high temperatures, light intensity (Antoine and Benson-Evans, 1984; Hill *et al.* 1995), low current velocity (typical of summer drought conditions (Stevenson 1996), high Ultraviolet (UV) radiation (Cabrol *et al.* 2004), increased nutrient concentrations (Nicolosi Gelis *et al.* 2020; Rosemond *et al.* 2000), and dissolved chemical substances in water, including organic or inorganic substances (Boisson and Perrodin, 2006; Cattaneo *et al.* 2008; Guasch *et al.* 2009; Hoagland *et al.* 1996; Moisset *et al.* 2015), herbicides (Debenest *et al.* 2008), and heavy metals (Cerisier *et al.* 2018; Cunningham *et al.* 2005; Falasco *et al.* 2009a; Gold *et al.* 2003; Morin *et al.* 2007; Pandey *et al.* 2018; Pandey and Bergey 2018). Interactions between these factors often occur in aquatic environments and are hard to disentangle (Falasco *et al.* 2021).

While changes in the composition of diatom communities have been shown to be a good mirror of freshwater ecosystem health thanks to many decades of comprehensive monitoring studies (Medley and Clements, 1998; Sabater, 2000), to date few experimental studies document the effect of abiotic factors on freshwater diatoms.

Olenici *et al.* (2017) have already revealed conductivity as a major cause for the occurrence of abnormal forms of epilithic diatoms in rivers. In this context, the main objective of the present study was to assess the effect of high salinity levels on freshwater diatoms under laboratory conditions. We focus particularly on the development of teratological forms, assessing the degree of deformation of valve outline, as well as on the dynamics of species richness in these assemblages.

2. Materials and methods


2.1. Experimental setup

The experiment was carried out in July 2020 by using plastic trays (length $\times$ width $\times$ height = 46 $\times$ 32 $\times$ 8 cm), filled with 4 L of water from an experimental freshwater pond

(Diatom Laboratory, Leon, Spain, 42.6047093° N, 5.5565376° W), to which 25 mL of a periphyton suspension collected from the same pond, and 80 mL of concentrated Alga-Gro® freshwater medium, were added. Two conductivity levels were maintained: the first one as a control treatment (low conductivity: LC: $\sim 460 \mu\text{S}\cdot\text{cm}^{-1}$: conductivity of water without any additions), and the second as a high conductivity (HC: $\sim 1400 \mu\text{S}\cdot\text{cm}^{-1}$) treatment by adding marine salt. Water conductivity was daily measured and adjusted. Trays were maintained at a constant water level under natural sunlight for 4 weeks.

Figure 3 shows a single trial. The same design was replicated 3 times (3 trials) in the same conditions. Each treatment was replicated three times. The experiment took place in a glass room receiving sunlight with natural intensity and photoperiod throughout the incubation period. The trays were kept at room temperature.

Microscope glass slides with a cavity were used as artificial substrata for the settlement and growth of periphytic diatoms. Each water tray contained three slides: two slides were removed from each treatment after two (slides LC₂ and HC₂), and four weeks (slides LC₄ and HC₄) respectively, whereas the third slide was translocated from LC to HC (slide Trans-H) and vice-versa (slide Trans-L) after the second week and removed at the end of the experiment (Figures 1–3). Removed slides were placed in 50 mL polypropylene screw-cap tubes (Falcon-BD, Franklin Lakes, NJ, USA) and stored in a freezer at -20 °C until analyses.



Figures 1, 2. Experiment in trays for attachment of diatoms




Figure 3. Schematic representation of the experiment consisting in two plastic trays, each contains three glass slides with a cavity, LC₂, Trans-H and LC₄ / HC₂, Trans-L and HC₄ respectively for Low conductivity (LC) and High conductivity (HC) trays. Points rectangle are the slides removed from each treatment, traits rectangle are the slides translocated from LC to HC (slide Trans-

H) and vice-versa (slide Trans-L). Removed slides were placed in 50 mL polypropylene screw-cap tubes (Falcon-BD, Franklin Lakes, NJ, USA) and stored in a freezer at -20 °C until analyses.

2.2. Identification

Slides were cleaned with hot hydrogen peroxide (30%) and a few drops of hydrochloric acid to remove organic material and dissolve calcium carbonates. The samples were then rinsed several times with distilled water. The cleaned samples were transferred to coverslips to dry for 24 hours. Once the samples were dry, permanent glass slides using high refractive resin Naphrax (RI=1.74) were mounted.

Diatom frustules were identified and counted under light microscopy (1000 × magnification, Olympus BX 60 microscope), with oil immersion, using standard references and segregating teratologic forms (Blanco Lanza *et al.* 2011; Hofmann *et al.* 2011). The relative abundances of diatom species were finally calculated.

2.3. Data processing

The relative abundance of teratological forms was set as the response variable. Due to the non-normality of this variable, statistical differences between treatments were analyzed by means of the Kruskal-Wallis test (Ostertagová *et al.* 2014).

Diatom assemblage composition was compared between treatments after 4 weeks using non-metric multidimensional scaling (NMDS), using Spearman rank correlation as distance measure, segregating normal and teratological forms within each genus. Singletons (genera occurring in a single sample) were previously removed. To test for significant differences between treatments, a one-way ANOSIM test (using Euclidean similarity coefficients) was carried between the Cartesian coordinates of the resulting groups.

All statistical analyses were performed with the free Past software, version 3.24 (Hammer *et al.* 2001).

2.4. Degree of teratologies

To evaluate the degree of valve deformation, geometric morphometry was used to analyze changes in valve outline caused by experimental treatments in *Pseudostaurosira brevistriata* (Grunow) D.M. Williams & Round, the dominant species in the samples.

A total of 141 individuals of *P. brevistriata* were photographed using Optikam digital camera and OptikaView7 software. Images were binarized and segmented using ImageJ software (Abràmoff *et al.* 2004). Valve outlines were then vectorized with Shape v.1.3 package (Iwata and Ukai, 2002), which uses Elliptical Fourier Analysis (EFA) to describe valve outline. EFA consists of fitting a given number of harmonics to the original valve outline, harmonics which are then analyzed by means of Principal component analysis (PCA).

Differences in valve outline between normal and teratologic *P. brevistriata* valves were tested by means of an ANOSIM test using Euclidean distances between the PCA scores.


3. Results


3.1. Identification

Diatom composition of all samples was similar with a strong dominance of *Encyonopsis subminuta* Krammer & Reichardt (20.63%), *Achnanthis minutissimum* (Kützing) Czarnecki (19.86%), *Pseudostaurosira brevistriata* (15.20%), *Staurosira venter* (Ehrenberg) Cleve & Moeller (11.01%), *Nitzschia palea* (Kützing) W. Smith (9.10%), and *Nitzschia dissipata* (Kützing) Grunow (6.06%).

Within the typology of deformations described in the literature (Falasco *et al.* 2021; Falasco *et al.* 2009b), the most widespread ones observed in our samples were the occurrence of irregular valve outlines, in species such as *A. minutissimum*, *A. pyrenaicum*, *Brachysira neoexilis*, *Cocconeis euglypta*, *C. lineata*, *Craticula subminuscula*, *E. subminuta*, *Eunotia sp.*, *Fragilaria gracilis*, several *Nitzschia* species (*N. palea*, *N. dissipata*, *N. solgensis*, *N. frustulum*, *N. inconspicua*, *N. amphibia*, *N. filiformis*, and *N. recta*), *Navicula tripunctata*, *N. veneta*, *Planothidium frequentissimum*, *P. brevistriata*, *S. venter*, *S. construens var. binodis*, *Punctastriata sp.*, and *Ulnaria acus*. Atypical raphes were found in *E. subminuta*, aberrant striae occurred in *F. gracilis* and several *Navicula* (*N. tripunctata*, *N. veneta*, and *N. trivialis*) and disrupted fibulae in *N. palea* and *N. filiformis*.

Combined teratologies have been also noted in *E. subminuta* (involving valve outline and in raphe), *F. gracilis*, *P. brevistriata*, *S. venter*, and *S. construens var. binodis* (irregular valve outline and aberrant striae), and *N. palea*, *N. filiformis*, and *N. frustulum* (disrupt fibulae and irregular valve outline) (Table 1). See Figures 4–68 for some deformed diatom species.





Figures 4–68. Examples of deformed frustules of some diatom species recorded in treatments exposed to a high level of water conductivity (LM microphotographs). Photos are not to scale

Figure 4–11. *Achnanthes minutissimum*: deformed valve outlines

Figure 12–16. *Staurosira venter*

Figure 12–15. Deformed valve outlines

Figure 16. Mixed teratologies: (deformed valve outline and aberrant striae)

Figure 17–19. *Eunotia* sp. deformed valve outlines

Figure 20–23. *Planorbulina freudentzenii* deformed valve outlines

Figure 24. *Punctastriatasp.*: deformed valve outline

Figure 25. *Cocconeis euglypta*: deformed valve outline

Figure 26–28. *Cocconeis lineata*: deformed valve outlines

Figure 29–32. *Navicula* sp. pl. deformed valve outlines

Figure 33–34. *Craticula subminuscula* deformed valve outlines

Figure 35–41. *Staurosira construens* var. *binodis* (deformed valve outlines)

Figure 42–50. *Encyonopsis subminuta*

Figure 42–48. Deformed valve outlines

Figure 49. Atypical raphe

Figure 50. Mixed teratologies (deformed valve outlines and atypical raphes)

Figure 51. *Brachysira neoxilis* (deformed valve outline)

Figure 52–66. *Nitzschia* sp. pl

Figure 52–62. Deformed valve outlines

Figure 63, 66. Mixed deformities (irregular valve outlines and disrupted fibulae)

Figure 64, 65. Disrupted fibulae

Figure 67. *Fragilaria gracilis*: deformed valve outline


Figure 68. *Ulnaria acus*: deformed valve outline

3.2. Teratology

The number of teratological forms was always significantly lower in control treatments, with no statistical differences between LC₂ and LC₄.

Figures 69–71 represent the percentage of teratological valves among different treatments. The exposure of the phototrophic biofilm to salt stress induced a significant increase ($H=9.46$, $p=0.023$) in the percentage of teratological frustules between treatments LC and HC. Control trays with low conductivity levels ($460 \pm 0.73 \mu\text{S}\cdot\text{cm}^{-1}$) harbored low numbers of teratological individuals during the first two weeks ($4.86 \pm 0.46\%$), raising to $6.41 \pm 0.35\%$ during the fourth week whereas the percentage of teratological forms increased to $11.68 \pm 1.87\%$ and $12.52 \pm 1.85\%$ in HC ($1400 \pm 40.32 \mu\text{S}\cdot\text{cm}^{-1}$) treatments, after 2 and 4 weeks, almost doubling the percentage in comparison to the LC treatment.

There was a significant difference in the number of abnormal forms comparing LC₂, LC₄, and Trans-H treatments ($H=7.2$, $p=0.027$, Figure 69), while no significant differences were found between Trans-L and Trans-H ($p=0.83$, Figure 70). Statistical analysis revealed also no significant differences ($p=0.96$) in the number of teratological forms comparing HC₂, HC₄, Trans-L, and Trans-H (Figure 71).



Figures 69–71. Boxplots showing the frequencies of teratological diatom valves across the different treatments. LC₂: Low conductivity after 2 weeks, LC₄: Low conductivity after 4 weeks, HC₂: High Conductivity after 2 weeks, HC₄: High Conductivity after 4 weeks, Trans-H: translocated from LC to HC after 2 weeks. Trans-L: translocated from HC to LC after 2 weeks. Error bars: standard error

No differences in terms of genera diversity (Shannon's H) were found between LC and Trans-H or between LC and HC levels, and no significant differences were observed either in species richness between low and high conductivity treatments after four weeks of colonization.

The comparison of diatom assemblages using NMDS (Figure 72) allowed clear segregation of normal and teratological individuals throughout the experiment in LC and HC treatments, revealing an evident effect of high conductivity levels on the overall biofilm response. Significant differences were observed ($p=0.0001$) between the position of normal and teratological individuals in the resulting plot at the end of the experiment.

Table 1. Diatom species found with the main deformity types

Species	Code genus	Code species	Type of teratology						
			One teratology				Mixed deformity in		
			Irregular valve outline	Atypical raphe	Aberrant striae	Disrupted fibulae	valve outline and raphe	valve outline and striae	valve outline and fibulae
Achnantheidium minutissimum (Kützing) Czarnecki	Acht	ADMI	+						
Achnantheidium pyrenaicum (Hustedt) H. Kobayasi		ADPY	+						
Brachysira neoxilis Lange-Bertalot	Brach	BNEO	+						
Cocconeis euglypta Ehrenberg	Cocco	CEUG	+						
Cocconeis lineata Ehrenberg	Crac	CLNT	+						
Craticula subminuscula (Manguin) Moser Lange-Bertalot & Metzeltin		ESBM	+						
Encyonopsis subminuta Krammer & Reichardt	Encyo	ESUM	+	+			+		
Eunotia sp.	Euno	EUNS	+						
Fragillaria gracilis Østrup	Fragi	FGRA	+		+			+	
Navicula tripunctata (O.F. Müller) Bory	Navi	NTPT	+		+				
Navicula trivialis Lange-Bertalot		NTRV			+				
Navicula veneta Kützing		NVEN	+		+				
Nitzschia amphibia Grunow	Nitz	NAMP	+						
Nitzschia palea (Kützing) W. Smith		NPAL	+			+			+
Nitzschia dissipata (Kützing) Grunow		NDIS	+						
Nitzschia frustulum (Kützing) Grunow		NIFR	+						+

Nitzschia filiformis (W.M. Smith) Van Heurck		NFIL	+		+		+
Nitzschia solgensis Cleve-Euler		NSOL	+				
Nitzschia inconspicua Grunow		NINC	+				
Nitzschia recta Hantzsch in Rabenhorst		NREC	+				
Planothidium frequentissimum (Lange-Bertalot) Lange- Bertalot	Plan	PLFR	+				
Pseudostaurosira brevistriata (Grun.in Van Heurck) Williams & Round		PSBR	+				+
Staurosira venter (Ehr.) Cleve & Moeller	Staur	SSVE	+				+
Staurosira construens var. binodis (Ehr.) Hamilton		SCBI	+				+
Punctastriata sp.	Punc	PUCS	+				
Ulnaria acus (Kützing) Aboal	Ulac	UACU	+				

The NMDS highlighted the different ecological behavior of the teratological forms with respect to the normal ones regardless of treatment, exposure time or substrate translocation status. The resulting graph reveals that actually teratological diatoms have different occurrences and abundances throughout the whole experimental material.





Figure 72. Non-metric multidimensional analysis of diatoms genera, normal (black) and aberrant forms (red). Achn : Achnanthes, Adla: Adlafia, Crac: Craticula, Ency: Encyonema, Euno: Eunotia, Gomph: Gomphonema, Reim: Reimeria, Rhop: Rhopalodia, Sella: Sellaphora. Other code's genera are included in Table 1.

3.3. Degree of teratology

Pseudostaurosira brevistriata was the most dominant species among the teratological forms. It was chosen as a model to quantify the degree of deformation in valve outline:

Geometric morphometry revealed few differences in shape between normal and teratological forms found in LC treatments, but differences in valve outline were evident at the end of the experiment in treatments exposed to high conductivity (Figures 73–80) (ANOSIM test on PCA scores: $p=0.001$, Figures 81, 82).




Figures 73–80. LM microphotographs of abnormal frustules of *Pseudostaurosira brevistriata* recorded at the end of the experiment in treatments exposed to a high level of water conductivity (HC). Scale bar = 10 µm

4. Discussion

The occurrence of abnormal individuals is among the most striking effects of environmental stress on diatom metabolism. While deformities can be initiated at different stages throughout the diatom life cycle, the processes leading to abnormal cell formation are yet unsolved (Falasco *et al.* 2021; Lavoie *et al.* 2017; Morin *et al.* 2012a). In particular, the deleterious effects of high salinity on photoautotrophs have been described by several

researchers (Allakhverdiev *et al.* 1999; Lyon *et al.* 2011; Mostaert 1995; Rijstenbil 2005; Schmid 1979; Schultz 1971; Sudhir and Murthy 2004; Vairavamurthy *et al.* 1985), and the literature gathers extensive research on the acclimation of photoautotrophic organisms to salt stress. For example, hypersaline conditions induce osmotic and ionic stress on cells, this tension disrupts photosynthetic activity, and increases respiration and growth arrest (Sudhir and Murthy, 2004). Salinity causes also indirect damage as a result of oxidative stress (Rijstenbil, 2005).



Figures 81, 82. PCA plot of the Elliptic Fourier descriptors obtained in the analyzed *P. brevistriata* populations: deformed (red), and normal (black) valves resulting from low (A) conductivity and high (B) conductivity after four weeks of exposure time

Noticeably, the altered valve outline was the only kind of teratology observed in control treatments. Although raphe modifications, altered striation patterns, disrupted fibulae, and mixed deformities (several deformities in the same frustule) were additionally observed under salt stress.

At the end of the experiment, Trans-L treatments harbored the highest proportion of teratological frustules (although not significantly different from HC or Trans-H treatments), even once translocated to low conductivity waters after two weeks of incubation under high conductivity conditions. This indicates that the deformed cells continue to replicate and transfer their teratologies for a number of generations. Hustedt. (1956) had already observed that particular ecological conditions at the time of zygotes' (auxospore) formation may induce changes in frustule morphology that are perpetuated during vegetative reproduction, leading to populations with a morphology quite different from that of the parent population. This new abnormal cell would then split by mitosis and pass on the abnormal shape to all subsequent daughter cells, as also suggested by Stoermer (1967).

It appears from the experiment that 2 weeks of stressful salinity conditions are sufficient to induce a maximum percentage of teratologies. The increase in teratology after 4 weeks compared to 2 weeks is not significant. This can be attributed to the fact that a plateau was reached after 2 weeks of exposure and no further increase in the percentage of teratologies will occur regardless of the increase in time.

Numerous studies (Clavero *et al.* 2000; Millan *et al.* 2020; Schmid, 1980; Tuchman *et al.* 1984) have found a positive correlation between increased salinity in freshwater environments and the number of individuals exhibiting teratological forms. Falasco *et al.* (2009b) noticed that in

long-term cultures, not only nutrients and waste but also salinity can stimulate the development of teratological forms, affecting primarily the production of structures involved in colony formation and cell anchorage to overcome floating difficulties, although teratological forms such as aberrant surface ornamentations (position and shape of areolae) and valve outline can also be formed (Falasco *et al.* 2009a). Salinity acts on diatoms primarily by osmotic pressure (Stoermer and Andresen, 2006) by impairing microtubule function (Falasco *et al.* 2009b), but sometimes the ionic water composition itself can be responsible for morphological variations in algae, as pointed out by Lewin and Roberston (1971), and Schmid (1979). The resulting deformation can be severe and identification becomes difficult even at the genus level (Stoermer and Andresen, 2006).

During her investigations, Schmid (1980) found a close relationship between changes in water salinity and sexual reproduction in *Anomoeoneis sphaerophora* (Ehrenberg) Pfitzer and *Surirella peisonis* Pantocsek. The raphe fissure in these diatoms was fragmented into short parts or tubules as salinity increased. Schultz (1971) also found that a modification in the valve pattern of *Cyclotella cryptica* Reimann, Lewin & Guillard and *C. meneghiniana* Kützing was related to salinity levels.

Likewise, Håkansson and Chepurnov. (1999) found that *Cyclotella meneghiniana* cells cultured at high salinity levels exhibited greater morphological variability, including differences in the pattern and the number of marginal striae, the number and placement of valve face fulcra, and the pattern of mantle fulcra and satellite pores.

Al-handal *et al.* (2014) in their study of Lake Sawa (a salt lake in Iraq), observed that water salinity was responsible for teratological traits in *Cocconeis sawensis* Al-Handal & Riaux-Gobin, with the occurrence of abnormalities in valve contour, deflection of the raphe, and distortion of areolae structure.

In our experiment, salt stress didn't induce deformities in some genera, particularly five out of 16 dominant genera (*Achnanthes*, *Encyonema*, *Epithemia*, *Gomphonema* and *Rhopalodia*) which didn't show any deformity in their frustules even under high conductivity conditions.

On the other hand, species such as *A. minutissimum*, *E. subminuta*, *N. palea*, *P. brevistriata*, *S. construens* var. *binodis*, and *S. venter* showed a high percentage of deformities under salt stress. Trobajo *et al.* (2004) and Vendrell-Puigmitja *et al.* (2021) noticed also these species-specific responses, but the reasons leading to such taxonomic differences remain unknown. Lavoie *et al.* (2017) think that certain genera are more likely to produce teratological forms than others when affected by "a certain kind" of disturbance.

The irregular valve outline showed the most prominent form of teratology in our samples, followed by aberrant striae, disrupted fibulae, and atypical raphe. Unlike several works (Jahn, 1986; Trobajo Pujadas, 2007; Trobajo *et al.*

2011, 2004) we did not notice a change in the length and width of cells throughout the experiment.

The present work demonstrates, particularly, a close relationship between salt stress and valve outline deformity in *P. brevistriata*. Peres-Weerts (2000) had already reported that the percentage of aberrant *P. brevistriata* increased with decreasing water quality conditions, but our study shows also a response in the degree of deformation in valve outline (loss of symmetry, bent, incised, swollen, or notched profile), assessed by means of EFA analysis. The results evidence a close relationship between salt stress and the deformation at the individual level. The degree of deformity is more marked in *P. brevistriata* growing in treatments exposed to salt stress compared to those collected from control trays. Our results are in agreement with Olenici *et al.* (2017) who detected a deformation gradient in *A. minutissimum* from acid mine drainage sites. They found a positive relationship between the deformation degree in valve outline and dissolved Zinc (Zn) concentrations. Mu *et al.* (2018) also focused their studies on the degree of teratology in *Halophora veneta* (Kützing) Levkov when exposed to cadmium (Cd) and lead (Pb) for a period of 96 hours. They found a slight deformation on *H. veneta* cells following cadmium exposure at a concentration of 1.42 mg L⁻¹, while obvious deformation occurred with cells largely expanded after 96 hours of lead exposure at a concentration of 15.35 mg L⁻¹.

Cells affected by mixed deformities are poorly viable and are unable to reproduce (Arini *et al.* 2013). According to many researchers, mixed teratologies are lethal, hence why they are rarely observed in natural biofilms (Falasco *et al.* 2009b), as well as in laboratory experiments (Arini *et al.* 2013). This may explain the low percentage of mixed teratologies compared to a single form of teratology (deformity) found in our experiment

It's interesting to note that diatom communities grown in our experiment under various conductivity levels did not differ significantly from one another; this suggests that high conductivity brought on by salt stress did not cause variations in the composition of diatom communities. Our findings are in contrast to those of earlier research that found metal pollution was responsible for changes in the species composition of diatoms (Cantonati *et al.* 2014; Cunningham *et al.* 2003; Duong *et al.* 2008; Morin *et al.* 2012b; Sgro *et al.* 2007). Venâncio *et al.* (2019) noticed that even small increases in salinity may be sufficient to induce structural changes in freshwater communities or to induce changes in trophic relationships. Accordingly, Vendrell-puigmitja *et al.* (2021) revealed that freshwater salinization, induced by effluent from potash mining caused a shift in the diatom community of the exposed biofilm.

Our results agree, in contrast, with those of (Millan *et al.* 2020) who found that mineralization and radioactivity did not induce any significant change in diatom communities.

5. Conclusion

Although our experiment used a simplistic experimental design, it demonstrated that high conductivity induced by marine salt addition has the potential to significantly alter

the structure of diatom frustules causing the appearance of teratological forms, this teratology continuing throughout the generations by vegetative reproduction. Irregular valve outlines, aberrant striae, disrupted fibulae, atypical raphes, and some mixed deformities are the most significant deformations observed during our research. The species *P. brevistriata* was a good example used to measure the teratology degree, correlated to conductivity levels. Our study also showed that salt stress did not induce any difference or change in the diatom community composition when comparing control and high conductivity treatments. These results can be complemented by experimentally establishing a conductivity gradient to establish the salinity thresholds that cause teratology in diatoms.

Future laboratory experiments on morphological changes associated with salinity fluctuations based on axenic cultures (i.e. examining the behavior of each species separately) will be interesting and helpful to refine the data presented in this work. Further studies at the molecular and proteome levels are also needed to understand the mechanism underlying the development of teratologies.

Acknowledgments

The present work is funded by the Algerian Ministry of Higher Education and Scientific Research and the University of 20 August 1955 (Algeria), and it fits into the university training research project (PRFU) carrying the code A16N01UN210120180002. We express our thanks to the team of laboratory of diatomology, university of Leon- Spain for their assistance during the experiment. Special thanks to Óscar Fernández and Adrián Llamazares for their precious technical help.

References

- Abràmoff M.D., Magalhães P.J., Ram S.J. (2004). Image processing with imageJ. *Biophotonics International*, **11**, 36–41. <https://doi.org/10.1201/9781420005615.ax4>
- Al-handal A.Y., Riaux-gobin C., Abdulla D.S., Ali M.H. (2014). *Cocconeis sawensis* sp. nov. (Bacillariophyceae) from a saline lake (Sawa Lake), South Iraq: comparison with allied taxa. *Phytotaxa*, **181**, 216–228. <https://doi.org/10.11646/phytotaxa.181.4.2>
- Allakhverdiev S.I., Nishiyama Y., Suzuki I., Tasaka Y. and Murata N. (1999). Genetic engineering of the unsaturation of fatty acids in membrane lipids alters the tolerance of *Synechocystis* to salt stress. *Proceedings of the National Academy of Sciences. U.S.A.*, **96**, 5862–5867. <https://doi.org/10.1073/pnas.96.10.5862>
- Antoine S.E. and Benson-Evans K. (1984). Teratological variations in the river Wye diatom flora, Wales, UK, in: *Proceedings of the 8th International Diatom Symposium, Paris*, 59–66.
- Arini A., Durant F., Coste M., Delmas F., Feurtet-Mazel A. (2013). Cadmium decontamination and reversal potential of teratological forms of the diatom *Planorhynchium frequentissimum* (Bacillariophyceae) after experimental contamination. *Journal of Phycology*, **49**, 361–370. <https://doi.org/10.1111/jpy.12044>
- Blanco Lanza S., Álvarez Blanco I., Cejudo Figueiras C. and Bécares Mantecón E. (2011). Guía de las diatomeas de la Cuenca del Duero. Ministerio de Medio Ambiente, Medio Rural y Marino / Confederación Hidrográfica del Duero.
- Boisson J.C. and Perrodin Y. (2006). Effects of road runoff on biomass and metabolic activity of periphyton in experimental streams. *Journal. Hazard. Mater*, **132**, 148–154. <https://doi.org/10.1016/j.jhazmat.2005.07.083>
- Cabrol N.A., Grin E.A., Hock A., Kiss A., Borics G., Kiss K., Acs E., Kovacs G., Chong G., Demergasso C., others. (2004). Investigating the Impact of UV Radiation on High-Altitude Shallow Lake Habitats, Life Diversity, and Life Survival Strategies: Clues for Mars' Past Habitability Potential?. *Lunar & Planetary Science*. 1049.
- Cantonati M., Angeli N., Virtanen L., Wojtal A.Z., Gabrieli J., Falasco E., Lavoie I., Morin S., Marchetto A., Fortin C. and Smirnova S. (2014). *Achnanthes minutissimum* (Bacillariophyta) valve deformities as indicators of metal enrichment in diverse widely-distributed freshwater habitats. *Science. Total Environment*, **475**, 201–215. <https://doi.org/10.1016/j.scitotenv.2013.10.018>
- Cattaneo A., Asioli A., Comoli P. and Manca M. (1998). Organisms' response in a chronically polluted lake supports hypothesized link between stress and size. *Limnology and Oceanography*, **43**, 1938–1943.
- Cattaneo A., Couillard Y. and Wunsam S. (2008). Sedimentary diatoms along a temporal and spatial gradient of metal contamination. *Journal. Paleolimnol*, **40**, 115–127. <https://doi.org/10.1007/s10933-007-9159-1>
- Cerisier A., Vedrenne J., Lavoie I. and Morin S. (2018). Assessing the severity of diatom deformities using geometric morphometry. *Botany Letters*, 1–9. <https://doi.org/10.1080/23818107.2018.1474800>
- Clavero E., Grimalt J.O., Garcia-pichel F., Clavero E. and Al E.T. (2000). Salinity tolerance of diatoms from thalassic hypersaline environments. *Journal. Phycology*, **36**, 1021–1034.
- Cunningham L., Raymond B., Snape I., Riddle M.J. (2005). Benthic diatom communities as indicators of anthropogenic metal contamination at Casey Station, Antarctica. *Journal of Paleolimnology*, **33**, 499–513. <https://doi.org/10.1007/s10933-005-0814-0>
- Cunningham L., Stark J.S., Snape I., Mcminn A. and Riddle M.J. (2003). Effects of metal and petroleum hydrocarbon contamination on benthic diatom communities near Casey station, Antarctica: an experimental approach. *Journal of Phycology*, **39**, 490–503.
- Debenest T., Silvestre J., Coste M., Delmas F. and Pinelli E. (2008). Herbicide effects on freshwater benthic diatoms: Induction of nucleus alterations and silica cell wall abnormalities. *Aquatic Toxicology*, **88**, 88–94. <https://doi.org/10.1016/j.aquatox.2008.03.011>
- Duong T.T., Morin S., Coste M., Herlory O., Feurtet-Mazel A. and Boudou A. (2010). Experimental toxicity and bioaccumulation of cadmium in freshwater periphytic diatoms in relation with biofilm maturity. *Science. Total Environment*, **408**, 552–562. <https://doi.org/10.1016/j.scitotenv.2009.10.015>
- Duong T.T., Morin S., Herlory O., Feurtet-Mazel A., Coste M. and Boudou A. (2008). Seasonal effects of cadmium accumulation in periphytic diatom communities of freshwater biofilms. *Aquatic Toxicology*, **90**, 19–28. <https://doi.org/10.1016/j.aquatox.2008.07.012>
- Dziengo-Czaja M., Koss J. and Matuszak A. (2008). Teratological forms of diatoms (Bacillariophyceae) as indicators of water

- pollution in the western part of Puck Bay (southern Baltic Sea). *Oceanol Hydrobiol Stud*, **37**, 119–132. <https://doi.org/10.2478/v10009-007-0042-1>
- Falasco E., Bona F., Badino G., Hoffmann L., Ector L. (2009). Diatom teratological forms and environmental alterations: A review. *Hydrobiologia*, **623**, 1–35. <https://doi.org/10.1007/s10750-008-9687-3>
- Falasco E., Bona F., Ginepro M., Hlúbíková D., Hoffmann L., Ector L. (2009b). Morphological abnormalities of diatom silica walls in relation to heavy metal contamination and artificial growth conditions. *Water SA*, **35**, 595–606. <https://doi.org/10.4314/wsa.v35i5.49185>
- Falasco E., Ector L., Wetzel C.E. and Badino G. (2021). Looking back , looking forward : a review of the new literature on diatom teratological forms (2010 – 2020). *Hydrobiologia*, **848**, 1675–1753. <https://doi.org/10.1007/s10750-021-04540-x>
- Gold C., Feurtet-Mazel A., Coste M. and Boudou A. (2003). Impacts of Cd and Zn on the development of periphytic diatom communities in artificial streams located along a river pollution gradient. *Archives of Environmental Contamination and Toxicology*, **44**, 189–197. <https://doi.org/10.1007/s00244-002-2024-3>
- Gonçalves S., Almeida S.F.P., Figueira E. and Kahlert M. (2019). Valve teratologies and Chl c in the freshwater diatom *Tabellaria flocculosa* as biomarkers for metal contamination. *Ecological Indicators*, **101**, 476–485. <https://doi.org/10.1016/j.ecolind.2019.01.032>
- Granetti B. (1968). Alcune forme teratologiche comparse in colture di *navicula minima* grun. e *navicula seminulum* grun. *Giornale Botanico Italiano*, **102**, 469–484. <https://doi.org/10.1080/11263506809426483>
- Guasch H., Leira M., Montuelle B., Geislinger A., Roulier J.L., Tornés E. and Serra A. (2009). Use of multivariate analyses to investigate the contribution of metal pollution to diatom species composition: Search for the most appropriate cases and explanatory variables. *Hydrobiologia*, **627**, 143–158. <https://doi.org/10.1007/s10750-009-9721-0>
- Håkansson H. and Chepurnov V. (1999). A Study of Variation in Valve Morphology of the Diatom *Cyclotella Meneghiniana* in Monoclonal Cultures: Effect of Auxospore Formation and Different Salinity Conditions. *Diatom Re*, **14**, 251–272. <https://doi.org/10.1080/0269249X.1999.9705469>
- Hammer Ø., Harper D.A.T., Ryan P.D. and others. (2001). PAST: Paleontological statistics software package for education and data analysis. *Palaeontology. Electrones*, **4**, 9.
- Hill W.R., Ryon M.G. and Schilling E.M. (1995). Light limitation in a stream ecosystem: responses by primary producers and consumers. *Ecology*, **76**, 1297–1309.
- Hoagland K.D., Carder J.P. and Spawn R.L. (1996). Effects of organic toxic substances, in: *Algal Ecology*. Elsevier, pp. 469–496.
- Hofmann G., Werum M. and Lange-Bertalot H. (2011). Diatomeen im Süßwasser-Benthos von Mitteleuropa: Bestimmungsflora Kieselalgen für die ökologische Praxis; über 700 der häufigsten Arten und ihrer Ökologie. Gantner.
- Hustedt F. (1956). Kieselalgen (Diatomeen). Einführung in die Kleinlebewelt. Kosmos. Gesellschaft der Natur freunde. Fr. Verlagshandlung Stuttgart 1–70.
- Iwata H. and Ukai Y. (2002). SHAPE: a computer program package for quantitative evaluation of biological shapes based on elliptic Fourier descriptors. *Journal of Heredity*, **93**, 384–385. <https://doi.org/10.1093/jhered/93.5.384>
- Jahn R. (1986). A study of Gomphonema augur Ehrenberg. The structure of the frustule and its variability in clones and populations, in: *8th Diatom Symposium 1984*. 191–204.
- Kelly M.G. (2003). Short term dynamics of diatoms in an upland stream and implications for monitoring eutrophication, *Environmental Pollution*, **125**, 117–122. [https://doi.org/10.1016/S0269-7491\(03\)00075-7](https://doi.org/10.1016/S0269-7491(03)00075-7)
- Lavoie I., Hamilton P.B., Morin S., Kim Tiam S., Kahlert M., Gonçalves S., Falasco E., Fortin C., Gontero B., Heudre D., Kojadinovic-Sirinelli M., Manoylov K., Pandey L.K. and Taylor J.C. (2017). Diatom teratologies as biomarkers of contamination: Are all deformities ecologically meaningful? *Ecology. Indica*, **82**, 539–550. <https://doi.org/10.1016/j.ecolind.2017.06.048>
- Lewin Ralph A and Roberston A. (1971). Influence of salinity on the form of Asterocytis in pure culture. *Journal of Phycology*, **7**, 236–238.
- Lyon B.R., Lee P.A., Bennett J.M., DiTullio G.R., Janech M.G. (2011). Proteomic analysis of a sea-ice diatom: salinity acclimation provides new insight into the dimethylsulfoniopropionate production pathway. *Plant Physiology*, **157**, 1926–1941.
- Mccormick P.V. and Cairns J. (1994). Algae as indicators of environmental change. *Journal of Applied Phycology*, **6**, 509–526.
- Medley C.N. and Clements W.H. (1998). Responses of diatom communities to heavy metals in streams: The influence of longitudinal variation. *Ecological Applications*, **8**, 631–644. [https://doi.org/10.1890/1051-0761\(1998\)008\[0631:RODCTH\]2.0.CO;2](https://doi.org/10.1890/1051-0761(1998)008[0631:RODCTH]2.0.CO;2)
- Millan F., Izere C., Breton V., Voldoire O., Biron D.G., Wetzel C.E., Miallier D., Allain E., Ector L., Beauger A. (2020). The effect of natural radioactivity on diatom communities in mineral springs. *Bot. Letter*, **167**, 95–113.
- Moisset S., Tiam S.K., Feurtet-Mazel A., Morin S., Delmas F., Mazzella N. and Gonzalez P. (2015). Genetic and physiological responses of three freshwater diatoms to realistic diuron exposures. *Environmental Science and Pollution Research*, **22**, 4046–4055. <https://doi.org/10.1007/s11356-014-3523-2>
- Morin S., Cordonier A., Lavoie I., Arini A., Blanco S., Becares E., Coste M., Faggiano L., Laviale M., Pe F., Fortin C., Guasch H. and Sabater S. (2012a). Consistency in diatom response to metal-contaminated environments, in: *Emerging and Priority Pollutants in Rivers*. Springer, pp. 117–146. https://doi.org/10.1007/978-3-642-25722-3_5
- Morin S., Gómez N., Tornés E., Licursi M., Rosebery J. (2016). Benthic diatom monitoring and assessment of freshwater environments: standard methods and future challenges. *Aquat. Biofilms Ecol. Water Qual. Wastewater Treat* 111–124.
- Morin S., Lambert A.S., Artigas J., Coquery M. and Pesce S. (2012b). Diatom immigration drives biofilm recovery after chronic copper exposure. *Freshwater Biology*, **57**, 1658–1666. <https://doi.org/10.1111/j.1365-2427.2012.02827.x>
- Morin S., Vivas-Nogues M., Thi T.D., Boudou A., Coste M. and Delmas F. (2007). Dynamics of benthic diatom colonization in a cadmium/zinc-polluted river (Riou-Mort, France). *Fundam. Applied Limnology*, **168**, 179–187. <https://doi.org/10.1127/1863-9135/2007/0168-0179>

- Mostaert U. (1995). Inorganic ions and mannitol in the red alga *Caloglossa leprieurii* (Ceramiales, Rhodophyta): Response to salinity change. *Phycologia*, **34**, 501–507. <https://doi.org/10.2216/i0031-8884-34-6-501.1>
- Mu W., Chen Y., Liu Y., Pan X. and Fan Y. (2018). Toxicological effects of cadmium and lead on two freshwater diatoms. *Environmental Toxicology and Pharmacology*, **59**, 152–162. <https://doi.org/10.1016/j.etap.2018.03.013>
- Muhr E. (2014). Teratological Diatom Deformities in the Periphyton of Colorado Alpine Streams as Indicators of Acid Mine Drainage Contamination. University of Colorado, Boulder.
- Nicolosi Gelis M.M., Cocherio J., Donadelli J. and Gómez N. 2020. Exploring the use of nuclear alterations, motility and ecological guilds in epipelagic diatoms as biomonitoring tools for water quality improvement in urban impacted lowland streams. *Ecological Indicators*, **110**, 105951. <https://doi.org/10.1016/j.ecolind.2019.105951>
- Olenici A., Blanco S., Borrego-Ramos M., Momeu L., Baciú C. (2017). Exploring the effects of acid mine drainage on diatom teratology using geometric morphometry. *Ecotoxicology* **26**, 1018–1030. <https://doi.org/10.1007/s10646-017-1830-3>
- Ostertagová E., Ostertag O. and Kováč J. (2014). Methodology and application of the Kruskal-Wallis test. *Applied Mechanics and Materials*, **611**, 115–120. <https://doi.org/10.4028/www.scientific.net/AMM.611.115>
- Pandey L.K. and Bergey E.A. (2018). Metal toxicity and recovery response of riverine periphytic algae. *Science Total Environment*, **642**, 1020–1031. <https://doi.org/10.1016/j.scitotenv.2018.06.069>
- Pandey L.K., Lavoie I., Morin S., Park J., Lyu J., Choi S., Lee H. and Han T. (2018). River water quality assessment based on a multi-descriptor approach including chemistry, diatom assemblage structure, and non-taxonomical diatom metrics. *Ecological Indicators*, **84**, 140–151. <https://doi.org/10.1016/j.ecolind.2017.07.043>
- Peres-Weerts F. (2000). Mise en évidence des effets toxiques des métaux lourds sur les diatomées par l'étude des formes tératogènes. Agence l'Eau Artois Picardie, Douai.
- Pujadas R.T. (2007). Ecological analysis of periphytic diatoms in Mediterranean coastal wetlands (Empordà wetlands, NE Spain). ARG Gantner Verlag.
- Ragueneau O., Conley D.J., Leynaert A., Longphuiet S.N. and Slomp C.P. (2006). Role of diatoms in silicon cycling and coastal marine food webs, In *The Silicon Cycle, Human Perturbations and Impacts on Aquatic Systems*. eds. V. Ittekkot D. Unger C. Humborg and N.T. An (Washington, DC: Island Press), 163–195.
- Riato L., Leira M., Della Bella V. and Oberholster P.J. (2018). Development of a diatom-based multimetric index for acid mine drainage impacted depressional wetlands. *Science Total Environment*, **612**, 214–222. <https://doi.org/10.1016/j.scitotenv.2017.08.181>
- Rijstenbil J.W. (2005). UV- and salinity-induced oxidative effects in the marine diatom *Cylindrotheca closterium* during simulated emersion. *Marine Biology*, **147**, 1063–1073. <https://doi.org/10.1007/s00227-005-0015-4>
- Rosemond A.D., Mulholland P.J. and Brawley S.H. (2000). Seasonally shifting limitation of stream periphyton: response of algal populations and assemblage biomass and productivity to variation in light, nutrients, and herbivores. *Canadian Journal of Fisheries and Aquatic Sciences*, **57**, 66–75. <https://doi.org/10.1139/cjfas-57-1-66>
- Sabater S. (2000). Diatom communities as indicators of environmental stress in the Guadamar River, S-W. Spain, following a major mine tailings spill. *Journal of Applied Phycology*, **12**, 113–124. <https://doi.org/10.1023/A:1008197411815>
- Schmid A.M. (1980). Valve morphogenesis in diatoms: a pattern-related filamentous system in pennates and the effect of APM, colchicine and osmotic pressure. *Nova Hedwigia*, **33**, 811–847.
- Schmid A.M.M. (1979). Influence of environmental factors on the development of the valve in diatoms. *Protoplasma*, **99**, 99–115. <https://doi.org/10.1007/BF01274072>
- Schultz M.E. (1971). Salinity-related polymorphism in the brackish-water diatom *Cyclotella cryptica*. *Canadian Journal of Botany*, **49**, 1285–1289. <https://doi.org/10.1139/b71-182>
- Sgro G.V., Poole J.B. and Johansen J.R. (2007). Western North American Naturalist Diatom species composition and ecology of the Animas River Watershed, Colorado, USA. *West. North American. National*, **67**, 510–519.
- Sládeček V. (1986). Diatoms as Indicators of Organic Pollution. *Acta Hydrochim. Hydrobiol*, **14**, 555–566. <https://doi.org/10.1002/aheh.19860140519>
- Smol J.P., Stoermer E.F. (2010). The diatoms: applications for the environmental and earth sciences. Cambridge University Press.
- Stevenson J. (2014). Ecological assessments with algae: a review and synthesis 1. *Journal. Phycology*, **50**, 437–461. <https://doi.org/10.1111/jpy.12189>
- Stevenson R.J. (1996). An introduction to algal ecology in freshwater benthic habitats. *Algal Ecology in Freshwater Benthic Ecosystems*. 3–30.
- Stevenson R.J., Pan Y. and Dam H.V. (1999). Assessing environmental conditions in rivers and streams with diatoms, in: *The Diatoms: Applications for the Environmental and Earth Sciences, Second Edition*. 57–85. <https://doi.org/10.1017/CBO9780511763175.005>
- Stoermer E.F. (1967). Polymorphism in *Mastogloia*. *Journal. Phycology*, **3**, 73–77.
- Stoermer E.F. and Andresen N.A. (2006). Atypical tabularia in coastal Lake Erie, USA. Foss. *Recent Phycology. Study*, 351–361.
- Sudhir. and Murthy S.D.S. (2004). Effects of salt stress on basic processes of photosynthesis. *Photosynthetica*, **42**, 481–486.
- Trobajo R., Cox E.J. and Quintana X.D. (2004). The effects of some environmental variables on the morphology of *Nitzschia frustulum* (Bacillariophyta), in relation its use as a bioindicator. *Nov. Hedwigia*, **79**, 433–445. <https://doi.org/10.1127/0029-5035/2004/0079-0433>
- Trobajo R., Rovira L., Mann D.G. and Cox E.J. (2011). Effects of salinity on growth and on valve morphology of five estuarine diatoms. *Phycological Research*, **59**, 83–90. <https://doi.org/10.1111/j.1440-1835.2010.00603.x>
- Tuchman M.L., Theriot E. and Stoermer E.F. (1984). Effects of low level salinity concentrations on the growth of *Cyclotella meneghiniana* Kütz. (Bacillariophyta). *Archangel. fur Protistenkd*, **128**, 319–326. [https://doi.org/10.1016/S0003-9365\(84\)80003-2](https://doi.org/10.1016/S0003-9365(84)80003-2)
- Vairavamurthy A., Anssdreare M.O. and Iverson R.L. (1985). Biosynthesis of dimethylsulfide and dimethylpropiothetin by

- Hymenomonas carterae in relation to sulfur source and salinity variations. *Limnol. Oceanogr*, **30**, 59–70. <https://doi.org/10.4319/lo.1985.30.1.0059>
- Venâncio C., Castro B.B., Ribeiro R., Antunes S.C., Abrantes N., Soares A.M.V.M. and Lopes I. (2019). Sensitivity of freshwater species under single and multigenerational exposure to seawater intrusion. *Philosophical Transactions of the Royal Society. B Biological Sciences*. **374**. <https://doi.org/10.1098/rstb.2018.0252>
- Vendrell-puigmitja L., Llenas L., Proia L., Ponsa S., Espinosa C., Morin S., Abril M. (2021). Effects of an hypersaline effluent from an abandoned potash mine on freshwater biofilm and diatom communities. *Aquatic Toxicology*, **230**, 105707. <https://doi.org/10.1016/j.aquatox.2020.105707>
- Weitere M., Erken M., Majdi N., Arndt H., Norf H., Reinshagen M., Traunspurger W., Walterscheid A. and Wey J.K. (2018). The food web perspective on aquatic biofilms. *Ecological Monographs*, **88**(4), 543–559. <https://doi.org/10.1002/ecm.1315>
- Windler M., Bova D., Kryvenda A., Straile D., Gruber A. and Kroth P.G. (2014). Influence of bacteria on cell size development and morphology of cultivated diatoms. *Phycological Research*, **62**, 269–281. <https://doi.org/10.1111/pre.12059>