

Improvement of the photo-Fenton performance at near neutral pH by the addition of tannic acid from tea leaves waste as a complexing agent in the Pb(II) photo-oxidation

Endang Tri Wahyuni^{1*}, Hardiana Kuncoro¹, Early Zahwa Alharrisa¹ and Adhitasari Suratman¹


¹Department of Chemistry, Faculty of Mathematic and Natural Sciences, Gadjah Mada University, Yogyakarta, Indonesia

Received: 21/08/2023, Accepted: 08/04/2024, Available online: 12/06/2024

*to whom all correspondence should be addressed: e-mail: endang_triw@ugm.ac.id

<https://doi.org/10.55555/gnj.005314>

Graphical abstract






Abstract

Introducing tannic acid from tea leaves waste as a complexing agent of Fe(III) into photo-Fenton process to prevent $\text{Fe}(\text{OH})_3$ precipitation at pH 7, and further to increase photo-Fenton activity in the near neutral pH for Pb(II) photo-oxidation is systematically addressed. The photo-Fenton process for Pb(II) photo-oxidation involving Fe^{2+} and H_2O_2 solutions as well as UV light was conducted by batch technique, both in the absence and present of tannic acid. Furthermore, the optimization of tannic acid concentration, solution pH, and reaction time were also conducted. The research results assign that the addition of tannic acid in the photo-Fenton process for photo-oxidation of Pb(II) is able to considerably enhance the effectiveness at pH 6-8 (near neutral pH). The activity of the tannic acid from tea leaves waste is comparable to the that of the commercial tannic acid. The best condition of Pb(II) photo-oxidation having $10 \text{ mg}\cdot\text{L}^{-1}$ in 50 mL of the solution can be reached by using Fe^{2+} as high $10 \text{ mmol}\cdot\text{L}^{-1}$, H_2O_2 $100 \text{ mmol}\cdot\text{L}^{-1}$, with the addition of tannic acid as much as $30 \text{ mg}\cdot\text{L}^{-1}$ at pH 7 and 60 min of the time that gave about 85% of the photo-oxidation effectiveness. It is also indicated that the Pb(II) photo-oxidation has resulted in the harmless PbO_2 solid. Hence, it is obviously assigned that the priceless tea waste could significantly contribute in the improvement of the photo-Fenton performance to prevent environmental pollution.

Keywords: Photo-Fenton, near neutral pH, tannic acid, Pb(II) photooxidation.

1. Introduction

Photo-Fenton is one of the advanced oxidation processes (AOPs) utilizing hydroxyl radical (noted as $\cdot\text{OH}$) as a strong oxidizing agent. The $\cdot\text{OH}$ is resulted from the reactions between Fe^{2+} and H_2O_2 as well as UV light with H_2O_2 as presented as equation (1) and (4) below [(Buitrago *et al.*, 2020; Choquehuanca *et al.*, 2021; Ebrahiem *et al.*, 2017a; Miranzadeh *et al.*, 2016; Ortega-Gómez *et al.*, 2012; Saldaña-Flores *et al.*, 2021; Thomas *et al.*, 2021; Wahyuni *et al.*, 2019; Wahyuni *et al.*, 2021):



Due to its easiness, practical and high performance, the method has been intensively examined for photo-degradation of various organic pollutants to form smaller and safer compounds (Ebrahiem *et al.*, 2017a; Miranzadeh *et al.*, 2016)], for photo-oxidation of toxic metal ions such as Pb(II) [(Wahyuni *et al.*, 2019) and As(V) (Wahyuni *et al.*, 2021) to be harmless compounds, as well as for bacterial disinfection (Ortega-Gómez *et al.*, 2012).

Unfortunately, the method works effectively only at pH around 3 or less active in the near neutral pH (Choquehuanca *et al.*, 2021; Ebrahiem *et al.*, 2017b; Miranzadeh *et al.*, 2016; Ortega-Gómez *et al.*, 2012; Saldaña-Flores *et al.*, 2021; Thomas *et al.*, 2021; Wahyuni *et al.*, 2019) due to the precipitate formation of $\text{Fe}(\text{OH})_3$ as seen in the Eq. 5. The precipitate can considerably reduce the quantity of Fe^{3+} present and also inhibit the light penetration. These drawbacks constrain the method in the wastewater treatment applications, since most wastewater has pH around 6-8 or near neutral pH (Ebrahiem *et al.*, 2017b; Miralles-Cuevas *et al.*, 2014).

An intensive effort has been demonstrated to extend the method activity in the wider range of pH, that is by introducing a complexing agent (Ahile *et al.*, 2020; Buitrago *et al.*, 2020; De Luca *et al.*, 2014; Fiorentino *et al.*, 2018; Guo *et al.*, 2021; Helal *et al.*, 2013; Miralles-Cuevas *et al.*, 2014; O'Dowd & Pillai, 2020; Pan *et al.*, 2019; Subramanian & Madras, 2016; Villegas-Guzman *et al.*, 2017). The complexing agents are usually organic compounds having several carboxylate groups (Ahile *et al.*, 2020; O'Dowd & Pillai, 2020), including humic acid (Ortega-Gómez *et al.*, 2012), citric acid (Guo *et al.*, 2021; Miralles-Cuevas *et al.*, 2014; O'Dowd & Pillai, 2020; Villegas-Guzman *et al.*, 2017), saccharic acid (Subramanian & Madras, 2016), ethylene diamine tetra acetic acid /EDTA (De Luca *et al.*, 2014; Pan *et al.*, 2019), oxalic acid (De Luca *et al.*, 2014), glutamic acid (Helal *et al.*, 2013), ascorbic acid (Guo *et al.*, 2021; Villegas-Guzman *et al.*, 2017), tartaric acid [(De Luca *et al.*, 2014; Guo *et al.*, 2021; Villegas-Guzman *et al.*, 2017), succinic acid (Guo *et al.*, 2021), caffeic acid (Villegas-Guzman *et al.*, 2017), nitriloacetic acid/NTA (De Luca *et al.*, 2014), (S,S)-ethylenediamine-N,N'-di-succinic acid (EDDS) (Miralles-Cuevas *et al.*, 2014; O'Dowd & Pillai, 2020), and tannic acid (Bolobajev *et al.*, 2016). These complexing agents also named as chelating agents (symbolized as L), are able to react with Fe^{3+} to form Fe-L_3 soluble complex compounds in the wider range pH. Chelation is useful to extend the pH range because the chelating ligand competes favourably with hydroxide ion, for complexes formation that are typically soluble in water (Ahile *et al.*, 2020; O'Dowd & Pillai, 2020). Hence, they can inhibit the precipitate $\text{Fe}(\text{OH})_3$ formation, and further enhance the effectiveness of the photo-Fenton process in the near neutral pH.




Figure 1. Chemical structure of tannic acid

Another complexing agent is tannic acid, having chemical structure as seen in Figure 1. The acid is widely distributed in several plants species such as tea, chocolate, cinnamon, chestnut, and grape, and are also found in wood, bark, leaves and fruit (Baldwin & Booth, 2022). Among them, tea leave waste is one of the priceless sources that has not been explored yet. The use of the tannic acid from very cheap sources will make photo-Fenton a cheaper process, allowing it to be widely applied.

Under the circumstance, in the present research, the addition of tannic acid from the tea leaves waste into conventional photo-Fenton process to increase its activity at the near neutral pH for Pb(II) photo-oxidation is addressed. The subject selected for evaluation of the photo-Fenton modified with tannic acid is Pb(II) ion, since this heavy metal ion is widely dispersed in the water environment due to the

disposal of various industrial wastewater (Ebraheim *et al.*, 2022; Wahyuni *et al.*, 2022; Wahyuni *et al.*, 2021). In fact, the heavy metal ion is reported to be hazard for environment and human health (Ebraheim *et al.*, 2022; Pan *et al.*, 2019). Oxidation seems to be the most interesting method for decreasing Pb(II) concentration (Wahyuni *et al.*, 2019), where the harmless and handleable PbO_2 precipitate is resulted (Ebraheim *et al.*, 2022; Wahyuni *et al.*, 2021; Wahyuni *et al.*, 2022). Currently, lack information of the research regarding the Pb(II) photo-oxidation through photo-Fenton process in the presence of tannic acid from the teabag waste is traceable.

Generally, the efficiency of the photo-Fenton process strongly depends on the parameters controlling the process, such as concentrations of Fe^{2+} (Buitrago *et al.*, 2020; De Luca *et al.*, 2014; Helal *et al.*, 2013.; O'Dowd & Pillai, 2020), H_2O_2 (Buitrago *et al.*, 2020; De Luca *et al.*, 2014; Saldaña-Flores *et al.*, 2021), and the complexing agent (De Luca *et al.*, 2014; Miralles-Cuevas *et al.*, 2014), as well as the solution pH (Guo *et al.*, 2021; O'Dowd & Pillai, 2020; Villegas-Guzman *et al.*, 2017; Wahyuni *et al.*, 2019; Wahyuni *et al.*, 2021) and reaction time (Buitrago *et al.*, 2020; Choquehuanca *et al.*, 2021; De Luca *et al.*, 2014; Guo *et al.*, 2021; Helal *et al.*, 2013.; Miralles-Cuevas *et al.*, 2014; O'Dowd & Pillai, 2020; Saldaña-Flores *et al.*, 2021; Villegas-Guzman *et al.*, 2017; Wahyuni *et al.*, 2019; Wahyuni, *et al.*, 2021). Hence, in the present research, the optimal condition of the photo-Fenton process is systematically determined. This study is hoped to valuably contribute to the development of photo-Fenton process and toxic metal remediation technology.

2. Experimental Section

2.1. Materials

FeCl_2 , H_2O_2 , commercial tannic acid, and $\text{Pb}(\text{NO}_3)_2$ purchased from Merck Company were used as received. Waste of teabags was chosen as a source of tannic acid. A set of photo-process apparatus was employed for photo-Fenton process. The analysis instruments including UV-Visible spectrophotometer, atomic absorption spectrophotometer (AAS) and transmission electron microscope equipped with electron diffraction spectrometer (SEM-EDX) machines were operated.

2.2. Research Procedures

2.2.1. Tannic acid extraction from the teabag waste

The teabag waste here is a teabag after being soaked in the hot water for 2 min, which is usually people do to make hot tea. The tea in the teabag waste was taken out from the paper bag and dried in an oven at 80 °C for 1 h. Then the dry tea waste was grounded and sieved into fine powder about 100 μm in size.

The fine powder of the tea waste about 10 g was mixed with 100 mL of ethanol 70%, accompanied by stirring for 18 h to allow the tannic acid dissolved from the tea waste powder into the water media. Then the solution containing tannic acid was separated from the powder by filtration to get a clear solution. It was continued with drying the clear solution at 80 °C for 1 h, so that the viscous liquid was obtained.

Tannic acid concentration was determined by using UV-Visible spectrophotometer machine. For that purpose, 50 mg of the viscous tannic acid was dissolved in 10 mL ethanol 70% to form clear yellow solution. Then the absorbance of the yellow solution was observed in the range 200-400 nm of the wavelength, which was observed maximally at 295 nm of the wavelength. The concentration was determined by interpolating its absorbance into the respective standard curve. From the calculation, it was found that the content of the tannic acid in the teabag waste was $\approx 42 \text{ mg}\cdot\text{g}^{-1}$.

2.2.2. Photo-oxidation of Pb(II) by photo-Fenton

Photo-Fenton processes was conducted through batch experiment in the photo-process apparatus that is equipped with 3 UV lamp with 40 watt of each, as illustrated in Figure 2. In this typical process, 50 ml of a solution containing Fe^{2+} 10 mmol. L^{-1} , H_2O_2 100 mmol. L^{-1} and Pb (II) 10 mg. L^{-1} was placed in the photo-process apparatus at room temperature, then was irradiated by UV light for a certain period of time. After the desired time, the solution was measured by AAS machine to find the concentration of the Pb(II) left in the solution. The effectiveness of the photo-oxidation (E) is presented as the oxidized Pb(II) in % that is calculated by following equation below:

$$E(\%) = \frac{C_o - C_t}{C_o} \times 100\%$$

C_o represents the initial Pb^{2+} concentration ($\text{mg}\cdot\text{L}^{-1}$) while C_t assigns the concentration of the un-oxidized or the left Pb(II) ($\text{mg}\cdot\text{L}^{-1}$).

The same procedure was repeated for the processes with various conditions as :

H_2O_2 concentrations were varied as: 25, 50, 75, 100, 125, 150, and 200 mmol. L^{-1} , with Fe^{2+} 10 mmol. L^{-1} , Pb(II) 10 mg. L^{-1} in 50 mL, pH 7, tannic acid 30 mg. L^{-1} and time 60 min.




Figure 2. Apparatus for photo-Fenton process

Fe^{2+} concentrations were adjusted as: 2, 4, 6, 8, 10, 12, 14, 16, 18, and 20 mmol. L^{-1} , with H_2O_2 100 mmol. L^{-1} , Pb(II) 10 mg. L^{-1} in 50 mL, pH 7, tannic acid 30 mg. L^{-1} and time 60 min.

The variation of tannic acid concentrations was set as : 10, 20, 30, 40, and 50 mg. L^{-1} with Fe^{2+} 10 mmol. L^{-1} , H_2O_2 100 mmol. L^{-1} , Pb (II) 10 mg. L^{-1} in 50 mL, pH 7, and time 60 min.

The reaction time was altered as : 15, 30, 45, 60, 75, 90, and 150 min, with Fe^{2+} 10 mmol. L^{-1} , H_2O_2 100 mmol. L^{-1} , Pb (II) 10 mg. L^{-1} in 50 mL, pH 7, and tannic acid 30 mg. L^{-1} .

The solution pH was varied as : 1, 3, 5, 7, 9, and 11, with Fe^{2+} 10 mmol. L^{-1} , H_2O_2 100 mmol. L^{-1} , Pb(II) 10 mg. L^{-1} in 50 mL, tannic acid 30 mg. L^{-1} and time 60 min.

3. Results and Discussion

3.1. Removal of Pb(II) through photo-Fenton process

The Pb(II) ion removal from the aqueous media under photo-Fenton process can be stimulated by precipitation of insoluble $\text{Pb}(\text{OH})_2$, and/or oxidation by $\bullet\text{OH}$ provided by Fenton agent and light exposure. The precipitation of Pb^{2+} with OH^- anionic into $\text{Pb}(\text{OH})_2$, written as Eq. 6, can only occur at pH higher than 7.5 assigned by its solubility product constant (K_{sp}) = $1.43 \cdot 10^{-20}$ (Wahyuni *et al.*, 2022).



In fact, the photo-Fenton was conducted at pH 7, suggesting that no precipitation of $\text{Pb}(\text{OH})_2$ formed. The second possibility is oxidation by $\bullet\text{OH}$ having high oxidation potential, that is 2.8 Volt (Wahyuni *et al.*, 2021; Wahyuni *et al.*, 2022). It is known that the standard reduction potential of $\text{Pb}(\text{IV})/\text{Pb}(\text{II})$ is -0.67 V (Wahyuni *et al.*, 2022), implying that the reduction is unlikely to proceed. In contrast, with such reduction potential, photo-oxidation of Pb(II) is more favourable. The photo-oxidation of Pb(II) by $\bullet\text{OH}$ producing PbO_2 is represented by Eq. 7. The more possible oxidation of Pb(II) is in agreement with the study reporting that Pb(II) could be oxidized by chlorine and was accelerated by Mn(VII) (Pan *et al.*, 2019). Accordingly, in the paragraph below, the decrease of the Pb(II) concentration is presented as photo-oxidation degree (in %).

3.2. Effect of the H_2O_2 concentration

In the photo-Fenton process, H_2O_2 is the main reagent to provide $\bullet\text{OH}$ both through catalysis decomposition by Fe^{2+} , and simultaneously under UV light exposure, as represented by Eq. 1 and Eq.4 respectively (Ahile *et al.*, 2020; Buitrago *et al.*, 2020; Choquehuanca *et al.*, 2021; De Luca *et al.*, 2014; Ebrahiem *et al.*, 2017b; Fiorentino *et al.*, 2018; Guo *et al.*, 2021; Helal *et al.*, 2013.; Miralles-Cuevas *et al.*, 2014; Miranzadeh *et al.*, 2016; O'Dowd & Pillai, 2020; Ortega-Gómez *et al.*, 2012; Pan *et al.*, 2019; Saldaña-Flores *et al.*, 2021; Subramanian & Madras, 2016; Thomas *et al.*, 2021; Villegas-Guzman *et al.*, 2017; Wahyuni *et al.*, 2019; Wahyuni *et al.*, 2021). The quantity of $\bullet\text{OH}$ produced strongly depends on the H_2O_2 concentration (Ahile *et al.*, 2020; Buitrago *et al.*, 2020; O'Dowd & Pillai, 2020), hence it is essential to find the most efficient concentration. Figure 3 exhibits the influence of the H_2O_2 concentration on the Pb(II) photo-oxidation degree.





Figure 3. The effect of H₂O₂ concentration, a) in the absent and b) in the present of tannic acid 30 mg.L⁻¹. (Pb(II) : 10 mg.L⁻¹ Volume = 50 mL, pH 7, Fe²⁺ = 10 mmol. L⁻¹, time = 60 min).

From Figure 3, it is notified that the elevation of H₂O₂ concentration results in the enhancement of the photo-oxidation. The highest level is reached by applying H₂O₂ as much 100 mmol. L⁻¹. However, the higher concentration of H₂O₂ than 100 mmol. L⁻¹ is observed to apparently decline the photo-oxidation degree. It is clearly implied that the amount of •OH produced is proportional to the H₂O₂ concentration up to 100 mmol. L⁻¹. A decrease in the photo-oxidation with excessive H₂O₂ occurred because the excess H₂O₂ could capture the •OH present, as seen in Eq. 8 (Buitrago *et al.*, 2020) that forms radical of •OOH. The consecutive reaction of scavenging •OH by the •OOH occurred, resulting in water and oxygen gas (Eq. 9). These reactions caused the •OH depletion, that was consequently less conducive for the Pb(II) photo-oxidation. The finding trend is well matched with previous reports (Ahile *et al.*, 2020; O'Dowd & Pillai, 2020; Wahyuni *et al.*, 2019; Wahyuni *et al.*, 2021).



3.3. Effect of the Fe²⁺ concentration

The role of Fe²⁺ is also very important, that is as a catalyst to promote •OH formation from H₂O₂ decomposition. The amount of •OH produced are also dependent on the quantity of Fe²⁺ ions. Accordingly it is important to get the optimum Fe²⁺ concentration, that was conducted by observing the effect of Fe²⁺ concentration on the Pb(II) photo-oxidation degree.

That result is displayed in Figure 4, notifying that the increase of the Fe²⁺ concentration can promote the Pb(II) oxidation which attain the maximum level at 10 mmol. L⁻¹ of Fe²⁺ concentration. The extension of Fe²⁺ concentration could generate more •OH, that promoted more effective photo-oxidation. In contrast, the descent photo-oxidation is notified when Fe²⁺ concentration was further elevated into more than 10 mmol. L⁻¹. The excessive Fe²⁺ was able to act as •OH scavenger, as following Eq.10 (Buitrago *et al.*, 2020; Ebrahiem *et al.*, 2017a), that led to the •OH

dismissed. Simultaneously, large quantity of Fe³⁺ and •OH resulted from Eq. 1, were allowed to mutually react to form insoluble Fe(OH)₃ (Eq.10). This insoluble hydroxide could screen the light penetration (Ahile *et al.*, 2020; De Luca *et al.*, 2014; O'Dowd & Pillai, 2020; Saldaña-Flores *et al.*, 2021; Wahyuni *et al.*, 2019; Wahyuni *et al.*, 2021). These conditions were less conducive for the photo-oxidation of Pb(II) process. Some others also found same trend data (Ahile *et al.*, 2020; De Luca *et al.*, 2014; O'Dowd & Pillai, 2020; Saldaña-Flores *et al.*, 2021; Wahyuni *et al.*, 2019; Wahyuni *et al.*, 2021).




Figure 4. The effect of Fe²⁺ concentration, a) in the absent, and b) in the present of tannic acid (Pb(II) : 10 mg. L⁻¹, volume = 50 mL, pH 7, H₂O₂ = 100 mmol. L⁻¹, time = 60 min).

From the data above, it can be found that the optimal Fe²⁺ concentration is 10 mmol.L⁻¹ and for H₂O₂ is 100 mmol. L⁻¹, giving mole ratio of 1/10. It has been frequently reported that the amount of Fe²⁺ was always lower than that of H₂O₂, with the various mole ratio ranging from 1/25 to 1/4 (Buitrago *et al.*, 2020; Ebrahiem *et al.*, 2017a; Wahyuni *et al.*, 2019; Wahyuni *et al.*, 2021). The lesser Fe²⁺ concentration is beneficial in the technical application, since it can inhibit the large amount the production of iron sludge (Buitrago *et al.*, 2020; Ebrahiem *et al.*, 2017a).

3.4. Effect of the tannic acid concentration

Figures 3, 4, 7, and 8 illustrate the effect of the tannic acid addition into the photo-Fenton process. It shows that the addition of tannic acid at pH 7 can distinctly raise the photo-oxidation. The enhancement of the photo-oxidation is continued to proceed when the tannic acid concentration was raised up to 30 mg. L⁻¹, as depicted in Figure 5. In the solution with pH 7, the tannic acid with pK_a = 6, is dissociated into its anionic form and hydrogen cation. The anionic tannic acid can easily react with Fe³⁺ ion resulted from the reaction of Fe²⁺ with H₂O₂ (Eq.1.) to form soluble Fe(III)-tannic acid complex compound, as illustrated by Figure 6. This formation of the soluble Fe(III)-tannic acid can strongly prevent the formation of insoluble Fe(OH)₃ (Bolobajev *et al.*, 2016). Hence, the more tannic anion provided, the more effective prevention of the of Fe(OH)₃ formation can be attained.

The contrary photo-oxidation results are notified when the tannic acid concentration was continued to enlarge into larger than 30 mg. L⁻¹. With higher tannic concentration than 30 mg. L⁻¹, there might be free anionic tannic acid that had no chance to react with Fe³⁺. The free anionic tannic acid was allowed to be degraded by •OH, that dismissed the quantity of the •OH (Bolobajev *et al.*, 2016). Consequently, the photo-oxidation has become less effective (Ahile *et al.*, 2020; Bolobajev *et al.*, 2016; De Luca *et al.*, 2014; O'Dowd & Pillai, 2020).




Figure 5. Influence of concentration of the tannic acid: a) from teabag waste, and b) commercial product (Pb(II) = 10 mg.L⁻¹, volume = 50 mL, pH 7, Fe²⁺ = 10 mmol. L⁻¹, H₂O₂ = 100 mmol. L⁻¹, time = 60 min).




Figure 6. The structure of Fe(III)-tannic complex compound (Bolobajev *et al.*, 2016).

Moreover, it also appears in the Figure 5, that the effectiveness of Pb(II) photo-oxidation promoted by tannic acid from teabag waste is seen to be similar, both in trend and effectiveness, to that of by tannic acid commercial. It is evidence that the waste seems to be potential as a cheap and effective chelating agent, that can replace the costly commercial tannic acid.

3.5. Effect of the solution pH

In the photo-Fenton process, pH is a key factor determining the effectiveness of the process. It demonstrates in Figure 7, in the process without tannic acid, the photo-oxidation is less effective at very low pH, due to the formation of the stable Fe(H₂O₂)₂ complex compound (Ahile *et al.*, 2020; O'Dowd & Pillai, 2020). Such complex formation could neutralize the Fe²⁺ ions, to form inactive catalyst, that prevented the •OH generation (Ahile *et al.*, 2020; O'Dowd & Pillai, 2020). Consequently, the effectiveness of the photo-oxidation was diminished.

It is also notable, the Pb(II) photo-oxidation proceeds more effectively, and reaches the highest level when the pH moves up into 3. In such range pH, the complex of Fe(H₂O₂)₂ was dissociated giving more free Fe²⁺ and H₂O₂. The more number of Fe²⁺ and H₂O₂ could stimulate them to mutually react to form large number of •OH, that was conducive for photo-oxidation. The opposite results are shown at pH higher than 3, due to the higher precipitate formation of Fe(OH)₃, that filtered the penetration of light. Consequently, the lesser light was penetrated, leading to the lack of •OH available. This data obtained well agreed with the finding reported by some others (De Luca *et al.*, 2014; Guo *et al.*, 2021; Helal *et al.*, 2013.; Miralles-Cuevas *et al.*, 2014; Villegas-Guzman *et al.*, 2017).




Figure 7. Influence of the solution pH on the photo-oxidation, a) in the absent b) in the present of tannic acid (Pb(II) : 10 mg. L⁻¹, Volume = 50 mL, Fe²⁺ = 10 mmol. L⁻¹, H₂O₂ = 100 mmol. L⁻¹, time = 60 min, tannic acid = 30 mg. L⁻¹).

The Figure 7 also assigns clearly that the addition of tannic acid at lower pH, can inhibit the photo-oxidation. In such low pH, tannic acid was formed as non-ionic molecule that is difficult to form complex compounds with Fe²⁺ or Fe³⁺ ions, allowing them to exist as free acid molecules. These free molecules were possible to be degraded by •OH available, that subtracted the radicals present (Bolobajev *et al.*, 2016). The more effective photo-oxidation is observable as the increasing pH from 3 into 7. However, it shows the contrast photo-oxidation results, when the pH was further elevated into higher than 7. In the solution with pH higher than 5, the tannic acid was dissociated into its anions and hydrogen cation, as presented in equation (10) (Bolobajev *et al.*, 2016). The anions are easier to react coordinatively with Fe²⁺ and Fe³⁺ forming soluble complex compound, that could prevent Fe³⁺ from the precipitation with the OH anions present. This condition should maintain the large amount of Fe²⁺ and Fe³⁺ in the solution, that generate more •OH, and finally promoted the higher photo-oxidation. However, at higher pH, the OH anions were formed in excess, that competed with tannic anion in the reaction with Fe³⁺. Consequently, the number of Fe³⁺ ion was depleted and further declined the photo-oxidation. The similar finding has also been intensively reported (Ahile *et al.*, 2020; Bolobajev *et al.*, 2016; De Luca *et al.*, 2014; Guo *et al.*, 2021; O'Dowd & Pillai, 2020; Villegas-Guzman *et al.*, 2017).

3.6. Effect of the reaction time

Figure 8 represents the alteration time influencing the effectiveness of the photo-oxidation. It is seen in the figure that the prolong reaction time results in the higher photo-oxidation and the photo-oxidation kept to increase as the reaction time was further lengthen up to 60 min. The photo-oxidation is not depended on the time after 60 minutes of the reaction. The longer reaction time, the more OH radicals could be provided and the more effective contact between the OH radicals with Pb^{2+} ions also proceeded. After 60 mins, more PbO_2 solids may be formed in the solution that prevented the light entering into the solution (Wahyuni *et al.*, 2019). Consequently, the photo-oxidation insignificantly was altered or even slightly went down. This data is in a good agreement with other reports (Wahyuni *et al.*, 2019; Wahyuni *et al.*, 2021).




Figure 8. Influence of the reaction time on the photo-oxidation, a) in the absent and b) in the present of tannic acid (Pb(II): 10 $mg.L^{-1}$, Volume = 50 mL, Fe^{2+} = 10 $mg.L^{-1}$, H_2O_2 = 100 $mg.L^{-1}$, pH = 7, tannic acid = 30 $mg.L^{-1}$).

3.7. Detection of the iron and Pb(IV) present after the photo-Fenton process

It has been reported that the Pb(II) photo-oxidation has produced PbO_2 solid material (Wahyuni *et al.*, 2019; Wahyuni *et al.*, 2021).

In order to confirm the formation of the oxide, SEM image accompanied with the EDX spectra have been taken. Figure 8 depicts the result of SEM and EDX spectra of slurry material from photo-Fenton process. From the EDX spectra, it is notable the presence of Fe and Pb elements. The presence of Fe element was originated from the insoluble $Fe(OH)_3$ whereas the Pb element may be attributed to PbO_2 . From the EDX spectra, it is obviously implied that the oxidation has resulted in PbO_2 .

To find the content of Pb in more quantitatively, X-ray fluorescence (XRF) analysis was conducted and the results from the photo-Fenton process with various tannic acid concentration were exhibited in Table 1.

It is notable from the table that the content of Pb appears to increase, while the Fe amount decreases, This data is consistent with the degree of the Pb(II) photo-oxidation. The data provides clearly evidence that the photo-oxidation of Pb(II) into PbO_2 was successfully proceeded.





Figure 9. The SEM image and EDX spectra of the slurry from the photo-Fenton process.

Table 1. Elemental composition of the slurry from photo-Fenton in the Pb(II) photo-oxidation conducted by XRF.

Tannic acid addition ($mg.L^{-1}$)	Pb(II) photo-oxidation from the solution (%)	Pb content from slurry (% mol)	Fe content from slurry (% mol)
10	60.25	6.26	12.52
30	85.50	15.74	7.70
50	75.12	10.52	9.80

4. Conclusion

It is concluded that the addition of tannic acid from the teabag waste into the photo-Fenton process can considerably enhance the process performance at the near neutral pH in the Pb(II) photo-oxidation. Furthermore, the enhancement of the photo-oxidation result is found to be controlled by tannic acid concentration, and the tannic acid as much 30 $mg.L^{-1}$ showed the highest effect in the improvement. The ability of the tannic acid from the teabag waste is found to be comparable with the commercial tannic acid. In addition, the highest photo-oxidation of 100 mL of Pb(II) 10 $mg.L^{-1}$, about 86%, can be achieved by employing Fe^{2+} 10 $mmol.L^{-1}$, H_2O_2 100 $mmol.L^{-1}$, pH 7 and time 60 min in the presence of 30 $mg.L^{-1}$ tannic acid. Hence, it is beneficial from the utilizing the priceless waste to enhance photo-Fenton performance in the reducing the toxic contaminant.

References

Ahile U. J., Wuana R. A., Itodo A. U., Sha'Ato R. and Dantas R. F. (2020). A review on the use of chelating agents as an alternative to promote photo-Fenton at neutral pH: Current trends, knowledge gap and future studies. *In Science of the*

- Total Environment* (Vol. 710). Elsevier B.V. <https://doi.org/10.1016/j.scitotenv.2019.134872>
- Baldwin A. and Booth B. W. (2022). Biomedical applications of tannic acid. In *Journal of Biomaterials Applications* (Vol. 36, Issue 8, pp. 1503–1523). SAGE Publications Ltd. <https://doi.org/10.1177/08853282211058099>
- Bolobajev J., Trapido M. and Goi A. (2016). Interaction of tannic acid with ferric iron to assist 2,4,6-trichlorophenol catalytic decomposition and reuse of ferric sludge as a source of iron catalyst in Fenton-based treatment. *Applied Catalysis B: Environmental*, 187, 75–82. <https://doi.org/10.1016/j.apcatb.2016.01.015>
- Buitrago J. L., Sanabria J., Gutiérrez-Zapata H. M., Urbano-Ceron F. J., García-Barco A., Osorio-Vargas, P. and Rengifo-Herrera J. A. (2020). Photo-Fenton process at natural conditions of pH, iron, ions, and humic acids for degradation of diuron and amoxicillin. *Environmental Science and Pollution Research*, 27(2), 1608–1624. <https://doi.org/10.1007/s11356-019-06700-y>
- Choquehuanca A., Ruiz-Montoya J. G. and La Rosa-Toro Gómez A. (2021). Discoloration of methylene blue at neutral pH by heterogeneous photo-Fenton-like reactions using crystalline and amorphous iron oxides. *Open Chemistry*, 19(1), 1009–1020. <https://doi.org/10.1515/chem-2021-0077>
- De Luca A., Dantas R. F. and Esplugas S. (2014). Assessment of iron chelates efficiency for photo-Fenton at neutral pH. *Water Research*, 61, 232–242. <https://doi.org/10.1016/j.watres.2014.05.033>
- Ebraheim G., Karbassi A. R. and Mehrdadi N. (2022). Employing speciation of metals to assess photo-assisted electrochemical efficiency for improving rainwater quality in Tehran, Iran, *International Journal of Environmental Science and Technology*, 19(1), 261–280. <https://doi.org/10.1007/s13762-021-03127-2>
- Ebrahiem E. E., Al-Maghrabi M. N. and Mobarki A. R. (2017a). Removal of organic pollutants from industrial wastewater by applying photo-Fenton oxidation technology. *Arabian Journal of Chemistry*, 10, S1674–S1679. <https://doi.org/10.1016/j.arabjc.2013.06.012>
- Ebrahiem E. E., Al-Maghrabi M. N. and Mobarki A. R. (2017b). Removal of organic pollutants from industrial wastewater by applying photo-Fenton oxidation technology. *Arabian Journal of Chemistry*, 10, S1674–S1679. <https://doi.org/10.1016/j.arabjc.2013.06.012>
- Fiorentino A., Cucciniello R., Di Cesare A., Fontaneto D., Prete P., Rizzo L., Corno G. and Proto A. (2018). Disinfection of urban wastewater by a new photo-Fenton like process using Cu-iminodisuccinic acid complex as catalyst at neutral pH. *Water Research*, 146, 206–215. <https://doi.org/10.1016/j.watres.2018.08.024>
- Guo Q., Zhu W., Yang D., Wang X., Li Y., Gong C., Yan J., Zhai J., Gao X. and Luo Y. (2021). A green solar photo-Fenton process for the degradation of carbamazepine using natural pyrite and organic acid with in-situ generated H₂O₂. *Science of the Total Environment*, 784. <https://doi.org/10.1016/j.scitotenv.2021.147187>
- Helal M., Ali O. and Hussein Mady A. (2013). Catalytic Degradation of Phenol Using Different Chelating Agents at Near Neutral pH in Modified-Fenton process Anti-inflammatory heterocyclic Steroids View project wastewater treatment using modified Fenton reaction as applicable method View project. <https://www.researchgate.net/publication/282750928>
- Miralles-Cuevas S., Oller I., Pérez J. A. S. and Malato S. (2014). Removal of pharmaceuticals from MWTP effluent by nanofiltration and solar photo-Fenton using two different iron complexes at neutral pH. *Water Research*, 64, 23–31. <https://doi.org/10.1016/j.watres.2014.06.032>
- Miranzadeh M. B., Zarjam R., Dehghani R., Haghghi M., Badi H. Z., Marzaleh M. A. and Tehrani A. M. (2016). Comparison of linear alkyle benzene sulfonate (LAS) from aqueous solutions. *Polish Journal of Environmental Studies*, 25(4), 1639–1648. <https://doi.org/10.15244/pjoes/61824>
- O’Dowd K. and Pillai S. C. (2020). Photo-Fenton disinfection at near neutral pH: Process, parameter optimization and recent advances. In *Journal of Environmental Chemical Engineering* (Vol. 8, Issue 5). Elsevier Ltd. <https://doi.org/10.1016/j.jece.2020.104063>
- Ortega-Gómez E., Fernández-Ibáñez P., Ballesteros Martín M. M., Polo-López M. I., Esteban García B. and Sánchez Pérez J. A. (2012). Water disinfection using photo-Fenton: Effect of temperature on *Enterococcus faecalis* survival. *Water Research*, 46(18), 6154–6162. <https://doi.org/10.1016/j.watres.2012.09.007>
- Pan W., Pan C., Bae Y. and Giammar D. (2019). Role of Manganese in Accelerating the Oxidation of Pb(II) Carbonate Solids to Pb(IV) Oxide at Drinking Water Conditions. *Environmental Science and Technology*, 53(12), 6699–6707. <https://doi.org/10.1021/acs.est.8b07356>
- Pan Y., Guo H., Zhou M., Zhang Y., Tian Y. and Wang, W. (2019). EDTA enhanced removal of sulfamethazine by pre-magnetized Fe₀ without oxidant addition. *Chemical Engineering Journal*, 372, 905–916. <https://doi.org/10.1016/j.cej.2019.04.211>
- Saldaña-Flores K. E., Flores-Estrella R. A., Alcaraz-Gonzalez V., Carissimi E., de Souza B. G., Ruotolo L. A. M. and Urquieta-Gonzalez E. (2021). Regulation of hydrogen peroxide dosage in a heterogeneous photo-fenton process. *Processes*, 9(12). <https://doi.org/10.3390/pr9122167>
- Subramanian G. and Madras G. (2016). Introducing saccharic acid as an efficient iron chelate to enhance photo-Fenton degradation of organic contaminants. *Water Research*, 104, 168–177. <https://doi.org/10.1016/j.watres.2016.07.070>
- Thomas N., Dionysiou D. D. and Pillai S. C. (2021). Heterogeneous Fenton catalysts: A review of recent advances. In *Journal of Hazardous Materials* (Vol. 404). Elsevier B.V. <https://doi.org/10.1016/j.jhazmat.2020.124082>
- Villegas-Guzman P., Giannakis S., Torres-Palma R. A. and Pulgarin C. (2017). Remarkable enhancement of bacterial inactivation in wastewater through promotion of solar photo-Fenton at near-neutral pH by natural organic acids. *Applied Catalysis B: Environmental*, 205, 219–227. <https://doi.org/10.1016/j.apcatb.2016.12.021>
- Wahyuni E. T., Nurhikmatillah A., Kurniasari H. and Siswanta D. (2021). Detoxification of As(III) in aqueous media by using photo-Fenton method. *Global Nest Journal*, 23(4), 550–555. <https://doi.org/10.30955/gnj.003265>
- Wahyuni E. T., Pratama N. A., Lestari N. D. and Suherman S. (2022). Enhancement of TiO₂ activity under visible light by N,S codoping for Pb(II) removal from water. *Journal of*

Engineering and Applied Science, **69**(1).
<https://doi.org/10.1186/s44147-022-00069-5>

Wahyuni E. T., Rahmaniati T., Hafidzah A. R., Suherman S. and Suratman A. (2021). Photocatalysis over N-doped TiO₂ driven by visible light for Pb(II) removal from aqueous media. *Catalysts*, **11**(8). <https://doi.org/10.3390/catal11080945>

Wahyuni E. T., Siswanta D., Kunarti E. S., Supraba D. and Budiraharjo S. (2019). Removal of Pb(II) ions in the aqueous solution by photo-Fenton method. *Global Nest Journal*, **21**(2), 180–186. <https://doi.org/10.30955/gnj.002936>