

Ammonium nitrate enhances salinity stress tolerance by inducing physiological and phytohormonal regulation changes in *Vigna radiata* L.

Shilin Fu¹, Babar Iqbal¹*, Tahani Awad Alahmadi², Mohammad Javed Ansari³, Arshad Jalal⁴, Muhammad Junaid Nazir¹, Guanlin Li^{1,5}* and Daolin Du⁶*

¹School of Environment and Safety Engineering, Jiangsu University, Zhenjiang 212013, People's Republic of China

²Department of Pediatrics, College of Medicine and King Khalid University Hospital, King Saud University, Medical City, PO Box-2925, Riyadh -11461, Saudi Arabia

³Department of Botany, Hindu College Moradabad (Mahatma Jyotiba Phule Rohilkhand University Bareilly) 244001- India

⁴School of Engineering, Department of Plant Health, Rural Engineering and Soils, São Paulo State University - UNESP-FEIS, Ilha Solteira, 15385-000, São Paulo, Brazil.

⁵Jiangsu Collaborative Innovation Center of Technology and Material of Water Treatment, Suzhou University of Science and Technology, Suzhou 215009, People's Republic of China

⁶Jingjiang College, Institute of Environment and Ecology, School of Emergency Management, School of Environment and Safety Engineering, School of Agricultural Engineering, Jiangsu University, Zhenjiang 212013, People's Republic of China

Received: 02/04/2025, Accepted: 25/04/2025, Available online: 16/05/2025

*to whom all correspondence should be addressed: e-mail: babar@ujs.edu.cn, liguanlin@ujs.edu.cn, ddl@ujs.edu.cn https://doi.org/10.30955/gnj.07514

Graphical abstract



Highlights

- Soil salinity reduced growth and ultimately diminished the final productivity
- Salinity stress reduced aboveground fresh and dry biomass and chlorophyll contents
- Salinity stress increased H₂O₂, MDA, organic solutes, and antioxidant enzymes
- Nitrogen supplementation reduced H₂O₂ and MDA but improved other parameters
- Ammonium nitrate improved salinity tolerance and mitigating damage due to salinity

Abstract

Mungbean is a key leguminous crop species highly regarded for its nutritious properties. However, soil salinity significantly reduced its growth and yield. Our study investigated whether applying exogenous nitrogen (N) (ammonium nitrate; AN) could improve salinity tolerance in mungbean seedlings by supplementing them with N under different concentrations of NaCl stress. Results revealed that the increased NaCl stress levels significantly reduced shoot height, root length, fresh and dry weight, and chlorophyll pigment. In contrast, increased hydrogen peroxide (H₂O₂), malondialdehyde (MDA), organic solutes (soluble protein, proline, and soluble sugar), antioxidant enzymes, i.e., superoxide dismutase, ascorbate peroxidase, catalase, and peroxidase, and abscisic acid were significantly elevated. Jasmonic acid (JA) increased at 40 mM stress and decreased at 120 mM stress level. There was also a significant reduction in the concentration of cytokinin, indole acetic acid, and N-metabolizing enzymes, viz. nitrate reductase (NR) and glutamate synthase (GOGAT) activity. In contrast, N supplementation reduced H₂O₂ and MDA but improved chlorophyll levels, osmolytes, phytohormones, N assimilation potential, and enzymatic antioxidant activity. Exogenous N supplementation reduced the impairments by enhancing their growth and metabolism. Conclusively, AN application may provide an effective strategy for improving salinity tolerance and mitigating damage caused by salinity by modulating physiological responses.

Keywords: Abiotic stress; salinization; nitrogen fertilization; antioxidant mechanism; salt-sensitivity

Shilin Fu, Babar Iqbal, Tahani Awad Alahmadi, Mohammad Javed Ansari, Arshad Jalal, Muhammad Junaid Nazir, Guanlin Li and Daolin Du. (2025), Ammonium nitrate enhances salinity stress tolerance by inducing physiological and phytohormonal regulation changes in *Vigna radiata* L., *Global NEST Journal*, **27**(6), 07514.

1. Introduction

The mung bean is an economically significant crop species rich in minerals, protein, vitamins, fibers, essential amino acids, and carbohydrates (Sehrawat *et al.* 2019). Globally, mungbean accounts for approximately 5% of all pulse production (World Vegetable Centre 2018). Mung bean plants are impacted by salinity in several ways, viz seed germination, growth characteristics, relative water content (RWC), nutrient acquisition, excessive reactive oxygen species (ROS), photosynthesis, nodulation, and nodule respiration. It has also been reported that under 50 mM NaCl, its yield decreased to almost 70%, adversely affecting crop quality (Sehrawat *et al.* 2019). Hence, it is imperative to improve salt tolerance to maintain its productivity in salinity-affected soils worldwide.

Increasing soil salinity generally causes a higher accumulation of toxic salt ions, including namely sodium ion (Na⁺) and Chloride ion (Cl⁻), leading to ionic toxicity, osmotic imbalance, impaired nutrient acquisition, metabolic disruption, excessive ROS, and cellular oxidative stress damage (Ullah et al. 2022a; Wang et al. 2025). It is evident that plants adopt several mechanisms to protect themselves from the adverse effects associated with salinity, including the production of osmolytes, the exclusion or compartmentalization of Na⁺ and Cl⁻ into vacuoles and old tissues, the biosynthesis and regulation of antioxidant mechanisms (non-enzymatic and enzymatic) (Solangi et al. 2019; Zaid and Wani 2019). As a consequence of excessive salinity causes ionic toxicity and osmotic stress effects, resulting in the generation of ROS such as singlet oxygen $({}^{1}O_{2})$, superoxide radicals $(O_{2}^{\bullet-})$, and hydrogen peroxide (H₂O₂), hydroxyl radicals (•OH) (Ullah et al. 2022b). The higher ROS accumulation causes oxidative stress, leading to metabolic changes and degradation of cellular membranes (lipid peroxidation), RNA, DNA, and proteins (Mignolet-Spruyt et al. 2016; Azeem et al. 2020; Zhang et al. 2020).

When plants undergo oxidative stress, one of their immediate responses is to eliminate excess ROS via physiological and molecular mechanisms that stimulate antioxidant enzymes viz. peroxidase (POD), superoxide dismutase (SOD), catalase (CAT), and ascorbate peroxidase (APX) and antioxidant metabolites (Sarker and Ercisli 2022; Sarker et al. 2023), that remove ROS. Therefore, plants can reduce oxidative stress damage by upregulating their antioxidant system to neutralize ROS and continue average growth. Hence, the upregulation of the antioxidant mechanism has been reported to correlate positively with the salt stress condition in the soil (Wang et al. 2017; Sahab et al. 2021). In stressful conditions, plants accumulate saltrelieving solutes such as soluble sugars, glycine betaine, and proline. To maintain salinity homeostasis, osmolytes regulate water balance and cell division, prevent ion toxicity and chlorophyll degradation, stabilize cellular structures, and eliminate excess ROS (Ullah et al. 2020; Singh et al. 2022).

Phytohormones (PGRs) are natural signalling molecules that act as plant growth regulators. As phytohormone signalling pathways cross-talk extensively, phytohormones can have several effects on metabolic and signalling pathways, such as additive, synergistic, and antagonistic effects (Aerts et al. 2021), and they can also have significant effects on plant responses to biotic and abiotic stresses (Peleg and Blumwald 2011; Wei et al. 2021). A deeper understanding of phytohormone signaling pathways regulating salt stress responses may provide a better understanding of abiotic stress responses and the interplay between phytohormones, potentially revealing new strategies to enhance salt stress adaptation in crop species. Nitrogen (N) is an essential constituent of organic molecules, and in the process of N metabolism, nitrate reductase (NR) function to reduce absorbed nitrate (NO₃-) into nitrite (NO₂⁻) and then into ammonia (NH₄⁺) (Liu et al. 2014). Following its synthesis, NH₄⁺ is converted into amino acids either through the action of glutamine synthetase and glutamate synthase (GS/GOGAT) or through another enzyme known as glutamate dehydrogenase (GDH) (Khan et al. 2023). The increase in salinity has adverse effects on N metabolism due to reduced uptake of NO3⁻ and NH4⁺ or decreased activity of N-metabolizing enzymes (Ullah et al. 2019, 2022a; Abile et al. 2024).

Salinization affects ≈3600 Mha of arable land annually, causing USD 27.5 billion in losses (Zhang and Shi 2013; Wani et al. 2020). Humanity's significant challenge is the rapid rise in food demand and soil salinization. Besides, rapid industrialization, global warming, environmental fluctuations, irrigation with salty water, and unsustainable fertilizer consumption will further exacerbate the future salinization problem (Zhu 2001; Zuo et al. 2021; Ma et al. 2022). So far, researchers have suggested several management strategies for reducing the adverse impacts of salinity on plants, including cultivating salt-tolerant varieties and improving soil salinity. Nevertheless, the implementation of these measures will take a very long time. In previous studies, mineral nutrition and supplemental plant growth regulators are the most effective methods to improve plant tolerance (Singh et al. 2014; Guo et al. 2017; Ravi et al. 2022; Raveena et al. 2024). It is well-documented that N assimilation plays a vital role in determining a plant's ability to withstand salinity. However, a complex relationship exists between N metabolism and salinity due to several factors, such as salt stress intensity, plant species' susceptibility, and soil availability (Dai et al. 2015). Besides being a vital macronutrient, N is a key player contributing to better crop production after water (Kant 2018). Further, N participates in many cellular functions within plants owing to its presence in enzymes, proteins, amino acids, nucleic acids, osmolytes, and hormones (Singh et al. 2014, 2019; Liu et al. 2023). all of which function differently and are involved in salinity tolerance mechanisms (Arghavani et al. 2017).

Thus, this study speculates that additional N applications will increase mungbean salt tolerance. Exogenous N has been shown to significantly modulate the growth of Brassica (Siddiqui *et al.* 2010), wheat (Arghavani *et al.* 2017), tomato plants (Singh *et al.* 2019), and cotton (Sikder *et al.* 2020), subjected to salt stress. Based on the findings of the studies discussed above, N-applied positive modulation occurred via the upregulation of

photosynthesis and antioxidant systems and the reduction of oxidative stress damage. Despite numerous studies investigating the effects of salinity on mung beans, little is known about the interaction between salinity stress and exogenous N application in mung beans. Accordingly, this study investigated the role of N in improving mungbean salt tolerance. Our study hypothesized that (a) increasing salinity could adversely affect mungbean plants by disturbing their physio-biochemical functions and growth, and (b) exogenous N supplementation could reduce salinity-stress-induced damages in mungbeans by modulating their physio-biochemical characteristics. A comprehensive study of morphological and physiobiochemical changes in mungbean plants was conducted to test our hypothesis, including a detailed analysis of the changes in shoot and root biomass, photosynthetic pigments, salt ion concentration, antioxidant mechanisms, osmolytes accumulation, endogenous phytohormones, and N metabolism under salinity and N supplementation.

2. Materials and methods

2.1. Study area and experimental steps

A greenhouse experiment was conducted at the campus of Jiangsu University, Zhenjiang City, China. Two kilograms of soil (Silt-loamy soil; pH 6.9 and EC 0.288 dS/m) were placed in small plastic pots with a bottom opening (2 cm in diameter). Following a complete randomized block design (CRBD), pots containing uniform seedlings (n = 2 seedlings per pot) were selected and divided into eight groups for salt stress treatments and N application. Four treatments were conducted for salinity stress (0 mM, 40 mM, 80 mM, and 120 mM NaCl). The four remaining groups were subjected to the same salinity stress levels but supplied with exogenous N. A solution of ammonium nitrates i.e., during irrigation, 5 mM of ammonium/nitrate (50/50 ratio) was applied to the root medium. Ammonium sulfate ((NH₄)₂SO₄) and calcium nitrate (Ca(NO₃)₂) were used as a source of NH_4^+ and NO_3^- (50/50) and were mixed with water (-AN, +AN) and applied to the pots during irrigation. The vapor-transpired water was replenished every three days to maintain the target volume. Following is the formula for calculating soil relative water content (SRWC):

$$SRWC = \begin{pmatrix} [(Wsoil - Wpot - DWsoil)/\\ ((WFC - Wpot - DWsoil)] * 100 \end{pmatrix}$$
(3)

In this case, Wsoil corresponds to the weight of the soil, pot, and water at the time of saturation; Wpot corresponds to the weight of the empty plastic pot; DW soil corresponds to the weight of the dry soil; and WFC corresponds to the weight of the soil at capacity (soil + pot + water). Each treatment was replicated five times. The growth characteristics of the mungbean plants were determined six weeks after harvesting. The remaining pots were harvested and frozen in liquid N the same day before being stored at -80 °C for laboratory analysis.

2.2. Determination of growth characteristics

Root length, shoot height, fresh weight (FW), and dry weight (DW) were measured after the plants were harvested. Carefully uprooted the plants and divided them into different parts. The FW and length of the root and

shoot were calculated. After this, keep the samples at 60 °C constant temperature in the oven to dry for about three to four days. Lastly, dry weight was calculated.

2.3. Determination of chlorophyll pigments

A chlorophyll extract (0.1 to 0.3 g) was prepared using ethanol (95 %, v/v), and the absorbances were measured at 665 nm and 649 nm (Lichtenthaler and Buschmann 2001).

2.4. Determination of mineral elements

Leaf powder samples were digested with concentrated HNO₃ and a volume of 15 ml of deionized water. An inductively coupled plasma-optical emission spectrometry (ICP-OES) was used to estimate Na⁺ and K⁺ contents (Yousfi *et al.* 2010).

2.5. Determination of phytohormones

We determined the endogenous phytohormones, including abscisic acid (ABA), indole acetic acid (IAA), cytokinin (CTK), and jasmonic acid (JA) in fresh leaf samples using ELISA according to the instructions included with the Plant ABA ELISA Kit, Plant IAA ELISA Kit, Plant JA ELISA Kit, and Plant CTK ELISA Kit (MiBio, Shanghai).

2.6. Oxidative stress indicators

A standardized technique was used to measure the concentration of H_2O_2 (Patterson *et al.* 1984). Fresh samples (0.2 g) were homogenized in trichloroacetic acid (5 ml, 0.1%) and then centrifuged at $5000 \times g$ (10 min, 4 °C). An absorbance measurement was conducted at 410 nm (Patterson *et al.* 1984). MDA content was measured using the thiobarbituric acid (TBA) test (Zhou *et al.* 2007). Leaf samples (0.5 g) were homogenized in trichloroacetic acid (TCA; 1 ml and 5%) solution and then centrifuged at 5,000 $\times g$ (4°C, 10 min). The OD450, OD532, and OD600 nm were measured using a spectrophotometer

2.7. Determination of antioxidant enzyme activities

We ground and homogenized fresh leaf samples in phosphate buffer (0.1 M; pH 7.3) containing EDTA (0.6 mM). We centrifuged the mixture at $12000 \times g$ (10 min; 4°C). Next, the enzyme extract in the supernatant was used for the assays. The catalase (CAT) activity was assessed by monitoring the disappearance of H_2O_2 (Sabra *et al.* 2012), and the absorbance was read at 240 nm using a spectrophotometer. The absorbance was read at 460 nm for peroxidase (POD) estimation following standard methods (Wang et al. 2018). Moreover, using a spectrophotometer, we measured the superoxide dismutase (SOD) activity based on the decline rate of nitroblue tetrazolium at 560 nm (Giannospolitis and Ries 1977). The ascorbate peroxidase (APX) activity was measured following a standard method (Iqbal et al. 2024a). A rise in absorption (290 nm) was noted following ascorbate oxidation using a spectrophotometer.

2.8. Determination of proline and soluble sugar, soluble protein, and glycine betaine

After homogenizing 0.3 g of samples in sulfosalicylic acid (3%; 5 ml) and acetic acid (10%; 2 ml), they were centrifuged at 5,000 × g (10 min). The absorbance at 520 nm was measured using a spectrometer (Liu *et al.* 2014). We calculated the soluble sugar using the anthrone sulphuric acid (Irigoyen *et al.* 1992). Leaf samples were extracted in distilled water and centrifuged to determine

the concentration of glycine betaine. The absorbance at 530 nm was measured using a spectrophotometer (Grieve and Grattan 1983). Using glucose as a standard, we measured the absorbance spectrophotometrically at 620 nm. Following Bradford's (Bradford 1976), we determined

the soluble protein concentration using the Coomassie brilliant blue G-250 reagent.



Figure 1. Changes in growth feature (A) shoot height, (B) shoot fresh weight, (C) shoot dry weight, (D) root length, (E) root fresh weight, and (F) root dry weight, following salinity stress and N supplementation. The bars represent the mean data (n = 3) and SD. T1, T2, T3 and T4 represent 0, 40, 80, and 120 mM salinity stress, respectively. Following the LSD Tukey test, different letters indicate significant differences at *P* < 0.05.



Figure 2. Changes in growth feature (A) Chl a, (B) Chl b, and (C) Chl/Chlb, following salinity stress and N supplementation. The bars represent the mean data (n = 3) and SD. T1, T2, T3 and T4 represent 0, 40, 80, and 120 mM salinity stress, respectively. Following the LSD Tukey test, different letters indicate significant differences at *P* < 0.05.



Figure 3. Changes in the potassium (K⁺) concentration (mg g⁻¹) (A), sodium (Na⁺) concentration (mg g⁻¹) (B), and K⁺/Na⁺ ratio (C) under salinity stress and AN application. The bars represent the mean data (n = 3) and SD. T1, T2, T3 and T4 represent 0, 40, 80, and 120 mM salinity stress, respectively. Following the LSD Tukey test, different letters indicate significant differences at *P* < 0.05.



Figure 4. Changes in the concentration of (A) Abscisic acid concentration, (B) Jasmonic acid concentration, (C) Indole acetic acid concentration and (D) Cytokinin concentration following salinity stress and N supplementation. The bars represent the mean data (n = 3) and SD. T1, T2, T3 and T4 represent 0, 40, 80, and 120 mM salinity stress, respectively. Following the LSD Tukey test, different letters indicate significant differences at *P* < 0.05.

2.9. Determination of N metabolizing enzymes

The resultant supernatant assessed the NR activity using a diazo-coupling technique with Griess reagent (Sánchez-Rodríguez *et al.* 2011). In addition, GS activity was determined according to a previous study (Ullah *et al.* 2019). The NADH oxidation was continuously monitored by measurement of absorbance at 340 nm using a spectrophotometer.

2.10. Statistical analysis

The measurements were repeated three times. Statistical analysis and one-way analysis of variance (ANOVA) were performed using SPSS version 16.0 (Chicago, IL, United States). We used Duncan's multiple range tests at a significance level of p > 0.05 to compare the mean values. The figures were created using Microsoft Excel 2013. Pearson correlation and principle component analysis were carried out using OriginPro 2023.

3. Results

3.1. Hazardous effects of salinity on growth attributes of Vigna radiata L. and its alleviation by N-application

The results showed that salinity stress substantially reduced the growth parameters of the Vigna radiata L

crop. The plant shoot height, shoot FW (fresh weight), shoot DW (dry weight), root length, root FW, and root DW experienced a 31, 38, 32, 26, 67, and 42% decline under 40 mM, while 37, 55, 62, 35, 74, and 71% at 80 mM, and 54, 62, 71, 48, 77, and 71% inhibition under 120 mM salinity stress compared to control treatment, respectively (**Figure 1A-F**). However, the N-application improved the growth indices such as shoot height, shoot FW, shoot 2.4, 2.1, 1.4, 2.9, and 2.4% at 80 mM, while 2.4, 3.0, 1.8, 1.8, 5, and 4% at 120 mM of salinity stress, respectively (**Figure 1A-F**).

3.2. Interactive effect of salinity stress and N-application on Vigna radiata L chlorophyll pigments.

The physiological attributes are significantly reduced under salinity stress. Therefore, ChI a and ChI b declined up to 13.2 and 35% at 40 mM, 58 and 55% at 80 mM, and 41 and 63% at 120 mM of salinity stress, respectively, compared to the control treatment. Interestingly, N-application did not significantly affect ChI a concentration under either control or salinity stress treatments (**Figure 2A-B**). Although Napplication improved ChI b concentration under 40 mM stress, it had no significant impact under 80 mM and 120 mM stress levels. Additionally, the ratio of ChI a/ChI b increased across all salinity treatments compared to the control. However, N-application had no significant effect on the Chl a/Chl b ratio under salinity treatments compared to their untreated counterparts.

3.3. Increment in nutrient uptake by N-application under salt stress

The K⁺ and K⁺/Na⁺ reduced up to 27 and 52% at 40 mM; 41 and 71% at 80 mM, whereas 49-and 80% declined following 120 mM of salt stress, respectively (**Figure 3AC**). Contrary to others, Na⁺ uptake significantly increased by 1.5-, 2.0-, and 2.5-fold under all salinity levels compared to control

(Figure 3B). However, N supplementation decreased Na⁺ by 1.4-fold at 40 mM, 1.7-fold at 80 mM, and 2.7-fold at 120 mM salinity, respectively (3B). Furthermore, the results revealed that N application enhanced the concentration of K⁺ and K⁺/Na⁺ by 1.1-, 1.6-fold at 40 mM, 1.1-, and 2.6-fold at 80 mM, while 1.6-, and 4.4-fold at 120 mM of salinity, respectively (3AC).



Figure 5. Changes in the concentration of (A) hydrogen peroxide (H₂O₂), and (B) Malondialdehyde (MDA), and enzymatic acitiity of (C) S uperoxide dismutase (SOD), (D) Catalase (CAT), (E) Peroxidase (POD), and (F) Ascorbate peroxidase (APX) following salinity stress and N supplementation. The bars represent the mean data (n = 3) and SD. T1, T2, T3 and T4 represent 0, 40, 80, and 120 mM salinity stress, respectively. Following the LSD Tukey test, different letters indicate significant differences at P < 0.05.</p>

3.4. Changes in endogenous phytohormones

The concentration of abscisic acid (ABA) increased by 6.0, 21.7, and 47.9% at 40, 80, and 120 mM salinity stress, whereas jasmonic acid (JA) enhanced at 40 mM by 3.3%, compared to controlled conditions (**Figure 4 A-B**). In contrast, indole acetic acid (IAA) and cytokinin (CTK) decreased by 31.8 and 23.8 at 40 mM, 30.2 and 36.5% at

80 mM, and 42.2, and 33.8% at 120 mM salinity stress, respectively (Figure 4 C-D). Similarly, compared to their controlled conditions, JA decreased by 7.2 and 36.0% at 80 and 120 mM Salinity stress levels. Exogenous N supplementation improved the phytohormone concentrations. For instance, the concentrations of JA, IAA, ABA, and CTK enhanced by 5, 13.4, 6.3, and 12.5% at 40

mM, and 4.3, 21, 1.1, and 17% at 80 mM, and 7, 11, 6, and 14% at 120 mM, compared to the control treatments (**Figure 4 A-D**). In conclusion, jasmonic acid (JA) increased at 40 mM stress and decreased at 120 mM stress level.



Figure 6. Changes in the concentration of (A) Proline, (B) Soluble protein, (C) Soluble sugar, (D) Glycine betaine following salinity stress and N supplementation. The bars represent the mean data (n = 3) and SD. T1, T2, T3 and T4 represent 0, 40, 80, and 120 mM salinity stress, respectively. Following the LSD Tukey test, different letters indicate significant differences at P < 0.05.



Figure 7. Changes in the concentration of (A) NO_3^- , (B) NH_4^+ , and enzymatic activities of (C) NR, and (D) GS following salinity stress and N supplementation. The bars represent the mean data (n = 3) and SD. T1, T2, T3 and T4 represent 0, 40, 80, and 120 mM salinity stress, respectively. Following the LSD Tukey test, different letters indicate significant differences at P < 0.05.

3.5. Mitigation of Oxidative Stress Through N-Application Under Saline Conditions

Under salinity stresses, Vigna radiata L. plants produced a considerable amount of reactive oxygen species. The H₂O₂ and MDA contents increased by 1.3 and 1.4-fold at 40 mM, 1.9, and 1.5-fold under 80 mM, and 2.4 and 2.3-fold following 120 mM of salinity stress compared to the control treatment (Figure 5 A-B). Additionally, the N-application decreased the concentration of H_2O_2 and MDA under controlled and salinity stress treatments compared to their unfertilized pots. Additionally, the antioxidant enzymes such as SOD, CAT, POD, and APX improved by 1.1, 1.2, 1.05, and 1.2-fold at 40 mM, 1.3, 1.4, 1.4, and 1.3-fold at 80 mM, whereas 1.4, 1.46, 1.5, and 1.4-fold at 120 mM of salinity stress, respectively (Figure 5 C-F). Exogenous N supplementation further strengthened the enzymatic antioxidant mechanism. For instance, the N-application caused an increment of 1.1, 1.36, 1.2, and 1.15-fold at 40 mM, 1.3, 1.3, 1.5, and 1.2-fold at 80 mM, while 1.4, 1.5, 1.6, and 1.3-fold at 120 mM of salinity stress was noticed, respectively (Figure 5 C-F).

3.6. The negative impact of salinity on biochemical changes and its mitigation by N-application

The metabolites study showed a different pattern since proline, protein, and sugar contents were enhanced by 2.21-, 1.35-, and 1.08-fold at 40 mM of salt stress. Meanwhile, proline and protein improved 3.2- and 1.6-fold at 80 mM and 1.6- and 1.1-fold under 120 mM salinity stress compared to the control treatment (Figure 6 A-C). Furthermore, the sugar content faced a significant reduction at 80 and 120 mM in salt stress compared to the control. Like salt stress, the N-application improved the proline and protein content by 2.3- and 1.2-fold at 40 mM, whereas 2.6- and 1.1-fold at 80 mM salinity stress, respectively (Figure 6 A-B). On the other hand, at 120 mM of salt stress, N-application had no significant effect on proline and protein concentration (Figure 6 B). The application of N improved soluble sugar by 1.21 and 1.1fold under control and 80 mM stress, respectively (Figure 6 C). The concentration of GB decreased by 21.3, and 36.5% at 80 and 120 mM salinity stress levels compared to the control treatments (Figure 6 D). In contrast, N supplementation improved the GB concentration by 1.27, 1.54, and 1.11-fold at 40, 80, and 120 mM stress levels compared to their untreated peers (Figure 6 D).

3.7. Interactive impact of salinity and N-application on N metabolism

The NO₃⁻ and NH₄⁺ reduced up to 17 and 17% at 40 mM, while 60 and 44% at 80 mM, whereas 67 and 72% declined following 120 mM of salt stress, respectively (**Figure 7 A-B**). Furthermore, the results revealed that N-application enhanced the nutrient uptake, including NO₃ and NH₄^{+,} by 1.2-, 1.2-fold at 40 mM, 2.6-, and 1.8-fold at 80 mM salinity. Moreover, the NR and GS significantly reduced under all stress levels (**Figure 7 C-D**). Compared to the control treatment, both parameters were reduced to 21-and- 22% at 40 mM, 32- and 43% at 80 mM, and 62 and 63% at 120 mM of salinity stress. Similarly, N-application significantly enhanced the NR under control treatments (1.4-fold) compared to its untreated peer. Meanwhile, GS activity increased by 1.10, 1.27, and 2.1-fold at control, 40 mM, and 1.20 mM treatments, respectively (**Figure 7 C-D**).

3.8. Correlation analysis and principal component analysis

The correlation analysis revealed that SHL (shoot height length), SHFW (shoot fresh weight), SHDW (shoot dry weight), RTL (root length), RTFW (root fresh weight), and RTDW (root dry weight) exhibited a significant positive correlation (P < 0.05) with photosynthetic pigments such as Chl a, Chl b, and the Chl a/Chl b ratio, as well as with osmoregulatory compounds like proline and glycine betaine (Figure 8). These growth parameters also positively correlated with enzymes associated with N metabolism, including NR and GS, and N forms such as NO₃⁻ and NH₄⁺. On the contrary, a significant negative correlation was observed among antioxidant enzymes, including SOD, POD, CAT, and APX, with N assimilation enzymes (NR and GS). Moreover, H₂O₂ and MDA, oxidative stress markers, displayed a significant negative correlation with photosynthetic pigments and growth parameters, indicating that higher stress markers are associated with lower growth and pigment levels. Additionally, the plant hormones ABA showed a significant negative correlation with the growth parameters and photosynthetic pigments (Figure 8). At the same time, JA, IAA, and CTK exhibited a strong positive correlation, suggesting their role in promoting growth and pigment synthesis (Figure 8).

4. Discussion

The seedling stage in a crop's life cycle is vital and particularly vulnerable, and vigorous seedlings assure a high yield. At the early stage of rice seedlings, salt stress causes significant reductions in root number, root length, and shoot length (Hussain et al. 2017). Likewise, another experiment demonstrated that tomato seedlings treated with salt experienced a significant decline in growth attributes (such as plant height, reduced biomass, and reduced amounts of photosynthetic pigments) and gross photosynthesis (Singh et al. 2019). Similarly, we observed that the root-shoot length, fresh and dry weight as well and chlorophyll pigments significantly decreased when salinity stress was increased, which could be directly correlated with a proportionally increased cellular Na⁺ (Ullah et al. 2022a), H₂O₂ and MDA (Figure 4A-B). Supplemental N improved growth and reduced NaCl-associated toxicity by increasing chlorophyll pigments, K⁺, and the K⁺/Na⁺ ratio and enhancing N-metabolism and antioxidant activity.

Moreover, mungbean growth and biomass were improved by N supplementation under controlled conditions compared with untreated plants, indicating the high N demands of young seedlings with a more significant improvement in physiological and biochemical production. The biological systems of plants require adequate amounts of N for a wide range of vital metabolic functions (Gong *et al.* 2013; Raveena and Surendran, 2023). Consequently, N availability is central in regulating physiological mechanisms under unstressed and stressed conditions, including salinity. Excessive salinity leads to an imbalance of cellular ions, which leads to excess production of ROS, including H₂O₂. Under stressful environments, such as salinity, MDA is commonly used to assess cellular oxidative damage (Saidimoradi *et al.* 2019). As salinity stress levels increased, we noticed increased H_2O_2 and MDA contents in mung bean leaves. Our findings suggest that ROS interacts with macromolecules (proteins and lipids) and critical metabolic enzymes, leading to the degradation of photosynthetic performance and decreased mungbean plant growth and biomass. Under salinity stress, N supplementation to mungbean plants significantly reduced H_2O_2 and MDA contents, suggesting N can help protect membranes (Singh *et al.* 2019; Sikder *et al.* 2020).

Scavenging ROS during stressful conditions is essential for preventing oxidative damage (Zhao *et al.* 2022). In plants, antioxidant mechanisms enable them to eliminate ROS and minimize or prevent damage caused by ROS to cells (Li *et al.* 2024). Considering that N supplementation increases the antioxidant potential and salinity tolerance (Ahanger *et al.* 2019), we assessed the H₂O₂ and the SOD, APX, POD, and CAT activities. H₂O₂ levels were increased by salinity in comparison with the control. The N supplementation, however, inhibited H₂O₂ levels significantly compared to unfertilized plants. The decline in H₂O₂ and MDA could be attributed to the immediate strengthening of enzymatic antioxidants and osmolytes (such as proline, soluble protein, and soluble sugar) under exogenous N supplementation. It has been shown that N protects against oxidative stress induced by external or internal stresses (Sikder et al. 2020; Singh et al. 2019). The N may have a positive effect due to its role as a component of the antioxidant mechanism. Furthermore, several other studies have found that co-applying N significantly improved antioxidant enzyme levels in tomatoes (Singh et al. 2014), soybean (Borella et al. 2019), blueberry (Yañez-Mansilla et al. 2014), and cotton (Sikder et al. 2020), which corroborates our findings. Moreover, plants must regulate osmotic potential to minimize the adverse effects of osmotic stresses and dehydration. The osmolytes (such as glycine betaine, proline, soluble protein, and soluble sugar) play a critical role in maintaining salt balance, cell division, preventing ion toxicity, chlorophyll degradation, stabilizing cell structures, and eliminating excess ROS (Singh et al. 2022).



Figure 8. Analysis of correlations between all parameters examined. The colors blue and red indicate negative and positive correlations, respectively. Shoot height (SHL), shoot fresh weight (SHFW), shoot dry weight (SHDW), root length (RTL), root fresh weight (RTFW), root dry weight (RTDW), sodium (Na⁺), potassium (K⁺), chlorophyll a (ChI a), chlorophyll b (ChI b), chlorophyll a (chl a), chlorophyll b ratio (ChI a/ChI b), soluble sugar (SS), sodium/potassium ratio (Na⁺/K⁺), proline (Pro), soluble protein (SP), nitrate reductase (NR), glutamine synthetase (GS), nitrate (NO₃⁻), ammonium (NH₄⁺), ascorbate peroxidase (APX), and superoxide dismutase (SOD), malondialdehyde (MDA), hydrogen peroxide (H₂O₂), peroxidase (POD), catalase (CAT), including abscisic acid (ABA), indole acetic acid (IAA), cytokinin (CTK), and jasmonic acid (JA).

We noticed a significant increase in proline and soluble proteins under salinity stress, similar to previous research (Meriem *et al.* 2014; Sharif *et al.* 2018; Ullah *et al.* 2022a). Additionally, N application enhanced osmotic regulation by significantly increasing the osmoregulation substance. For instance, exogenous N supplementation improved the concentration of soluble protein, sugar, and proline. Another researcher observed improvements in relative water content, antioxidant potential, and osmolytes in

wheat subjected to salinity, including proline and lower membrane permeability (Agami *et al.* 2018).

Mungbean plants appear to use osmolytes to combat salinity stress, especially at low and medium stress (40 and 80 mM). The intersectional point is that despite the increased concentration of proline and soluble protein at all stress levels compared to the control, their concentration was more significant at 80 mM (80 > 40 > 120 > 0 mM NaCl). In contrast, the activity of the antioxidant enzyme increased as salinity stress levels (i.e., ascending). Hence, we suggest that mungbean seedlings spend more energy adjusting to osmotic conditions under low-tomedium salinity stress conditions. However, under high salinity stress, they concentrate on antioxidant defence mechanisms. Proline concentrations have been suggested to protect against salt damage/osmotic stress in plants by maintaining cellular osmotic potential. Additionally, they stabilize other functional units, such as membranes, electron transport complex II, enzymes, and proteins (Igbal et al. 2024a). Salt stress tolerance is boosted by the accumulation of soluble proteins, which can act as osmosis (Zhang et al. 2013; Sarker et al. 2018), corroborates our results. Proline, in addition to its function in osmotic regulation, also plays several other functions, including the scavenging of ROS, stabilizing subcellular structures, and buffering cellular redox potential under salinity stress conditions (Singh et al. 2015; Khanna-Chopra et al. 2019).

The growth-promoting hormone IAA is an important signalling molecule in plants. Auxin has been linked previously to salt stress responses in plants; however, it is unclear how these responses are controlled (Ryu and Cho 2015). The reduced IAA levels in salinity-treated mungbean plants in our study are consistent with previous findings on different plants species, including tomato (Albacete et al. 2008), Iris hexagona (Wang et al. 2001) and wheat (Shakirova et al. 2003). There may be a connection between this and the simultaneous suppression of growth in leaves and roots found in these studies. CKs also play a role in regulating plant growth and development, similar to auxins. Plant cytokinin levels indicate their ability to resist environmental stresses, making plants more tolerant of abiotic stresses such as high salinity and temperatures (Javed et al. 2023). A study found that increasing the level of CK in cereals improved their resistance to salinity stress (Iqbal et al. 2024b). It has been demonstrated in numerous studies that salt stress can lead to a reduction in overall CK levels (Vanková et al. 2014), which corroborates our findings. ABA promotes the closure of stomata and is a crucial stress indicator and mediator of the stress response in plants. In our study, all stress levels increased ABA content compared to controlled conditions. The increase in ABA levels following adverse environmental conditions has been reported in several previous publications (Albacete et al. 2008; Dobra et al. 2010; Vanková et al. 2014). Few studies have examined the effects of salt stress on endogenous JA levels. Compared to the control condition, the jasmonic content increased remained unchanged and decreased at 40-, 80-, and 120 mM stress levels, respectively. The current study revealed that N fertilization

enhanced the concentrations of JA, ABA, CTK and IAA under salinity stress levels compared to their untreated peers. In previous studies, N applications have been found to improve plant growth by regulating endogenous auxin levels (Ye et al. 2021). Plant hormones can be altered by both N and nitrate, which serve as N sources and signal molecules (Garnica et al. 2010; Kumar et al. 2018). Moreover, exogenous N fertilizer enhances the concentration of ABA and IAA in the crops by revealing a more significant fluctuation in growth and productivity (Borella et al. 2019). In conclusion, we suggest that N supplementation could enhance mungbean resistance to stress by upregulating endogenous salinity phytohormones, leading to improved growth compared to untreated plants.

The regulation of N metabolism plays a vital role in salt stress tolerance through several pathways (Arghavani et al. 2017; Yan et al. 2021). Plants use inorganic N from NO3⁻ and NH4⁺ for metabolic processes through their roots (Luo et al. 2013). Increasing salinity stress levels significantly reduced NO3⁻ ions in mungbean leaves. This could be attributed to NO₃⁻ uptake and loading of the root xylem is sensitive to salinity (Debouba et al. 2007). Plants absorb inorganic N, which can be converted into organic N through the action of NR, which is the starting factor and rate-limiting enzyme in the N metabolism. Thus, N accumulation and protein content positively relate to its activity. Through the GS/GOGAT cycle, approximately 95% of NH₄⁺ is converted to amino acids. The main form of N in plants and transportation is amino acids. GS is an essential enzyme in this cycle (Giunta et al. 2020).

Moreover, the foliar NR activity declined under increased salinity stress levels. Therefore, the decline in NO_3^- concentration may occur because of reduced NR activity or a direct reduction in enzyme synthesis under increased salinity stress (Iqbal 2018; Ullah *et al.* 2019). Hence, we suggest that the reduced NO_3^- uptake and accumulation may have affected N metabolism, which could have resulted in a significantly decreased amino acid and protein synthesis, thus considerably reducing the dry weight of mungbean plants under salinity stress (Xie *et al.* 2020). Further, a decreased NH₄⁺ was observed under increased salinity stress. In salt-induced NO₃⁻ reduction reactions, NH₄⁺ cannot be supplied to GS/GOGAT, inhibiting amino acid synthesis (Wang *et al.* 2012).

Moreover, the GS enzyme decreased with increased salinity stress. Several studies reported salt-induced reductions in plant NH₄⁺ through the impaired GS/GOGAT cycle (Wang *et al.* 2012; Meng *et al.* 2016; Ullah *et al.* 2019). Furthermore, a decrease in NH₄⁺ accumulation under salinity stress may result from gene downregulation related to the assimilation of NH₄⁺ due to NO₃⁻ deficiency, which reduces N assimilation (Wang *et al.* 2012). Furthermore, increased salinity stress reduced the ability of NO₃⁻ to be reduced (or NR- inhibition) and to be assimilated (GS), thus impairing N assimilation and amino acid metabolism and protein synthesis (Ullah *et al.* 2019). Therefore, biomass growth and accumulation are significantly reduced due to this process. In contrast, N

supplementation increased NO₃⁻, NH₄⁺ concentrations and NR and GS enzyme activities. A recent study demonstrated that N supplementation increases NR activity (Rehman et al. 2020), corroborating our findings. NO_{3⁻} and amino acids are controlled by plant cells through NR, which plays a critical role in the conversion of NO3- to amino acids. Because the availability of NO3 regulates NR activity, we suggest that an increase in NO3⁻ levels might enhance NR activity under N application. Increasing N-metabolizing enzymes can enhance plant N metabolism and facilitate plant protein synthesis and transformation (Chen et al. 2022). The uptake of water and minerals necessary for carbon and N metabolisms enhanced the osmotic regulation of mungbean plants under salinity stress due to the exogenous N supplementation. Therefore, the plants were able to grow under salinity conditions in a relatively normal manner compared to their untreated counterparts.

5. Conclusions

This study demonstrates that nitrogen supplementation effectively mitigates salinity stress in mungbean (Vigna radiata L.) through multi-faceted physiological improvements. Under 120 mM NaCl stress, we observed severe growth reductions (shoot-root biomass, 32-42% under 40 mM and 54-71% under 120 mM) that were alleviated by 1.4- 1.8-fold, 2.4- and 4-fold with N application. The treatment enhanced chlorophyll b and improved the K⁺/Na⁺ ratio (1.6-, 2.6- and 4.4-fold at 40-, 80and 120 mM, respectively). Moreover, growth hormones such as IAA and CTK improved by 13.4- and 12.5% at 40 mM, 21- and 17% at 80 mM, and 11- and 14% at 120 mM stress, respectively. Notably, N supplementation reduced oxidative stress markers (H₂O₂/MDA) while boosting antioxidant enzymes (SOD/CAT/POD increased 1-, 1.36-, and 1.2-, at 40 mM, 1.3-, 1.3-, and 1.5-, at 80 mM, while 1.4, 1.5, and 1.6-fold at 120 mM, respectively) and N metabolism. These findings establish that N application prioritizes metabolic repair over stress defense, offering a practical solution (50 mg/kg soil N) for saline agriculture. These findings establish a physiological framework for developing climate-resilient mungbean cultivation strategies in salinity-prone regions while highlighting the need for crop-specific nutrient management approaches in legume production systems. Future studies should elucidate the molecular mechanisms underlying nitrogenmediated salinity stress tolerance in mungbean, particularly focusing on nutrient-hormone crosstalk and stress adaptation. Moreover, field validation of optimized nitrogen supplementation regimes (timing, dosage, and formulation) under variable climatic conditions is essential for practical implementation.

Author contributions

Fu Shilin: Writing original draft; Software; Data curation; Babar Iqbal: Conceptualization; Funding acquisition; Investigation; Tahani Awad Alahmadi: Data curation; Formal analysis; Software; Mohammad Javed Ansari: Data curation; Writing - review & editing; Arsahd Jalal: Formal analysis; Writing - review & editing; Muhammad Junaid Nazir: Formal analysis; Software; Guanlin Li: Writing review & editing; Funding acquisition; Daolin Du: Resources; Writing - review & editing. All authors contributed to the article and approved the submitted version.

Declarations

Funding sources

This work was supported by the National Natural Science Foundation of China (32350410400), the International Science and Technology Cooperation Program of Jiangsu Province (BZ2024025), the Zhenjiang Municipal Policy Guidance Project of International Science and Technology Cooperation (Grant GJ2023005), the Postgraduate Research & Practice Innovation Program of Jiangsu Province (KYCX24_4012), and the Zhenjiang Major Project of Social Development in Research and Development (Grant SH2022004). This work was also supported by Zhenjiang Deweilepu Energy Environmental Protection Technology Co., Ltd. The authors wish to express their gratitude for the support.

Acknowledgement:

This project was supported by Researchers Supporting Project number (RSP2025R230), King Saud University, Riyadh, Saudi Arabia.

Data availability

All data are included in this article.

Conflict of interest

The authors declare no conflict of interest.

References

- Ahanger MA, Qin C, Begum N, *et al.* (2019). Nitrogen availability prevents oxidative effects of salinity on wheat growth and photosynthesis by up-regulating the antioxidants and osmolytes metabolism, and secondary metabolite accumulation. *BMC Plant Biology* **19**:1–12. doi:10.1186/s12870-019-2085-3
- Albacete A, Ghanem ME, Martínez-Andújar C, et al. (2008). Hormonal changes in relation to biomass partitioning and shoot growth impairment in salinized tomato (Solanum lycopersicum L.) plants. *Journal of Experimental Botany* 59:4119–4131.
- Arghavani M, Zaeimzadeh A, Savadkoohi S, Samiei L. (2017). Salinity tolerance of Kentucky bluegrass as affected by nitrogen fertilization. *Journal of Agricultural Science and Technology*. **19**:173–183.
- Azeem A, Javed Q, Sun J, et al. (2020). Functional traits of okra cultivars (Chinese green and Chinese red) under salt stress. Folia Horticulturae. 32:159–170. doi: 10.2478/fhort-2020-0015
- Borella J, Becker R, Lima MC, et al. (2019). Nitrogen source influences the antioxidative system of soybean plants under hypoxia and re-oxygenation. Sciences Agri 76:51–62. doi: 10.1590/1678-992x-2017-0195
- Bradford MM. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72:248–254. doi: 10.1016/0003-2697(76)90527-3
- Chen Y, Liu Y, Ge J, *et al.* (2022). Improved physiological and morphological traits of root synergistically enhanced salinity tolerance in rice under appropriate nitrogen application rate.

Frontiers in Plant Science **13**:982637. doi: 10.3389/fpls.2022. 982637

- Dai J, Duan L, Dong H. (2015). Comparative effect of nitrogen forms on nitrogen uptake and cotton growth under salinity stress. *Journal of Plant Nutrition* 38:1530–1543. doi: 10.1080/01904167.2014.983126
- Debouba M, Maâroufi-Dghimi H, Suzuki A, *et al.* (2007). Changes in growth and activity of enzymes involved in nitrate reduction and ammonium assimilation in tomato seedlings in response to NaCl stress. *Annals of Botany* **99**:1143–1151. doi: 10.1093/aob/mcm050
- Dobra J, Motyka V, Dobrev P et al. (2010). Comparison of hormonal responses to heat, drought and combined stress in tobacco plants with elevated proline content. *Journal of Plant Physiology* 167:1360–1370. doi: 10.1016/j.jplph.2010.05.013
- Garnica M, Houdusse F, Zamarreño AM, Garcia-Mina JM. (2010). The signal effect of nitrate supply enhances active forms of cytokinins and indole acetic content and reduces abscisic acid in wheat plants grown with ammonium. *Journal of Plant Physiology* **167**:1264–1272. doi: 10.1016/j.jplph.2010.04.013
- Giunta F, Mefleh M, Pruneddu G, Motzo R. (2021). Role of nitrogen uptake and grain number on the determination of grain nitrogen content in old durum wheat cultivars. *Agron*. 11:42. doi: 10.3390/agronomy11010042
- Gong B, Wen D, VandenLangenberg K, *et al.* (2013). Comparative effects of NaCl and NaHCO3 stress on photosynthetic parameters, nutrient metabolism, and the antioxidant system in tomato leaves. *Scientia Horticulturae* **157**:1–12. doi: 10.1016/j.scienta.2013.03.032
- Grieve CM, Grattan SR. (1983). Rapid assay for determination of water soluble quaternary ammonium compounds. *Plant and Soil* **70**:303–307. doi: 10.1007/BF02374789
- Guo JS, Zhou Q, Li XJ, Yu BJ, Luo QY. (2017). Enhancing NO3- supply confers NaCl tolerance by adjusting Cl- uptake and transport in G. max & G. soja. *Journal of Soil Science and Plant Nutrition* 17:194–202. doi: 10.4067/S0718-95162017005000015
- Hussain S, Zhang J, Zhong C, et al. (2017). Effects of salt stress on rice growth, development characteristics, and the regulating ways: A review. Journal of Integrative Agriculture 16:2357– 2374. doi: 10.1016/S2095–3119(16)61608-8
- Iqbal MJ. (2018). Role of osmolytes and antioxidant enzymes for drought tolerance in wheat. Global Wheat Production. *InTech.* doi: 10.5772/intechopen.75926
- Iqbal B, Zhao X, Khan KY, et al. (2024a). Microplastics meet invasive plants: Unraveling the ecological hazards to agroecosystems. Science of the Total Environment 906:167756. doi: 10.1016/j.scitotenv.2023.167756
- Iqbal B, Khan I, Anwar S, et al. (2024b). Biochar and saline soil: Mitigation strategy by incapacitating the ecological threats to agricultural land. International Journal of Phytomedicine doi: 10.1080/15226514.2024.2310001
- Irigoyen JJ, Einerich DW, Sánchez-Díaz M. (1992). Water stress induced changes in concentrations of proline and total soluble sugars in nodulated alfalfa (Medicago sativa) plants. *Physiologia Plantarum* 84:55–60. doi: 10.1111/j.1399-3054.1992.tb08764.x
- Javed Q, Sun J, Rutherford S, et al. (2023). Soil pollution and the invasion of congener Sphagneticola in crop lands. Journal of Environmental Management 340:118013. doi: 10.1016/j. jenvman.2023.118013

- Kant S. (2018). Understanding nitrate uptake, signaling and remobilisation for improving plant nitrogen use efficiency. In: *Seminars in Cell & Developmental Biology*. Elsevier, 89–96.
- Khanna-Chopra R, Semwal VK, Lakra N, Pareek A. (2019). Proline–A key regulator conferring plant tolerance to salinity and drought. In: Plant tolerance to environmental stress. *CRC Press*, 59–80.
- Kumar R, Mukherjee S, Ayele BT. (2018). Molecular aspects of sucrose transport and its metabolism to starch during seed development in wheat: a comprehensive review. *Biotechnology Advances* 36:954–967. doi: 10.1016/j.biotechadv.2018.02.015
- Khan W, Shah S, Ullah A, Ullah S, Amin F, et al. (2023). Utilizing hydrothermal time models to assess the effects of temperature and osmotic stress on maize (Zea mays L.) germination and physiological responses. BMC Plant Biology 23: 414. doi: 10.1186/s12870-023-04429-y
- Lichtenthaler HK, Buschmann C. (2001). Chlorophylls and carotenoids: Measurement and characterization by UV-VIS spectroscopy. *Current Protocols in Food Analytical Chemistry* F4.3.1 doi: 10.1002/0471142913.faf0403s01
- Li G, Cui X, Tariq M, *et al.* (2024). Microplastic and cadmium contamination: Impact on the soil by inhibiting the growth of pak choi (Brassica rapa subsp. chinensis). *Process Safety and Environmental Protection* **189**:714–727. doi: 10.1016/j.psep. 2024.06.081
- Liu C, Wang Y, Pan K, et al. (2014). Carbon and nitrogen metabolism in leaves and roots of dwarf bamboo (Fargesia denudata Yi) subjected to drought for two consecutive years during sprouting period. *Journal of Plant Growth Regulation* 33:243–255. doi: 10.1007/s00344-013-9367-z
- Liu RX, Chen JA, Ren ZR, *et al.* (2023). Leaf traits of clonal grasses responding to the ratios of ammonium to nitrate in a semiarid grassland: Leaf order matters. *Journal of Plant Ecology* **16**(4):rtac108. doi: 10.1093/jpe/rtac108
- Luo J, Li H, Liu T, et al. (2013). Nitrogen metabolism of two contrasting poplar species during acclimation to limiting nitrogen availability. Journal of Experimental Botany 64:4207– 4224. doi: 10.1093/jxb/ert234
- Ma H, Cui LJ, Li W, et al. (2022). Effect of daily salinity fluctuation on the intraspecific interactions of a euhalophyte (suaeda salsa) along a salinity gradient. Journal of Plant Ecology 15(1):208–221. doi: 10.1093/jpe/rtac002
- Meng S, Su L, Li Y, et al. (2016). Nitrate and ammonium contribute to the distinct nitrogen metabolism of Populus simonii during moderate salt stress. PLOS One 11: e0150354. doi: 10.1371/journal.pone.0150354
- Meriem BF, Kaouther Z, Chérif H, *et al.* (2014). Effect of priming on growth, biochemical parameters and mineral composition of different cultivars of coriander (Coriandrum sativum L.) under salt stress. *Journal Of Stress Physiology And Biochemistry* **10**:84–109.
- Mignolet-Spruyt L, Xu E, Idänheimo N, et al. (2016). Spreading the news: subcellular and organellar reactive oxygen species production and signalling. Journal of Experimental Botany 67:3831–3844. doi: 10.1093/jxb/erw080
- Patterson BD, MacRae EA, Ferguson IB. (1984). Estimation of hydrogen peroxide in plant extracts using titanium (IV). Analytical Biochemistry **139**:487–492
- Raveena S, Surendran R, Sameer A, et al. (2024). Empowering coffee farming using counterfactual recommendation based RNN driven IoT integrated soil quality command system. *Scientific Reports* 14, 6269. doi: 10.1038/s41598-024-56954-x.

- Raveena S and Surendran R. (2023). Recommending the Right Biofertilizer Using Deep Collaborative Matrix Factorization in the Coffee Plantation. 7th International Conference on Electronics, Communication and Aerospace Technology (ICECA), Coimbatore, India, 2023, 476–482, doi: 10.1109/ICECA58529.2023.10395096.
- Ravi S, Bader MKF, Young T, et al. (2022). Are the well-fed less thirsty? Effects of drought and salinity on New Zealand mangroves. Journal of Plant Ecology 15(1):85–99. doi: 10.1093/jpe/rtab071
- Rehman M, Yang M, Fahad S, et al. (2020). Morpho-physiological traits, antioxidant capacity, and nitrogen metabolism in ramie under nitrogen fertilizer. Agronomy Journal **112**:2988–2997. doi: 10.1002/agj2.20212
- Ryu H, Cho YG. (2015). Plant hormones in salt stress tolerance. Journal of Plant Biology 58:147–155. doi: 10.1007/s12374-015-0103-z
- Sabra A, Daayf F, Renault S. (2012). Differential physiological and biochemical responses of three Echinacea species to salinity stress. *Scientia Horticulturae* **135**:23–31. doi: 10.1016/j.scienta.2011.11.024
- Saidimoradi D, Ghaderi N, Javadi T. (2019). Salinity stress mitigation by humic acid application in strawberry (Fragaria x ananassa Duch.). *Scientia Horticulturae* 256:108594. doi: 10.1016/j.scienta.2019.108594
- Sánchez-Rodríguez E, del Mar Rubio-Wilhelmi M, Ríos JJ, et al. (2011). Ammonia production and assimilation: its importance as a tolerance mechanism during moderate water deficit in tomato plants. *Journal of Plant Physiology* **168**:816–823. doi: 10.1016/j.jplph.2010.11.018
- Sahab S, Suhani I, Srivastava V, et al. (2021). Potential risk assessment of soil salinity to agroecosystem sustainability: current status and management strategies. Science of the Total Environment 764:144164. https://doi.org/10.1016/j.scitotenv.2020.144164.
- Sarker U, Ercisli S. (2022). Salt eustress induction in Red Amaranth (Amaranthus gangeticus) augments nutritional, phenolic acids and antiradical potential of leaves. *Antioxidants* 11:2434. doi: 10.3390/antiox11122434
- Sarker U, Hossain MN, Oba S, *et al.* (2023). Salinity stress ameliorates pigments, minerals, polyphenolic profiles, and antiradical capacity in Lalshak. *Antioxidants* **12**:173. doi: 10.3390/antiox12010173
- Sarker U, Islam MT, Oba S. (2018). Salinity stress accelerates nutrients, dietary fiber, minerals, phytochemicals and antioxidant activity in Amaranthus tricolor leaves. *PLOS One* **13**: e0206388. doi: 10.1371/journal.pone.0206388
- Sehrawat N, Yadav M, Sharma AK, et al. (2019). Salt stress and mungbean [Vigna radiata (L.) Wilczek]: effects, physiological perspective and management practices for alleviating salinity. Archives of Agronomy and Soil Science 65:1287–1301. https://doi.org/10.1080/03650340.2018.1562548
- Shakirova FM, Sakhabutdinova AR, Bezrukova MV, et al. (2003). Changes in the hormonal status of wheat seedlings induced by salicylic acid and salinity. *Plant Science* 164:317–322. doi: 10.1016/S0168-9452(02)00415-6
- Sharif P, Seyedsalehi M, Paladino O, et al. (2018). Effect of drought and salinity stresses on morphological and physiological characteristics of canola. International Journal of Environmental Science and Technology 15:1859–1866. doi: 10.1007/s13762-017-1508-7

- Siddiqui MH, Mohammad F, Khan MN, et al. (2010). Nitrogen in relation to photosynthetic capacity and accumulation of osmoprotectant and nutrients in Brassica genotypes grown under salt stress. Agricultural Sciences in China 9:671–680. doi: 10.1016/S1671-2927(09)60142-5
- Sikder RK, Wang X, Zhang H, *et al.* (2020). Nitrogen enhances salt tolerance by modulating the antioxidant defense system and osmoregulation substance content in Gossypium hirsutum. *Plants* **9**:450. doi: 10.3390/plants9040450
- Singh M, Kumar J, Singh S, et al. (2015). Roles of osmoprotectants in improving salinity and drought tolerance in plants: a review. Reviews in Environmental Science and Biotechnology 14:407–426. doi: 10.1007/s11157-015-9372-8
- Singh M, Kumar J, Singh VP, Prasad SM (2014) Plant tolerance mechanism against salt stress: the nutrient management approach. *Biochemical Pharmacology* 3:e165. doi: 10.4172/2167-0501.1000e165
- Singh M, Singh VP, Prasad SM. (2019). Nitrogen alleviates salinity toxicity in Solanum lycopersicum seedlings by regulating ROS homeostasis. *Plant Physiology and Biochemistry* 141:466– 476. doi: 10.(1016)/j.plaphy.2019.04.004
- Singh P, Choudhary KK, Chaudhary N, et al. (2022). Salt stress resilience in plants mediated through osmolyte accumulation and its crosstalk mechanism with phytohormones. Frontiers in Plant Science 13:1006617. doi: 10.3389/fpls.2022.1006617
- Solangi KA, Siyal AA, Wu Y, *et al.* (2019). An Assessment of the Spatial and Temporal Distribution of Soil Salinity in Combination with Field and Satellite Data: A Case Study in Sujawal District. *Agronomy*. **9**(12):869. doi: 10.3390/agronomy9120869
- Teshita A, Khan W, Ullah A, *et al.* (2024). Soil nematodes in agroecosystems: Linking cropping system's rhizosphere ecology to nematode structure and function. *Journal of Soil Science and Plant Nutrition* **24**, 6467–6482. doi: 10.1007/s42729-024-01982-9
- Ullah A, Li M, Noor J, et al. (2019a). Effects of salinity on photosynthetic traits, ion homeostasis and nitrogen metabolism in wild and cultivated soybean. PeerJ 7:e8191. doi: 10.7717/peerj.8191
- Ullah A, Tariq A, Sardans J, *et al.* (2022a). Alhagi sparsifolia acclimatizes to saline stress by regulating its osmotic, antioxidant, and nitrogen assimilation potential. *BMC Plant Biology* **22**:1–17. doi: 10.1186/s12870-022-03832-1
- Ullah A, Zeng F, Tariq A, *et al.* (2022b). Exogenous naphthaleneacetic acid alleviated alkalinity-induced morphophysio-biochemical damages in Cyperus esculentus L. var. sativus Boeck. *Frontiers in Plant Science* **13**:1018787. doi: 10.3389/fpls.2022.1018787
- Ullah I, Mao H, Shabbir A, et al. (2020). Physiological response of tomato plants under different irrigation levels and nutrient concentrations in greenhouse. Pakistan Journal of Agricultural Sciences 57:599–608. doi: 10.21162/PAKJAS/19.844
- Vanková R, Kosová K, Dobrev P, et al. (2014). Dynamics of cold acclimation and complex phytohormone responses in Triticum monococcum lines G3116 and DV92 differing in vernalization and frost tolerance level. *Environmental and Experimental Botany* **101**:12–25. doi: 10.1016/j.envexpbot.2014.01.002
- Wang H, Wu Z, Zhou Y, *et al.* (2012). Effects of salt stress on ion balance and nitrogen metabolism in rice. *Plant, Soil and Environment* **58**:62–67. doi: 10.17221/615/2011-PSE
- Wang J, Guo X, Zhao Q, et al. (2025). Subsurface drip irrigation with micro-nano bubble hydrogen water improves the salt

tolerance of lettuce by regulating the antioxidant system and soil bacterial community. *Applied Soil Ecology* **207**: 105948. doi: 10.1016/j.apsoil.2025.105948

- Wang J, Zhong XM, Lv XL, et al. (2018). Photosynthesis and physiology responses of paired near-isogenic lines in waxy maize (Zea mays L.) to nicosulfuron. *Photosynthetica* 56:1059–1068. doi: 10.1007/s11099-018-0816-6
- Wang Y, Mopper S, Hasenstein KH. (2001). Effects of salinity on endogenous ABA, IAA, JA, and SA in Iris hexagona. J Chem Ecol. 27:327–342. doi: 10.1023/A:1005632506230
- Wani SH, Kumar V, Khare T, *et al.* (2020). Engineering salinity tolerance in plants: progress and prospects. Planta 251: 1–29. https://doi.org/10.1007/s00425-020-03366-6.
- Wei W, Zhang X, Hou Z, et al. (2021). Microbial regulation of deterioration and preservation of salted kelp under different temperature and salinity conditions. Foods. 10(8):1723. doi: 10.3390/foods10081723
- Wang Z, Li X, Zhu X, et al. (2017). Salt acclimation induced salt tolerance is enhanced by abscisic acid priming in wheat. Plant Soil Environ. 63(7): 307–314. doi: 10.17221/287/2017-PSE

World Vegetable Centre (2018) Shanhua, tainan, Taiwan 74151.

- Xie W, Chen Q, Wu L, et al. (2020). Coastal saline soil aggregate formation and salt distribution are affected by straw and nitrogen application: a 4-year field study. Soil Tillage Res. 198:104535. https://doi.org/10.1016/j.still.2019.104535.
- Yan SH, Gao YM, Tian MJ, et al. (2021). Comprehensive evaluation of effects of various carbon-rich amendments on tomato production under continuous saline water irrigation: overall soil quality, plant nutrient uptake, crop yields and fruit quality. Agric Water Manage. 255:106995. https://doi.org/10.1016/j.agwat.2021.106995

Yañez-Mansilla E, Cartes P, Díaz MR, et al. (2014). Photosynthetic

concentrations, Δ 13C, and δ 15N of durum wheat and related amphiploids. *Journal of Experimental Botany* 61:3529–3542. doi: 10.1093/jxb/erq184

- Zaid A, Wani SH. (2019). Reactive oxygen species generation, scavenging and signaling in plant defense responses. In: Bioactive molecules in plant defense. Springer, 111–132.
- Zhang C, Li X, Yan H, et al. (2020). Effects of irrigation quantity and biochar on soil physical properties, growth characteristics, yield and quality of greenhouse tomato. Agri Water Manage. 241:106263. doi: 10.1016/j.agwat.2020.106263
- Zhang JL, Shi H. (2013). Physiological and molecular mechanisms of plant salt tolerance. Photosynthesis Res. 115:1–22. doi: 10.1007/s11120-013-9813-6
- Zhang M, Fang Y, Ji Y, et al. (2013). Effects of salt stress on ion content, antioxidant enzymes and protein profile in different tissues of Broussonetia papyrifera. South Afr J Bot. 85:1–9. doi: 10.1016/j.sajb.2012.11.005
- Zhao X, Xie H, Zhao X, et al. (2022). Combined inhibitory effect of Canada goldenrod invasion and soil microplastics on rice growth. Inter J Environ res Public Health. 19 (19), 11947. doi: 10.3390/ijerph191911947
- Zhou Y, Lam HM, Zhang J. (2007). Inhibition of photosynthesis and energy dissipation induced by water and high light stresses in rice. *Journal of Experimental Botany* 58:1207–1217. doi: 10.1093/jxb/erl291
- Zhu JK. (2001). Plant salt tolerance. *Trends in Plant Science* 6:66– 71. doi: 10.1016/S1360–1385(00)01838-0
- Zuo Z, Ye F, Wang Z, *et al.* (2021). Salt acclimation induced salt tolerance in wild-type and chlorophyll b-deficient mutant wheat. Plant Soil Environ. 67(1):26–32. doi: 10.17221/429/2020-PSE

sand antioxidant performance are differentially affected by short-term nitrogen supply in highbush blueberry cultivars. Cienc Inv Agr. 41:61–70. doi: 10.4067/S0718-16202014000100006

- Ye D, Shen Q, Guo Y, et al. (2021). Sufficient nitrogen promoted high phosphorus tolerance and phosphorus-accumulating capability of Polygonum hydropiper in relation to changes of phytohormones and phenols. Chemosphere 278:130318. doi: 10.1016/j.chemosphere.2021.130318
- Yousfi S, Serret MD, Voltas J, Araus JL. (2010). Effect of salinity and water stress during the reproductive stage on growth, ion