

Biomonitoring of soil quality in the Souk Ahras region (Northeast Algeria) using *Cornu aspersum* as a bioindicator: Trace metals and sulfates analysis

Bendjebbar Roumaïssa^{1*}, Berrouk Houda¹, Barour Choukri¹, Sleimi Noomene² and Merghad Amina¹

¹Laboratory of Aquatic and Terrestrial Ecosystems (LEAT), Department of Biology, Faculty of Natural and Life Sciences, Mohamed Chérif Messaâdia University, BP1553, Annaba Road, Souk Ahras, Algeria.


²Laboratory of Resources, Materials and Ecosystems (LR19ES20), Faculty of Sciences of Bizerte, University of Carthage, Tunisia.

Received: 31/01/2025, Accepted: 25/04/2025, Available online: 05/05/2025

*to whom all correspondence should be addressed: e-mail: r.bendjebbar@univ-soukahras.dz

<https://doi.org/10.30955/gnj.07315>

Graphical abstract



Abstract

The present study aims to assess the impact of physicochemical parameters on the availability of trace metal elements (TME) in soil by analyzing the accumulation of copper (Cu), zinc (Zn) and sulfates in both soil and edible tissues of *Cornu aspersum* using a passive biomonitoring approach at three sites in Souk Ahras province: Zaarouria, Mechrouha and ENAP (National Paint Enterprise). Statistical analysis of the data indicated that adsorption of TME and sulfates on soil particles inhibits their uptake by other terrestrial organisms such as snails. In addition, physicochemical parameters significantly influenced the distribution of TME in the soil, revealing marked spatial differences in the accumulation of Cu in the foot and viscera as well as Zn especially in the viscera. Furthermore, a significantly higher accumulation of sulfates was observed in the foot compared to the viscera especially at the ENAP site. These observations could confirm the potential of *C. aspersum* species as an effective bioindicator of metal pollution in soil.

Keywords: *Cornu aspersum*, bioindicator, bioavailability, accumulation, metal pollution, soil

1. Introduction

One of the main areas of current research is environmental pollution caused by the release of different types of

pollutants into water, air and soil (Baroudi 2020; N'Doua *et al.* 2024). These different pollutants originate from the expansion of human activities and chemical emissions, notably trace metal elements (TME) associated with industrial and agricultural needs (Dossou *et al.* 2022; Amlan *et al.* 2023; Hidouri *et al.* 2024). TME occur naturally in the environment and are considered among the most harmful pollutants, as they are non-biodegradable and their concentration is constantly increasing in different ecosystem compartments (Ibrahim *et al.* 2019; Ning *et al.* 2024). Due to their excessive presence in the ecosystem, heavy metals can also have a direct impact on the life cycles of animal and plant organisms, both terrestrial and aquatic. This dynamic could lead to a reduction in their density and abundance. This creates an imbalance in the food chain and poses a threat to biodiversity. As a result, the ecosystem becomes more vulnerable to environmental changes (Ibrahim *et al.* 2023; Shaffique *et al.* 2024). High concentrations of TME can cause serious problems for organisms, particularly due to the increased sensitivity of invertebrates to TME compared to vertebrates (Monchanin *et al.* 2021; Benhamdoun *et al.* 2024), such as disruption of the physiological balance in earthworms, as well as in snails and beetles resulting in an increase or decrease of certain oxidative stress biomarkers such as AchE, GST, MDA and antioxidant enzymes. This is due to the accumulation of TME in their tissues, which is directly caused by the presence of these metals in their natural environment, be it water, plants or soil (Du *et al.* 2019; Bennour and Soltani 2020; Larba *et al.* 2023; Bouzahouane *et al.* 2024; Ogbeide and Amayanbo 2024).

Currently, assessing the bioavailability of metals in soil and studying the multiple effects and hazards of polluted soils are a crucial issue in ecotoxicology (Chen *et al.* 2023; Selvanarayanan *et al.* 2024a) in order to protect potentially exposed populations, such as living fauna and flora organisms (Alsherif *et al.* 2022; El Mageed *et al.* 2023; Bici *et al.* 2024). Recent developments in real-time environmental monitoring using IoT-based systems have

shown promise in improving pollution management and water quality assessment (Maruthai *et al.* 2025; Periasamy *et al.* 2024). As well as in predicting and classifying water quality using advanced deep learning models such as Attention-based Deep Differential RecurFlowNet (Enkatraman *et al.* 2025).

Soil is considered to be a habitat for numerous microorganisms, plants and terrestrial fauna and their interactions within which diverse life communities develop and reproduce. It also provides a capacity to retain pollutants to prevent their transport especially by water, and subsequent transfer into food chains (Owojori *et al.* 2022; Soliman *et al.* 2022; Li *et al.* 2023; Mwelwa *et al.* 2023). Different species are able to indicate soil quality, among which we find terrestrial gastropods, due to their direct contact with the soil and their high capacity to accumulate TME, they are considered the most reliable bioindicators (Guessasma *et al.* 2020; Salih *et al.* 2021; Cheriti 2022; Nwagu *et al.* 2022; Owojori *et al.* 2022; Sargsyan *et al.* 2022; Ajayi and Oyewole 2023; Louzon *et al.* 2023). Due to its sedentary lifestyle and diet, it is easily contaminated by various pollutants present in the ecosystem (Al-Alam *et al.* 2024). The assessment of metal bioavailability in soils can be performed using this indicator in situ by active bioindication (by confining sentinel snail species from in vitro breeding, thus making them with a known biological history) or passive bioindication (based on direct sampling of wild snails from their habitats, which therefore have an unknown biological history) (Al-Alam *et al.* 2022; Sahraoui 2022). In a context where environmental conservation is a global concern, it is crucial to understand the mechanisms of metal contamination in ecosystems and the main reasons for it (Selvanarayanan *et al.* 2024b). How does this contamination affect human health, ecosystems and local biodiversity?

This study aims to monitor the distribution of TME in the soil of three different sites in the Souk Ahras province which are Zaarouria, Mechrouha and the paint factory (ENAP) under the influence of soil physicochemical parameters. This work is driven by growing environmental concerns over heavy metal contamination in regions affected by industrial and agricultural activities, and the need for effective biomonitoring tools to assess soil quality. We focused our

survey on measuring rates of two metals, copper (Cu) and Zinc (Zn) in the foot and viscera of *Cornu aspersum* as accumulation organs. In addition, a comparative analysis of sulfate concentrations was carried out between sites and snails' tissues of both foot and viscera using a passive biomonitoring approach which allowed us to assess the level of snail contamination by TME at the investigated sites and thereby evaluate soil quality.

2. Materials and methods

2.1. Sampling sites

This study was carried out in the geographical region of the extreme northeast of Algeria, which includes three different sampling sites in Souk Ahras province, bounded to the northwest by the cities of El Tarf and Guelma and to

the southwest by Oum El Bouaghi city, with Tébessa to the southeast and Tunisia to the east. Our sampling area is characterized by its mountainous terrain, it is also characterized by its topographical diversity, with altitudes ranging from 1000 m in the north to 650 m in the south. The area is divided into two geographical regions: in the north, mountainous and forests are distributed, characterized by a subhumid climate and siliceous or limestone soils, with an average annual rainfall of approximately 730 mm. On the other hand, the south is composed of high plains and pastures, with a semi-arid climate and brownish agricultural soils, often covered with a layer of limestone, with an average annual rainfall of 350 mm per year (Boudy 1955).

In addition, the sampling sites were selected according to a gradient of soil contamination compared to their proximity (distances) to factories of different industries. An area without any industrial activity was also selected, taking into account their accessibility, the ease of sampling and the abundance of *C. aspersum* species. Three monitoring sites were selected, as illustrated in **Figure 1**:

ENAP: National Enterprise of Paints of Souk Ahras located near the national road N° 16 (36° 17' 21" N, 7° 56' 19" E), the factory is responsible for manufacturing paints by using several raw materials: binders, pigments, solvents and glues.

Mechrouha: is located 21 km northwest of Souk Ahras city and 58 km southeast of Guelma (36° 21' 0" N, 7° 49' 60" E), this region is characterized by its rugged terrain, it consists mainly of mountains, contains stunning natural sites and is known for its dense forest cover.

Zaarouria: is located 10 km west of Souk Ahras city (36° 13' 38" N, 7° 57' 28" E), it comprises a natural mountainous area, an urban and an industrial area characterized by the presence of a brick factory. It is also distinguished by its variety of trees, including vineyards.




Figure 1. Map showing the sampling sites of *C. aspersum* in Souk Ahras province (northeastern Algeria).

2.2. Experimental approach

The use of sentinel bioindicator species has become a widely recognized phenomenon for assessing soil quality in recent decades. The bioindication method used in this study is passive bioindication. This approach is based on the principle that terrestrial gastropods due to their sensitivity to environmental changes can serve as reliable indicators of soil quality, thus allowing the monitoring of

concentrations of various analyses including metals (Dupont *et al.* 2023; Larba *et al.* 2023)

2.3. Biological model

The biological material used is a terrestrial gastropod, the *C. aspersum* snail (Müller 1774) randomly collected from three sites in the Souk Ahras province (northeastern Algeria). The snails (with an average weight of 8 ± 0.5 g) were placed in aestivation under dry conditions, then they were woken up by spraying with demineralized water and fed with fresh lettuce for two days before being deprived of food for another two days to ensure that their intestinal tract was empty.

2.4. Soil physicochemical characteristics

A physicochemical analysis was carried out on three soil samples collected from three different sites to determine common soil characteristics, in particular: hydrogen potential (pH) according to Baize and Jabiole (1995), electrical conductivity (EC) according to (Okalebo *et al.* 2002), salinity (Rhoades *et al.* 1982), nitrogen (N) based on Kjeldahl method (1883), phosphorus (P) according to Olsen *et al.* (1954), carbon (Walkley and Black 1934), total limestone content (Drouineau 1942), organic matter (OM) (Bonnefont *et al.* 1980), cation exchange capacity (CEC) according to Thomas (1982) and soil texture (Lag *et al.* 2008). This analysis was carried out at the Horizon private laboratory in Annaba city.

2.5. TME extraction by hot acid mineralization

After snails' dissection and drying of the viscera and feet of *C. aspersum* in an oven at 65 °C for 48 to 72 hours, the dry organic matter was ground to a fine powder using a ceramic mortar and pestle. 100 mg of dry matter was digested in 3 ml of an acid mixture consisting of: HNO₃: H₂SO₄: HClO₄ in a volume ratio of 10: 1: 0.5 at 110 °C for 2 h in Teflon bombs.

The extracts were recovered in 50 ml of 0.5% (v/v) HNO₃, then filtered through Whatman N° 1 filter paper and injected into the atomic absorption spectrometer (PinAAcle 900T, PerkinElmer, Waltham, MA, USA) to determine the content of the two metals (Cu and Zn). The Cu and Zn standards used to calibrate the instrument were prepared using PerkinElmer solutions of 1000 mg L⁻¹ (PerkinElmer, Waltham, MA, USA). The concentration range varied depending on the nature of element being analyzed and its expected abundance in the mineralized samples. The calibration ranges for Cu and Zn were 0-5 mg L⁻¹ and 0-1 mg L⁻¹ respectively (Sleimi *et al.* 2022; Bankaji *et al.* 2023; Bouzahouane *et al.* 2024).

2.6. Soil analysis by X-ray fluorescence spectrometry

The analyses were carried out at the common service unit for research "Atomic Absorption Spectrometer" and the Research Laboratory RME-Resources, Materials and Ecosystems, University of Carthage, Tunisia.

The studies presented here were carried out using an X-ray fluorescence (XRF) spectrometer. This analyzer offers the highest performance in XRF analysis, providing the lowest possible detection limits for nearly forty elements. It is specifically designed for the analysis of solid metals, alloys,

ores and soils. This last feature was implemented for our study. The analysis time lasts 30 s per sample. The results are expressed in parts per million (ppm), and they are particularly positive, allowing the direct use of the copper and zinc concentrations.

Soil samples (5 cm deep) were collected from the ENAP, Zaarouria and Mechrouha sites. Once collected, the soil to be analyzed must first be dried in an oven at 65 °C, then ground and sieved to a mesh size of 2 mm. They are then placed in capsules and covered with Mylar film. For analysis, the samples are placed in an automatic feeder of the XRF spectrometer (Téreygeol *et al.* 2010; Guessasma *et al.* 2020; Weindorf and Chakraborty 2020; Kim *et al.* 2023).

2.7. Sulfates analysis

For the quantification of sulfates, a photometric method was employed, sulfates are precipitated in a hydrochloric acid medium, forming barium sulfate. After mineralization of our samples and preparation of stock solutions, a precise dosing technique was followed. In test tubes, 2.5 ml of the solution (test sample) was combined with 17 ml of distilled water, 0.5 ml of HCL and 2.5 ml of the 20% BaCl₂ + PVP solution. These mixtures were vigorously stirred at a constant speed for exactly 1 minute, followed by a resting period of 15 minutes. The spectrometer cuvettes were then filled and the turbidity measured 30 seconds later at a wavelength of 650 nm. The sulfates transformed into suspended barium sulfate in a range solution were distributed in 6 tubes: 0-15-21-27-30-90 mg L⁻¹ of SO₄⁻². This detailed methodology ensures a meticulous analysis of the sulfates present in the sample (Gboko *et al.* 2022).

2.8. Statistical analysis

In the present study, all our statistical analyses were performed using R, version 4.2.2 (R Core Team 2022; Ihaka and Gentleman 1996) for MacOS (<http://cran.r-project.org>). Shapiro-Wilk test was used to test variables normality. Data were expressed as mean $\pm$ standard error (se). The comparisons between sites for the soil physicochemical characteristics and between tissues for trace metal elements (TME) and sulfates were assessed by the nonparametric Kruskal-Wallis (KW) rank sum test. The KW test was followed by the nonparametric pairwise Dunn's test (with Bonferroni adjusted *p*-value) to find post-hoc statistical differences at $\alpha = 0.05$ as significant level. Correlations between the analysed variables were also calculated by using Spearman's nonparametric correlation (with Bonferroni adjusted *p*-value). Finally, in order to characterise the soil samples according to their physicochemical characteristics and their possible impacts on the TME and sulfates concentrations in the snail's tissues; we carried out also a multivariate analysis by applying a principal component analysis (PCA) as ordination technique. Several R packages (libraries) were also used in our statistical analyses and to plot data results such as 'FactoMineR' (Lê *et al.* 2008), 'ggplot2' (Wickham 2016), 'dunn.test' (Dinno 2017), 'factoextra' (Kassambara and Mundt 2020), 'Hmisc' (Harrell 2023), 'ggcorrplot' (Kassambara 2023a), 'ggpubr' (Kassambara 2023b), 'psych' (Revelle 2023) and 'PMCMRplus' (Pohlert 2023).

3. Results

3.1. TME and sulfates analysis across distinct sites

We used the Principal Component Analysis (PCA) because it is a real versatile statistical method. In our study we applied it as an ordination technique and it allowed us to reduce the dimensionality of our dataset (based on 21 variables) and to preserve the most important patterns or relationships between the analyzed variables according to the target sites. The PCA provided an approximation of our original data on *C. aspersum* snails' TME: copper (CuF and CuV) and zinc (ZnF and ZnV) as well as sulfates (Sulf. F and Sulf. V) within three distinct sites in the Souk Ahras province (Zaarouria, Mechrouha and ENAP). In our case, the PCA enabled us the assessment of correlations between snails' TME (blue variables on the biplot, **Figure 2**) and with other soil variables (TME and texture-physicochemical parameters): copper (Cu), zinc (Zn), sulfates (Sulf), sand, silt, clay, cation exchange capacity (CEC), organic matter (OM), total lime (T. Lime), carbon (C), phosphorus (P), nitrogen (N), salinity (Sal.), electrical conductivity (EC) and pH. In a synthetic way, the PCA results indicate that the first two axes explain 100% of the total inertia (data variance) present in our dataset, indicating that the biplot represents the entirety of the variability (**Figure 2**).




Figure 2. Principal component analysis (PCA) biplot of the three sampled site characterization (Dim 1: 68.5% and Dim 2 = 31.5%).

The biplot shows the PCA scores of the fifteen explanatory variables as black arrows. Points on the same side as a given variable should be interpreted as having a high contribution on it. In this study, the six TME measured in snails' tissues of *C. aspersum* were treated as supplementary variables (the six blue arrows).

On the one hand, the 1st axis (Dim 1) alone, explained 68.5% of the total variation and showed strongly positive correlation especially with: Sulf ($r = 1$; $\cos^2 = 1$), CEC ($r = 0.95$; $\cos^2 = 0.91$), clay ($r = 0.95$; $\cos^2 = 0.90$), P ($r = 0.88$; $\cos^2 = 0.78$), Silt ($r = 0.85$; $\cos^2 = 0.73$), OM ($r = 0.72$; $\cos^2 = 0.52$), C ($r = 0.72$; $\cos^2 = 0.52$) and Sal. ($r = 0.71$; $\cos^2 = 0.71$). In addition, the 1st axis is also negatively correlated with the following variables: N ($r = -1$; $\cos^2 = 1$), Cu ($r = -1$; $\cos^2 =$

0.99), pH ($r = -0.90$; $\cos^2 = 0.80$) and sand ($r = -0.87$; $\cos^2 = 0.76$). On the other hand, the 2nd axis (Dim 2) just explained 31.5% of the total variation and it was also positively correlated with the variables: T. Lime ($r = 0.88$; $\cos^2 = 0.78$), EC ($r = 0.72$; $\cos^2 = 0.52$) and Sal. ($r = 0.71$; $\cos^2 = 0.50$) and as well as negative correlations with the variable Zn ($r = -0.91$; $\cos^2 = 0.83$).

Moreover, PCA showed different patterns of correlations between analyzed variables and the results revealed an important differentiation between sites due to fluctuations in the concentrations of various TME measured values from the soil as well as in the foot and viscera of *C. aspersum*.

The 1st axis generally distinguished Zaarouria and ENAP sites from the Mechrouha one which is characterized by silty-clay soil and high concentrations values, mostly of Sulf, CEC, OM, C, P, Sal., CuV, ZnV and CuF. By contrast, it exhibited low values for Cu, Sand, N and pH concentrations of Mechrouha in comparison with the two other sites. Furthermore, the 2nd axis mainly differentiated the Zaarouria site from the ENAP one. The Zaarouria site which is characterized by sandy texture and high values of pH, ZnF and T. Lime. By opposition, the ENAP site is marked by very high concentrations of Zn, Sulf. F and Sulf. V.

The PCA results also highlighted the relationship between the soil's physicochemical parameters and the distribution of sulfates as well as TME in the soil and the edible tissues of *C. aspersum*. The findings indicated that the soil Zn is completely independent of T.Lime, Sal. and EC. Conversely, these latter physicochemical parameters could promote the accumulation of ZnF. Additionally, ZnV appeared negatively correlated with the soil pH.

Likewise, soil sulfates appeared positively correlated with OM, CEC, P, C and the silty-clay textures. In contrast, T. Lime, EC and Sal. could inhibit the absorption of sulfates in the foot and viscera (Sulf. F and Sulf. V) in *C. aspersum* species. Copper exhibited a tendency to be present in higher concentrations in soils rich in N and sandy texture compared to silty-clay ones. Finally, the PCA biplot showed that copper absorption by snails' tissues (CuF and CuV) could be more pronounced in silty-clay soils rich in OM and C than in sandy soils.

3.2. Hierarchical cluster analysis (HCA)

The objective of HCA is to evaluate the similarities between soil samples according to the bioavailability of various TME in the soil as well as the physicochemical parameters analyzed across the three geographic sites, the cluster analysis dendrogram presented a plausible classification (in concordance with the PCA results). It clearly separated the Mechrouha site (cluster 1, high sulfates level according the PCA biplot), characterized by silty-clay soil, from the two others. The cluster 2 included two relatively homogeneous sites: Zaarouria and ENAP, mainly characterized by high N and Cu levels (according to the PCA biplot). Thus, the hierarchical analysis revealed heterogeneity levels among the soil samples, reflecting the north-south geographic transect. The observed heterogeneity could support the TME and sulfates distributions both in soil and in snails (**Figure 3**, see PCA biplot **Figure 2**).




Figure 3. Hierarchical classification of sampling sites of *C. aspersum* (Zaarouria, Mechrouha and ENAP) based on soil physicochemical parameters and TME.

3.3. TME and sulfates assessment in the snail's foot and viscera

The mean of the CuF concentrations (in dry weight) ranged from $46.91 \pm 2.83 \mu\text{g/g}$ to $78.58 \pm 5.06 \mu\text{g/g}$ at Zaarouria and Mechrouha sites; respectively (**Figure 4**); with a minimum of $36.44 \mu\text{g/g}$ at Zaarouria site and a maximum of $140.33 \mu\text{g/g}$ at ENAP site. The mean of the CuV concentrations ranged from $49.52 \pm 2.06 \mu\text{g/g}$ to $99.55 \pm 8.04 \mu\text{g/g}$ at Zaarouria and Mechrouha sites, respectively; with a minimum of $37.45 \mu\text{g/g}$ at Zaarouria site and a maximum of $146.66 \mu\text{g/g}$ at Mechrouha site. The nonparametric Kruskal-Wallis rank sum test revealed significant differences between sites only in viscera tissues (CuV, $p < 0.05$).




Figure 4. Variations in copper (Cu) concentrations measured in foot and viscera tissues of *C. aspersum*. Data are expressed as mean $\pm$ standard error (SE); the lowercase letters indicate significant differences (Dunn's test) among sites ($n = 5$).

Besides the Cu assessment, the mean of the ZnF concentrations ranged from $68.85 \pm 1.21 \mu\text{g/g}$ to $73.51 \pm 1.81 \mu\text{g/g}$ at ENAP and Zaarouria sites, respectively (**Figure 5**); with a minimum of $61.60 \mu\text{g/g}$ at Mechrouha site and a maximum of $84.10 \mu\text{g/g}$ at Zaarouria site. In addition, the mean of ZnV concentrations ranged from $161.61 \pm 8.38 \mu\text{g/g}$ and $327.11 \pm 32 \mu\text{g/g}$ at Zaarouria and Mechrouha sites, respectively; with a minimum of $123.68 \mu\text{g/g}$ at Zaarouria site and a maximum of $472.41 \mu\text{g/g}$ at Mechrouha site. The Kruskal-Wallis rank sum test also revealed significant differences among sites only in viscera tissues (ZnV, $p < 0.05$).

In this study we have also performed the nonparametric Mann-Whitney rank sum test to compare, within each site, TME and sulfates concentrations between viscera and feet tissues. The statistical results highlighted significant

differences in sulfates concentrations between snails' viscera and feet tissues only for ENAP site (**Figure 6**). Regarding to the TME data, the Mann-Whitney test also revealed, in each site, no significant differences in Cu concentrations between viscera and feet tissues. In contrast, significant differences in Zn concentrations were highlighted among viscera and feet tissues within each site. All these results will be explained in the discussion section.




Figure 5. Variations in zinc (Zn) concentrations measured in foot and viscera tissues of *C. aspersum*. Data are expressed as mean $\pm$ standard error (SE); the lowercase letters indicate significant differences (Dunn's test) among sites ($n = 5$).




Figure 6. Variations in Sulfates (Sulf) concentrations measured in foot and viscera tissues of *C. aspersum*. Data are expressed as mean $\pm$ standard error (SE); the lowercase letters indicate significant differences (Dunn's test) among sites ($n = 5$).

3.4. Correlation analysis

The results of the Spearman's correlation analysis showed that 68% of the correlations, calculated between the twenty-one studied variables, were not statistically significant. All correlation coefficients are shown in **Figure 7** as a corplot; the latter provided a visual exploratory tool on the correlation matrix relating to TME, sulfates and physicochemical parameters concentrations (measured in snails' tissues and soil samples) and it determined the statistical significance of the correlations.

The corplot highlighted that only 67 correlation coefficients were statistically significant ($p < 0.05$). As an example: the results clearly indicated highly significant positive correlations ($r = 1$) between several variables, such as Sulf-CuV, ZnV-CuV, Zn-Sulf.V, Sal-EC, P-N and CuF-ZnV. On the other hand, the corplot also highlighted highly significant negative correlations ($r = -1$) between many variables such as pH-CuF, Sulf.V-ZnF, EC-Sulf.F, C-pH, Zn-ZnF and Sulf.-Sand.




Figure 7. Spearman's correlation carried out on TME, sulfates and physicochemical parameters measured in both snails' tissues and soil samples in all sites.

4. Discussion

4.1. Impacts of soil physicochemical parameters on the bioavailability of TME and sulfates

Physicochemical parameters are among the main factors influencing the distribution and mobility of metals in soils. Several studies including those by van Gestel (2008), Laurent *et al.* (2020) and Kicińska *et al.* (2022) have demonstrated that an increase in pH reduces the concentrations of certain metals such as zinc, cadmium and lead. Włostowski *et al.* (2016) highlighted that under acidic soil conditions the concentration of cadmium tends to increase, which promotes its absorption by *Helix pomatia* and its accumulation in the intestinal gland. This accumulation results from various mechanisms, including ingestion of plants that have absorbed cadmium, direct ingestion of soil particles, and cutaneous diffusion through the snail's foot. According to Wei *et al.* (2021) acidic soils contribute significantly to the increased availability of Cu, Ni and Zn while simultaneously reducing the accumulation of Cd, Cu and Zn. These observations emphasize the influence of soil pH on the mobility and bioavailability of heavy metals as well as their uptake by terrestrial organisms. Our results are consistent with these findings showing that in alkaline soils with elevated pH levels the concentrations of ZnV, CuV and CuF tend to decrease. Heavy metals are generally insoluble and less available for plant uptake in alkaline soils, which could be explained by the fact that at high pH levels metals may form insoluble complexes or precipitate with other soil elements thereby reducing their mobility and bioavailability in soil (Yu *et al.* 2016; Sintorini *et al.* 2021). Shahid *et al.* (2017) demonstrated that soil pH plays a critical role in modulating the chemical forms of heavy metals by affecting various physicochemical processes, including adsorption/desorption, precipitation/dissolution and the formation of complexes with soil components. Conversely, the study by Li *et al.* (2014a) specifically mentions that soil acidity not only promotes the bioavailability of metals to terrestrial species such as *Eisenia fetida* but also increases their accumulation in soils.

The PCA analysis from this study revealed that the soil at Mechrouha is richer in sulfates compared to other sites. It is well known that this site experienced several significant fires two years ago, which likely led to substantial changes in soil composition including pH, organic matter content, and nutrient availability. Among the constituents of ash is sulfuric acid, which can increase soil acidity as evidenced by our findings for this site. Clemente *et al.* (2003) emphasized that soil acidification resulting from the oxidation of sulfides is a key factor in increasing the bioavailability of metals as demonstrated by the concentrations of Cu and Zn observed in the tissues of *C. aspersum* (CuF, CuV and ZnV). Furthermore, soil texture as discussed by Angelaki *et al.* (2022); Sungur *et al.* (2023) influences the distribution and availability of metals. Clayey soils exhibit a high adsorption capacity for various metals except for lead, which is in agreement with our results. We observed that CuF, CuV and ZnV tend to accumulate in silty clay soils rather than in sandy soils. The presence of Cu and Zn in the tissues of *C. aspersum* can be attributed to the organic richness and acidity of silty clay soils. These conditions promote increased metal desorption enhancing their solubility and availability not only to plants but also to other soil organisms such as snails (He *et al.* 2020; AbdelRahman *et al.* 2022; Zhu *et al.* 2023). Our results suggest that an increase in certain physicochemical parameters such as T. Lime, EC and salinity reduces the presence of zinc and copper in the soil. This observation contradicts the findings of Benahmed *et al.* (2016) and Bartkowiak *et al.* (2020) who reported an increase in zinc concentration in response to salinity, EC and T. Lime. A possible explanation for our findings could be competition between sodium cations (Na^+) in saline soils and other cations such as calcium (Ca^{2+}) in calcareous soils with zinc (Zn^{2+}) and copper (Cu^{2+}) ions for sorption sites on soil particles. This competition could reduce the amount of zinc retained by the soil (Acosta *et al.* 2011) increasing its concentration in the soil solution, making it more accessible for uptake by plants and other terrestrial organisms. This mechanism could also explain the results for ZnF, whose accumulation is promoted by these same three parameters (T. Lime, Sal. and EC). This observation aligns with studies by Zhang *et al.* (2020), Park *et al.* (2021) and which confirmed that salinity affects the mobility of metals particularly cadmium by increasing its mobility. Kadkhodaie (2012) also observed a significant increase in cadmium in different parts of sunflower and Sudan grass (roots, stems and leaves) in response to increased salinity. In addition, our results also reveal that the parameters T. Lime, Sal. and EC could influence the mobility and availability of sulfates inhibiting their uptake by the edible tissues of *C. aspersum* (Sulf.F and Sulf.V). This phenomenon could be explained by the fact that in calcium and sodium rich soils sulfates can form partially insoluble ionic complexes such as gypsum. These less soluble complexes tend to remain bound to the soil matrix. This hypothesis is supported by the study of Zunino and Scrivener (2020) which suggests that sulfates tend to form fewer mobile complexes in the presence of calcium. The study by Salao *et al.* (2017) confirmed that industrial emissions play a

significant role in the concentration of nitrogen in the environment, which corroborates our results highlighting the high levels of nitrogen in the Zaarouria site (urban and industrial). This pollution could be due to the release of nitrogen compounds such as ammonia and nitrogen oxides during industrial processes. These emissions can be deposited in the soil by precipitation, thus increasing the concentration of nitrogen in the soil (Stevenson 1965). The application of nitrogen fertilizers can alter the mobility and bioavailability of metals in the soil as confirmed by Li *et al.* (2019) whose study suggests that increasing nitrogen inputs to the soil could reduce the bioavailability of certain metals thus limiting their uptake by plants and microorganisms. These findings are consistent with our results which show that the presence of nitrogen promotes Cu accumulation in the soil while reducing its transfer to the tissues of *C. aspersum*.

4.2. TME (copper and zinc) in the foot and viscera of snails

The presence of TME in the soil can have a significant impact on organisms such as terrestrial gastropods which are considered bioindicators of pollution. Due to their sedentary lifestyle, these organisms have the ability to accumulate various pollutants including TME, thereby reflecting the contamination status of their environment (Mukhtorova *et al.* 2023; Rašković *et al.* 2023). Among the metals detected in the soils of the studied areas are zinc and copper which were also found in the edible tissues of the snails. Our analyses of the TME indicate that Cu and Zn were primarily accumulated in the foot and viscera of snails, revealing significant spatial variation among the three sampling sites for Cu in these parts, as well as for Zn in the viscera. This accumulation was particularly notable at the Mechrouha and ENAP sites compared to the Zaarouria site. The spatial variation of Cu can be explained by the different concentrations of this metal in the soils of three sites which may be influenced by the physicochemical parameters of the soil, leading to differences in absorption by the snail's foot and viscera. The spatial variation of Zn in the viscera only can be attributed to varying levels of Zn in the food sources of *C. aspersum* as well as the specific capacities of the viscera to absorb and accumulate this metal (Mitev *et al.* 2023) the absence of spatial variation of Zn in the snails' feet can be explained by similar or very close concentrations of Zn in the soils of the three sites. Our results also revealed a notable fluctuation in the distribution of Cu and Zn between the tissues of *C. aspersum*. The viscera exhibit a higher accumulation of metals compared to the foot which supports the findings of Gomot-de Vaufléury and Pihan (2002), Hoang *et al.* (2008) and Benhamdoun *et al.* (2024) who confirmed through their studies on metal bioaccumulation in land snails that the viscera accumulate significantly more metals than the foot. For example, Hoang *et al.* (2008) reported that copper was predominantly accumulated in the viscera, accounting for approximately 60% of the total burden, while the shell contained less than 4%. The foot represented about 40% of the total copper accumulation. This observation can be explained by the particularly active metabolism of the

viscera which is responsible for the accumulation, processing and storage of nutrients, leading to higher metal accumulation. In contrast, the foot is primarily composed of muscles dedicated to locomotion. However, this does not exclude the accumulation of metals in the foot or other parts of the snail, as some metals tend to concentrate in the foot or the shell (Aleksander-Kwaterczak and Gołas-Siarzewska 2015; Ćirić *et al.* 2018; Huang *et al.* 2018). In addition, the mode of exposure to these contaminants (whether by dermal contact or ingestion) plays a significant role in the distribution of metals within various parts of the snail body. This observation aligns with the study by (De-Vaufleury and Bispo, 2000) who supported this theory through their experiments. Sebban *et al.* (2022) revealed that certain metals exhibit a preference for accumulation in specific tissues, with Zn tending to accumulate more heavily in the viscera, while Cu is found in abundance in the foot of snails. Our results regarding Zn are consistent with this study but they are different for Cu. Indeed, we observed that Cu accumulates almost equally in both the viscera and the foot of the snails with a slight predominance in the viscera in all three study areas. This can be explained by a significant presence of Cu in the snails' food sources rather than in the soils in which they live.

4.3. Sulfates concentration in the feet and viscera of snails

Although sulfur is an essential macronutrient (Maruyama-Nakashita and Ohkama-Ohtsu 2017; Li *et al.* 2020) high concentrations can cause issues in soil, water and microorganisms (Geurts *et al.* 2009; Björnerås *et al.* 2019). There are several potential sources of sulfates in the soil as we hypothesized in our study. We detected sulfates at all three distinct sites, the presence of sulfates at the Zaarouria site could be attributed to vineyards where sulfur-based treatments are commonly used (Hermes *et al.* 2022). Additionally, this presence could also be explained by household waste as this site is located in an urban area. The study by Malik *et al.* (2020) showed that organic matter promotes the presence of sulfur in the soil. Therefore, it is reasonable to conclude that household waste which often contains organic matter and nutrients could also be a source of sulfur. For the Mechrouha site, we hypothesize that the presence of sulfates is a result of the wildfires that occurred during the summer of 2022 both at this site and in other areas of Souk Ahras province. This is consistent with the study by Hermes *et al.* (2021) which shows that wildfire can introduce sulfur into the soil especially in agricultural areas by altering soil chemistry and facilitating the release of sulfur into the environment. Regarding the ENAP site (industrial activities), the presence of sulfates could be attributed to sulfur in raw materials used in paints particularly pigments based on barium sulfates. This site is the only one showing a significant difference in sulfate accumulation between the foot and the viscera of the snails (p -value = 0.0079), with accumulation being much higher in the foot (533.18 mg/g in dry weight). This contradicts the study by Lobo-da-Cunha (2019) which indicated that the digestive gland of mollusks is the primary site for nutrient absorption. Our result could be explained

by a much higher concentration of sulfates in the soil compared to what is found in the snails' food sources. Additionally, this difference may be due to a high concentration of calcium (Ca^{2+}) and magnesium (Mg^{2+}) ions which compete with sulfates for absorption, resulting in different accumulation levels in the foot and viscera of the snails. In contrast, the accumulation of sulfates in both the foot and viscera of snails across the three study sites (inter-site) showed no significant variation. This could be due to consistent sulfate concentrations across the three study sites and a uniform absorption mechanism by the snails' feet and viscera.

5. Conclusion

The results of this research highlight the potential of utilizing metals concentrations as environmental indicators, while emphasizing the need to consider spatial variations as well as the physicochemical parameters of the soil that influence these indicators at our sites, which in turn affects the level of metal absorption by terrestrial gastropods. We also measured sulfate levels in the foot and viscera of *C. aspersum* based on preliminary soil analysis by X-ray fluorescence spectrometry which revealed the presence of sulfur both in the soil and our target snail species. The results demonstrate that physicochemical parameters significantly influence the availability of TME in the soil leading to differences in their accumulation within edible tissues. Specifically, it was observed that copper and zinc accumulate much more in the viscera than in the foot of *C. aspersum* while sulfate shows the opposite pattern with higher concentrations in the foot. These results confirm that *C. aspersum* is an excellent bioindicator of soil pollution. Accumulation in the different tissues of edible snails including our study species *C. aspersum* could have harmful effects on human health through the food chain. This study presents certain limitations, including the lack of seasonal monitoring, limited geographic scope, and a restricted range of analyzed contaminants. Future research should broaden the study area, incorporate a wider array of pollutants and bioindicator species, and include temporal monitoring to better understand bioaccumulation dynamics. However, such proposed work may also face constraints such as logistical challenges, environmental variability, and limited accessibility to target species.

Acknowledgements

The authors would like to express their sincere gratitude to the Research Laboratory RME – Resources, Materials and Ecosystems, University of Carthage, Faculty of Sciences, Bizerte, Tunisia, for their essential support in providing access to the necessary equipment. Special thanks are also extended to all the laboratory staff for their invaluable assistance throughout this work.

References

- AbdelRahman, M. A. E., Metwaly, M. M., Afifi, A. A., D'Antonio, P. and Scopa, A. (2022). Assessment of soil fertility status under soil degradation rate using geomatics in West Nile Delta, *Land*, **11**, 1256, <https://doi.org/10.3390/land11081256>.
- Acosta, J. A., Jansen, B., Kalbitz, K., Faz, A. and Martínez-Martínez, S. (2011). Salinity increases mobility of heavy metals in soils, *Chemosphere*, **85**, 1318–1324, <https://doi.org/10.1016/j.chemosphere.2011.07.046>.
- Ajayi, A. A. and Oyewole, B. O. (2023). Giant African land snails (*Achatina achatina* and *Archachatina marginata*) as bioindicators of heavy metal pollution, *African Journal of Environmental Science and Technology*, **17**, 80–88, <https://doi.org/10.5897/AJEST2022.3131>.
- Al-Alam, J., Millet, M., Khoury, D., Rodrigues, A., Akoury, E., Tokajian, S. and Wazne, M. (2024). Biomonitoring of PAHs and PCBs in industrial, suburban, and rural areas using snails as sentinel organisms, *Environmental Science and Pollution Research*, **31**, 4970–4984, <https://doi.org/10.1007/s11356-023-31493-6>.
- Al-Alam, J., Millet, M., Khoury, D., Rodrigues, A., Harb, M., Akoury, E., Tokajian, S. and Wazne, M. (2022). Snails as temporal biomonitors of the occurrence and distribution of pesticides in an apple orchard, *Atmosphere*, **13**, 1185, <https://doi.org/10.3390/atmos13081185>.
- Aleksander-Kwaterczak, U. and Gołas-Siarzewska, M. (2015). Comparative analysis of *Helix pomatia* L. shells found in soils with varying degrees of contamination (southern Poland), *Geology, Geophysics and Environment*, **41**, 299, <https://doi.org/10.7494/geol.2015.41.4.299>.
- Alsherif, E. A., Al-Shaikh, T. M. and Abdelgawad, H. (2022). Heavy metal effects on biodiversity and stress responses of plants inhabiting contaminated soil in Khulais, *Saudi Arabia, Biology*, **11**, 164, <https://doi.org/10.3390/biology11020164>.
- Amlan, P. G., Chouti, W. K., Dedjiho, C. A., Fangnon, K. R. and Chitou, N. E. (2023). Évaluation de la pollution chimique des eaux par les éléments traces métalliques (ETM): cas de la rivière Mekrou (nord-ouest Bénin), *Journal of Applied Bioscience*, **192**, 20331–20346, <https://doi.org/10.35759/JABs.192.4>.
- Angelaki, A., Dionysidis, A., Sihag, P. and Golia, E. E. (2022). Assessment of contamination management caused by copper and zinc cations leaching and their impact on the hydraulic properties of a sandy and a loamy clay soil, *Land*, **11**, 290, <https://doi.org/10.3390/land11020290>.
- Baize, D. and Jabiol, B. (1995). Guide pour la description des sols, INRA, Paris, *Techniques et Pratiques*.
- Bankaji, I., Kouki, R., Dridi, N., Ferreira, R., Hidouri, S., Duarte, B., Sleimi, N. and Caçador, I. (2023). Comparison of digestion methods using atomic absorption spectrometry for the determination of metal levels in plants, *Separations*, **10**, 40, <https://doi.org/10.3390/separations10010040>.
- Baroudi, F. (2020). Étude comparative de trois biomoniteurs (conifère, escargot et miel) pour évaluer la variabilité spatio-temporelle de polluants organiques dans l'atmosphère au Liban, PhD thesis, Université de Strasbourg, *Chimie analytique*, NNT : 2020STRAF023.
- Bartkowiak, A., Dąbkowska-Naskręt, H., Jaworska, H. and Rydlewska, M. (2020). Effect of salinity on the mobility of trace metals in soils near a soda chemical factory, *Journal of Elementology*, **25**, 501–512, <https://doi.org/10.5601/jelem.2019.24.2.1875>.
- Benhamdoun, A., Achtaq, H. and Dahbi, A. (2024). Bioaccumulation of trace metals in edible terrestrial snails, *Theba pisana* and *Otala* spp., in a dumpsite area in Morocco and assessment of human health risks for consumers, *Environmental Science and Pollution Research*, **31**, 42810–42826, <https://doi.org/10.1007/s11356-024-33945-z>.
- Bennour, A. and Soltani, N. (2020). Assessment of soil quality in Annaba area (northeast Algeria) using the earthworm

- Lumbricus terrestris: bioindicative stress responses and heavy metal contamination, *Fresenius Environmental Bulletin*, **29**, 9635–9643.
- Bici, M., Sahiti, H., Zogaj, M., Halili, J. and Bislimi, K. (2024). Data on heavy metals (Pb, Ni, Zn) in soil and biota (common nettle and Roman snail) around the power plant TC Kosova A in Obiliq (Kosovo), *Jordan Journal of Biological Sciences*, **17**, 1–7, <https://doi.org/10.54319/jjbs/170101>.
- Björnerås, C., Škerlep, M., Floudas, D., Persson, P. and Kritzbeg, E. S. (2019). High sulfate concentration enhances iron mobilization from organic soil to water, *Biogeochemistry*, **144**, 245–259, <https://doi.org/10.1007/s10533-019-00581-6>.
- Bonnefont, J. C., Bonneau, M. and Souchier, B. (1980). Constituants et propriétés du sol, *Revue Géographique de l'Est*, **20**, 263–264, https://www.persee.fr/doc/rgeest_0035-3213_1980_num_20_3_1477_t1_0263_0000_2.
- Boudy, P. (1955). Économie forestière nord-africaine, vol. 4, Description forestière de l'Algérie et de la Tunisie, *Larose*, Paris.
- Bouzahouane, H., Kouki, R., Amri, S., Barour, C., Sleimi, N. and Ouali, K. (2024). Investigating seasonal metal impact on Stramonita haemastoma gastropod along the Algerian East Coast: Understanding through various pollution indicators, *Marine Pollution Bulletin*, **199**, 116006, <https://doi.org/10.1016/j.marpolbul.2023.116006>.
- Chen, F., Wang, Q., Meng, F., Chen, M. and Wang, B. (2023). Effects of long-term zinc smelting activities on the distribution and health risk of heavy metals in agricultural soils of Guizhou province, China, *Environmental Geochemistry and Health*, **45**, 5639–5654, <https://doi.org/10.1007/s10653-020-00716-x>.
- Cheriti, O. (2022). Utilisation des gastéropodes pulmonés terrestres comme traceurs de la pollution métallique des sols dans le bassin du Kebir Rhumel: de l'inventaire à la bioindication, PhD thesis, Université Frères Mentouri Constantine 1, Département de Biologie et Écologie Végétale.
- Ćirić, J., Cerić, O., Marković, R., Janjić, J., Spirić, D., Popović, M., Pećanac, B., Baltić, B. and Baltić, M. (2018). Seasonal distributions of heavy metal concentrations in different snail (*Helix pomatia*) tissues from an urban environment in Serbia, *Environmental Science and Pollution Research*, **25**, 33415–33422, <https://doi.org/10.1007/s11356-018-3295-1>.
- Clemente, R., Walker, D. J., Roig, A. and Pilar, M. P. (2003). Heavy metal bioavailability in a soil affected by mineral sulphides contamination following the mine spillage at Aznalcóllar (Spain), *Biodegradation*, **14**, 199–205.
- Core Team R. (2022). R: A language and environment for statistical computing, Available at: <https://www.R-project.org/>.
- De-Vaufleury, G. A. and Bispo, A. (2000). Methods for toxicity assessment of contaminated soil by oral or dermal uptake in land snails. 1. Sublethal effects on growth, *Environmental Science & Technology*, **34**, 1865–1870, <https://doi.org/10.1021/es9907212>.
- Dinno, A. (2017). Dunn's Test of Multiple Comparisons Using Rank Sums, R package version 1.5.1.
- Dossou, D., Bokossa, H. K. J., Adanlokonon, E. A. S., Zounon, Y., Adanlokonon, S. M. I., Johnson, R. C., Fiogbe, E. D. and Edorth, P. (2022). Evaluation of metal contamination of sediments and crabs (*Callinectes amnicola*) from Lake Ahémé in southern Benin, *International Journal of Biological and Chemical Sciences*, **16**, 2424–2435.
- Drouineau, G. (1942). Dosage rapide du calcaire actif des sols. Nouvelles données sur la répartition et la nature des fractions calcaires, *Annales Agronomiques*, **2**, 441–450.
- Du, C., Wu, J., Bashir, M. H., Shaukat, M. and Ali, S. (2019). Heavy metals transported through a multi-trophic food chain influence the energy metabolism and immune responses of *Cryptolaemus montrouzieri*, *Ecotoxicology*, **28**, 422–428.
- Dupont, J. (2023). Le souffle polymétallique de la fonderie Horne: caractérisation des dépôts atmosphériques à l'aide d'approches passives et de biosurveillance, *PhD thesis*, Université du Québec à Montréal, Montréal.
- El Mageed, Y. S. M. A., Ghobashy, A. E. F. A., Al-Thomali, A. W., Soliman, M. F., Mohammadein, A. and El-Shenawy, N. S. (2023). Impact of heavy metals contamination on biology, biochemical, and histology of *Eobania vermiculata* and *Monacha obstructa*, *Toxicology and Environmental Health Sciences*, **15**, 19–30, <https://doi.org/10.1007/s13530-022-00233-8>.
- Enkatraman, M., Surendran, R., Senduru Srinivasulu., Vijayakumar K. (2025) "Water quality prediction and classification using Attention based Deep Differential RecurFlowNet with Logistic Giant Armadillo Optimization", *Global NEST Journal*. <https://doi.org/10.30955/gnj.0679>.
- Gboko, Y. D. A., Natchia, A., Keumeane, K. N. and Soro, N. (2022). Suivi de la qualité physico-chimique des eaux du Sassandra à la station hydrométrique de Gaoulou, Sud-ouest de la Côte d'Ivoire, *African Sciences*, **21**, 127–141.
- Geurts, J. J. M., Sarneel, J. M., Willers, B. J. C., Roelofs, J. G. M., Verhoeven, J. T. A. and Lamers, L. P. M. (2009). Interacting effects of sulphate pollution, sulphide toxicity, and eutrophication on vegetation development in fens: A mesocosm experiment, *Environmental Pollution*, **157**, 2072–2081, <https://doi.org/10.1016/j.envpol.2009.02.024>.
- Gomot-de Vaufleury, A. and Pihan, F. (2002). Methods for toxicity assessment of contaminated soil by oral or dermal uptake in land snails: metal bioavailability and bioaccumulation, *Environmental Toxicology and Chemistry*, **21**, 820–827, <https://doi.org/10.1002/etc.5620210419>.
- Guessasma, Z., Khaldi, F., Grara, N., Agouni, M., Sleimi, N. and Benslama, M. (2020). Assessment of heavy metal soil contamination in some northeastern Algerian biotopes by using the terrestrial snail, *Helix aspersa*, *Studia Univ. Vasile Goldis Arad, Seria Stiintele Vietii (Life Sciences Series)*, **30**, 55–63, <https://www.univ-soukahras.dz/wpuploads/eprints/2021-992-681d3.pdf>.
- Harrell, J. F. (2023). Hmisc: Harrell Miscellaneous, R package version 5.1-1.
- He, G., Zhang, Z., Wu, X., Cui, M., Zhang, J. and Huang, X. (2020). Adsorption of heavy metals on soil collected from lixisol of typical karst areas in the presence of CaCO₃ and soil clay and their competition behavior, *Sustainability (Switzerland)*, **12**, 7315, <https://doi.org/10.3390/SU12187315>.
- Hermes, A. L., Dawson, T. E. and Hinckley, E. L. S. (2022). Sulfur isotopes reveal agricultural changes to the modern sulfur cycle, *Environmental Research Letters*, **17**, 054032, <https://doi.org/10.1088/1748-9326/ac6683>.
- Hermes, A. L., Ebel, B. A., Murphy, S. F. and Hinckley, E. L. S. (2021). Fates and fingerprints of sulfur and carbon following wildfire in economically important croplands of California, US, *Science of the Total Environment*, **750**, 142179, <https://doi.org/10.1016/j.scitotenv>.

- Hidouri, N., Moussaoui, Z., Sleimi, N., Hamed, Y. and Hamzaoui-Azzaza, F. (2024). Soil contamination by trace elements and radioelements and related environmental risks in agricultural soils of the M'Dhilla Basin (southwestern Tunisia). *Environmental Monitoring and Assessment*, **196**(11), 1024.
- Hoang, T. C., Rogevich, E. C., Rand, G. M. and Frakes, R. A. (2008). Copper uptake and depuration by juvenile and adult Florida apple snails (*Pomacea paludosa*). *Ecotoxicology*, **17**, 605–615, <https://doi.org/10.1007/s10646-008-0243-8>.
- Huang, F., Peng, L., Zhang, J., Lin, W. and Chen, S. (2018). Cadmium bioaccumulation and antioxidant enzyme activity in hepatopancreas, kidney, and stomach of invasive apple snail *Pomacea canaliculata*. *Environmental Science and Pollution Research*, **25**, 18682–18692, <https://doi.org/10.1007/s11356-018-2092-1>.
- Ibrahim, O. Z., Tankari, D. B. A., Guero, Y., Maissoro, M. I. F., Feidt, C., Sterckeman, T. and Echevarria, G. (2019). Spatial distribution of metallic trace elements in soils of Komabangou gold zone in Niger. *International Journal of Biological and Chemical Sciences*, **13**(1), 557–573.
- Ihaka, R. and Gentleman, R. (1996). R: A language for data analysis and graphics. *Computational and Graphical Statistics*, **5**, 299–314.
- Kadkhodaie, A., Kelich, S. and Baghbani, A. (2012). Effects of salinity levels on heavy metals (Cd, Pb, and Ni) absorption by sunflower and sudangrass plants. *Bulletin of Environment, Pharmacology and Life Sciences*, **1**, 47–53.
- Kassambara, A. (2023a). ggcorrplot: Visualization of a Correlation Matrix using 'ggplot2', R package version 0.1.4.1.
- Kassambara, A. (2023b). ggpubr: 'ggplot2' Based Publication Ready Plots, R package version.
- Kassambara, A. and Mundt, F. (2020). factoextra: Extract and Visualize the Results of Multivariate Data Analyses, R package version 1.0.7.
- Kicińska, A., Pomykała, R. and Izquierdo-Díaz, M. (2022). Changes in soil pH and mobility of heavy metals in contaminated soils. *European Journal of Soil Science*, **73**, e13203, <https://doi.org/10.1111/ejss.13203>.
- Kim, S. H., Bae, S. and Hwang, Y. S. (2023). Comparative bioaccumulation, translocation, and phytotoxicity of metal oxide nanoparticles and metal ions in soil-crop system. *Science of the Total Environment*, **856**, 158938, <https://doi.org/10.1016/j.scitotenv.2022.158938>.
- Kjeldahl, J. (1883). Neue Methode zur Bestimmung des Stickstoffs in organischen Körpern. *Fresenius' Zeitschrift für Analytische Chemie*, **22**, 366–382, <https://doi.org/10.1007/BF01338151>.
- Lag, J., Hadas, A., Fairbridge, R. W., Nóvoa Muñoz, J. C., Pombal, X. P., Cortizas, A. M., Almendros, G., Chesworth, W., Arbestain, M. C., Macías, F., Martínez, C. A., Chesworth, W., Jaynes, W. F. and Winterkorn, H. F. (2008). Hygroscopicity hygroscopy constant. *Encyclopedia of Soil Science*, Springer.
- Larba, R., Zeraib, A. and Soltani, N. (2023). Assessment of soil metallic contamination in several sites from Northeast Algeria by use of terrestrial gastropod: *Cornu aspersum* (O.F. Müller, 1774) (Helicidae). *Biodiversity Journal*, **14**, 373–383, <https://doi.org/10.31396/biodiv.jour.2023.14.2.373.383>.
- Laurent, C., Bravin, M. N., Crouzet, O., Pelosi, C., Tillard, E., Lecomte, P. and Lamy, I. (2020). Increased soil pH and dissolved organic matter after a decade of organic fertilizer application mitigates copper and zinc availability despite contamination. *Science of the Total Environment*, **709**, 135927, <https://doi.org/10.1016/j.scitotenv.2019.135927>.
- Lê, S., Josse, J., Rennes, A. and Husson, F. (2008). FactoMineR: An R Package for Multivariate Analysis. *Journal of Statistical Software*, **25**, 1–18, <http://www.jstatsoft.org/>.
- Li, L., Wu, H., Van Gestel, C. A. M., Peijnenburg, W. J. G. M. and Allen, H. E. (2014). Soil acidification increases metal extractability and bioavailability in old orchard soils of Northeast Jiaodong Peninsula in China. *Environmental Pollution*, **188**, 144–152, <https://doi.org/10.1016/j.envpol.2014.02.003>.
- Li, Q., Gao, Y. and Yang, A. (2020). Sulfur homeostasis in plants. *International Journal of Molecular Sciences*, **21**, 1–16, <https://doi.org/10.3390/ijms21238926>.
- Li, R., Tan, W., Wang, G., Zhao, X., Dang, Q., Yu, H. and Xi, B. (2019). Nitrogen addition promotes the transformation of heavy metal speciation from bioavailable to organic bound by increasing the turnover time of organic matter: An analysis on soil aggregate level. *Environmental Pollution*, **255**, 113170, <https://doi.org/10.1016/j.envpol.2019.113170>.
- Li, X., Liang, H., Zeng, Y., Zheng, X., Ren, Z. and Mai, B. (2023). Trophic transfer of heavy metals in a wetland food web from an abandoned e-waste recycling site in South China. *Science of the Total Environment*, **890**, 164327.
- Lobo-da-Cunha, A. (2019). Structure and function of the digestive system in molluscs. *Cell and Tissue Research*, **377**, 475–503, <https://doi.org/10.1007/s00441-019-03085-9>.
- Louzon, M., de Vaufléury, A. and Capelli, N. (2023). Ecogenotoxicity assessment with land snails: A mini-review. *Mutation Research - Reviews in Mutation Research*, **792**, 108472, <https://doi.org/10.1016/j.mrrev.2023.108472>.
- Malik, K. M., Khan, K. S., Akhtar, M. S. and Ahmed, Z. I. (2020). Sulfur distribution and availability in alkaline subtropical soils affected by organic amendments. *Journal of Soil Science and Plant Nutrition*, **20**, 2253–2266, <https://doi.org/10.1007/s42729-020-00292-0>.
- Maruthai, S., Rajendran, S., Selvanarayanan, R. and Gowri, S. (2025). “Wastewater Recycling Integration with IoT Sensor Vision for Real-time Monitoring and Transforming Polluted Ponds into Clean Ponds using HG-RNN”. *Global NEST Journal*. **27**(4). <https://doi.org/10.30955/gnj.06758>.
- Maruyama-Nakashita, A. and Ohkama-Ohtsu, N. (2017). Sulfur assimilation and glutathione metabolism in plants, in *Glutathione in Plant Growth, Development, and Stress Tolerance*, Springer - International Publisher, pp 287–308, https://doi.org/10.1007/978-3-319-66682-2_13.
- Mitev, T., Dedov, I. K. and Hristovski, S. (2023). Contents of Zn, Cu, Mn and Cd in soil, plants and Turkish snails *Helix lucorum* Linnaeus, 1758 in Skopje, Republic of North Macedonia. *Acta Zoologica Bulgarica*, **75**, 2.
- Mohamed, B., Abdelkader, D. and Benchaben, H. (2016). Mobilité du plomb et du zinc issus de retombées atmosphériques dans le sol : Cas de la zone industrielle de Tiaret, Algérie. *European Scientific Journal*, *ESJ*, **12**, 131, <https://doi.org/10.19044/esj.2016.v12n18p131>.
- Monchanin, C., Devaud, J. M., Barron, A., Lihoreau, M. and Barron, A. B. (2021). Current permissible levels of metal pollutants harm terrestrial invertebrates. *Science of the Total Environment*, **779**, 146398, <https://doi.org/10.1016/j.scitotenv.2021.146398>.
- Mukhtorova, D., Hlava, J., Száková, J., Najmanová, J. and Tlustoš, P. (2023). Can mollusks or insects serve as bioindicators of the risk element polluted area? Gastropods (Gastropoda) versus

- leaf beetles (Coleoptera: Chrysomelidae), *Environmental Science and Pollution Research*, **30**, 78707–78717.
- Mwelwa, S., Chungu, D., Tailoka, F., Beesigamukama, D. and Tanga, C. (2023). Biotransfer of heavy metals along the soil-plant-edible insect-human food chain in Africa, *Science of the Total Environment*, **881**, 163150, <https://doi.org/10.1016/j.scitotenv.2023.163150>.
- N'Doua, A. D. Y., Adjessan, M. A. M. and Koffi, K. M. (2024). Détermination du niveau de contamination en métaux lourds (arsenic, cadmium, mercure et plomb) de quatre espèces de poissons consommées par les familles de pêcheurs de Jacqueville, *European Scientific Journal, ESJ*, **20**, 53, <https://doi.org/10.19044/esj.2024.v20n3p53>.
- Ning, Q., Shao, B., Huang, X., He, M., Tian, L. and Lin, Y. (2024). Bioaccumulation, biomagnification, and ecological risk of trace metals in the ecosystem around oilfield production area: A case study in Shengli Oilfield, *Environmental Monitoring and Assessment*, **196**, 87, <https://doi.org/10.1007/s10661-024-102566>.
- Nwagu, C. U., Olujimi, O. O. and Okoro, G. C. (2022). The use of Giant African Snails (*Archachatina marginata* and *Achatina fulica*) as bio-indicators of heavy metal pollution, *IAA Journal of Scientific Research*, **8**, 19–41, ISSN: 2736-7319, Available at: <https://www.researchgate.net/publication/364348221>.
- Ogbeide, O. and Amayanbo, O. (2024). Effect Of Heavy Metal Runoff from Ikpoba River on The Gonad and Antioxidant Properties of African Snail (*Bulinus Africanus*). *African Journal of Health, Safety and Environment*, **5**(2), 10–25.
- Okalebo, J. R., Gathua, K. W. and Woomer, P. L. (2002). Laboratory methods of soil and plant analysis: A working manual, second edition, Sacred Africa, *Nairobi*, pp. 21–26.
- Olsen, S. R. (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate (No. 939), US Department of Agriculture, Available at: <http://books.google.com>.
- Owojori, O. J., Awodiran, M., Ayanda, O. E. and Jegede, O. O. (2022). Toxicity and accumulation of lead and cadmium in the land snail, *Archachatina papyracea*, in a tropical Alfisol from Southwestern Nigeria, *Environmental Science and Pollution Research*, **29**, 44917–44927.
- Park, H. J., Kim, S. U., Jung, K. Y., Lee, S., Choi, Y. D., Owens, V. N., Kumar, S., Yun, S. W. and Hong, C. O. (2021). Cadmium phytoavailability from 1976 through 2016: Changes in soil amended with phosphate fertilizer and compost, *Science of the Total Environment*, **762**, 1–10.
- Periasamy, S., Subramanian, P. and Surendran, R. (2024) “An intelligent air quality monitoring system using quality indicators and Transfer learning based Lightweight recurrent network with skip connection”, *Global NEST*, **26**(5), pp. 1–10.
- Pohlert, T. (2023). PMCMRplus: Calculate Pairwise Multiple Comparisons of Mean Rank Sums Extended, R package version 1.9.7.
- Rašković, B., Poleksić, V., Jarić, I., Skorić, S., Topisirović, G. and Stojnić, B. (2023). Accumulation of metal trace elements in different body parts of terrestrial Roman snail *Helix pomatia* L., 1758 on three polluted sites in Serbia, *Environmental Science and Pollution Research*, **30**(8), 21853–21862, <https://doi.org/10.1007/s11356-022-23697-z>.
- Revelle, W. (2023). psych: Procedures for Psychological, Psychometric, and Personality Research, R package version 2(9).
- Rhoades, J. D. (1982). Soluble salts, Methods of Soil Analysis: Part 2 Chemical and Microbiological Properties, 2nd ed., *American Society of Agronomy, Madison, WI*, pp. 849–867.
- Sahraoui, A. S. and Sahli, L. (2022). Utilisation in situ et ex situ de *Helix aspersa* comme bioindicateur de la contamination des sols par le cadmium et le plomb, Thèse de doctorat, Université Frères Mentouri-Constantine 1.
- Salao, A. B., Diop, C., Livardjani, F. and Djokhane, A. M. (2017). Évaluation des risques de la pollution atmosphérique (CO, NOx, SO2, O3, PM10 et PM2.5) de la ville de Dakar, *Sénégal, Revue Ivoirienne des Sciences et Technologie*, **30**, 1–10, <http://www.revist.ci>.
- Salih, A. H. S., Hama, A. A., Hawrami, K. A. M. and Ditta, A. (2021). The land snail, *Eobania vermiculata*, as a bioindicator of the heavy metal pollution in the urban areas of Sulaimani, Iraq, *Sustainability (Switzerland)*, **13**(24), <https://doi.org/10.3390/su132413719>.
- Sargsyan, A., Hovhannisyan, G., Simonyan, A., Arakelyan, M., Arzumanyan, M. and Aroutiounian, R. (2022). Application of land snail *Helix lucorum* for evaluation of genotoxicity of soil pollution, *Mutation Research - Genetic Toxicology and Environmental Mutagenesis*, **878**, 503500.
- Sebban, H., Ait Belcaid, H., El Alaoui El Fels, A., Bouriqi, A., Pineau, A. and Sedki, A. (2022). Trace element bioaccumulation in the edible milk snail (*Otala lactea*) and Cabrilla (*Otala punctata*) in Marrakech, Morocco, *Applied Ecology and Environmental Research*, **20**(1), 875–892, https://doi.org/10.15666/aer/2001_875892.
- Selvanarayanan, R., Rajendran, S., Algburi, S., Khalaf, O. I. and Hamam, H. (2024a). “Empowering coffee farming using counterfactual recommendation based RNN driven IoT integrated soil quality command system,” *Scientific Reports* **14**, 6269. <https://doi.org/10.1038/s41598-024-56954-x>
- Selvanarayanan, R., Rajendran, S., Pappa, K. C. and Thomas, B. (2024b). “Wastewater Recycling to Enhance Environmental Quality using Fuzzy Embedded with RNNIoT for Sustainable Coffee Farming”. *Global NEST Journal*. Available at: <https://doi.org/10.30955/gnj.06346>.
- Shaffique, S., Kang, S. M., Ashraf, M. A., Umar, A., Khan, M. S., Wajid, M., Al-Ghamdi, A. A. and Lee, I. J. (2024). Research progress on migratory water birds: indicators of heavy metal pollution in Inland Wetland resources of Punjab, Pakistan, *Water (Switzerland)*, **16**(8), 1163, <https://doi.org/10.3390/w16081163>.
- Shahid, M., Dumat, C., Khalid, S., Niazi, N. K. and Antunes, P. M. (2017). Cadmium bioavailability, uptake, toxicity and detoxification in soil-plant system, *Reviews of Environmental Contamination and Toxicology*, **241**, 73–137.
- Sintorini, M. M., Widyatmoko, H., Sinaga, E. and Aliyah, N. (2021). Effect of pH on metal mobility in the soil, *IOP Conference Series: Earth and Environmental Science*, **737**(1), 012071, <https://doi.org/10.1088/1755-1315/737/1/012071>.
- Sleimi, N., Bankaji, I., Kouki, R., Dridi, N., Duarte, B. and Caçador, I. (2022). Assessment of extraction methods of trace metallic elements in plants: approval of a common method, *Sustainability (Switzerland)*, **14**(3), 1428, <https://doi.org/10.3390/su14031428>.
- Soliman, M. M., Hesselberg, T., Mohamed, A. A. and Renault, D. (2022). Trophic transfer of heavy metals along a pollution gradient in a terrestrial agro-industrial food web, *Geoderma*, **413**, 115748.

- Stevenson, F. J. (1965). Origin and distribution of nitrogen in soil, *Soil Nitrogen*, **10**, 1–42.
- Sungur, A., Temel, E., Everest, T., Soylak, M. and Özcan, H. (2023). Effects of soil texture on trace metal concentrations and geochemical fractions in the soil of apple orchards (Çanakkale, NW Turkey), *Archives of Agronomy and Soil Science*, **69**, 1–10.
- Téreygeol, F., Arles, A., Foy, E., Florsch, N. and Llubes, M. (2010). Dosages par fluorescence X portable d'ateliers médiévaux de production des métaux non-ferreux, *ArchéoSciences*, **34**, 243–252, <https://doi.org/10.4000/archeosciences.2802>.
- Thomas, G. W. (1982). Exchangeable cations, *Methods of Soil Analysis: Part 2 Chemical and Microbiological Properties*, **9**, 159–165.
- van Gestel, C. A. M. (2008). Physico-chemical and biological parameters determine metal bioavailability in soils, *Science of the Total Environment*, **406**(3), 385–395, <https://doi.org/10.1016/j.scitotenv.2008.05.050>.
- Walkley, A. and Black, I. A. (1934). An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method, *Soil Science*, **37**, 29–38.
- Wei, X., Zhang, P., Zhan, Q., Hong, L., Bocharnikova, E. and Matichenkov, V. (2021). Regulation of As and Cd accumulation in rice by simultaneous application of lime or gypsum with Si-rich materials, *Environmental Science and Pollution Research*, **28**(6), 7271–7280, <https://doi.org/10.1007/s11356-020-11053-y>.
- Weindorf, D. C. and Chakraborty, S. (2020). Portable X-ray fluorescence spectrometry analysis of soils, *Soil Science Society of America Journal*, **84**(5), 1384–1392, <https://doi.org/10.1002/saj2.20151>.
- Wickham, H. (2016). Data Analysis, in: ggplot2, Springer, Cham, pp. 145–183, https://doi.org/10.1007/978-3-319-24277-4_9.
- Włostowski, T., Kozłowski, P., Łaszkiewicz-Tiszczenko, B. and Oleńska, E. (2016). Cadmium accumulation and pathological alterations in the midgut gland of terrestrial snail *Helix pomatia* L. from a zinc smelter area: role of soil pH, *Bulletin of Environmental Contamination and Toxicology*, **96**, 484–489, <https://doi.org/10.1007/s00128-016-1748-0>.
- Yu, H. Y., Liu, C., Zhu, J., Li, F., Deng, D. M., Wang, Q. and Liu, C. (2016). Cadmium availability in rice paddy fields from a mining area: The effects of soil properties highlighting iron fractions and pH value, *Environmental Pollution*, **209**, 38–45.
- Zhang, S., Ni, X., Arif, M., Yuan, Z., Li, L. and Li, C. (2020). Salinity influences Cd accumulation and distribution characteristics in two contrasting halophytes, *Suaeda glauca* and *Limonium aureum*, *Ecotoxicology and Environmental Safety*, **191**, 110230.
- Zhu, Q., Ji, J., Tang, X., Wang, C. and Sun, H. (2023). Bioavailability assessment of heavy metals and organic pollutants in water and soil using DGT: A review, *Applied Sciences*, **13**(17), 9760.