

Evaluation of Pollution Load Index: A case study from Bollaram Industrial area Vamsee Krishna Kodali¹, Randhi Uma Devi^{2*}, Balachandu Koya³

¹ GITAM Deemed To Be University, Visakhapatnam, Andhra Pradesh State, India - 530045

² GITAM Deemed To Be University, Hyderabad, Telangana State, India - 502329

³ BV Raju Institute of Technology, Narasapur, Medak District, Telangana State, India - 502313


*Corresponding Author: urandhi@gitam.edu

Abstract

This study has been conducted to comprehensively evaluate ecological risk through pollution indices of chemical water quality parameters in the aquatic body of the Bollaram industrial area. For the data, the water quality parameters were analyzed seasonally throughout the year and depicted the significant enrichment of heavy metals. Our results indicated a very high degree of contamination (CDR>24). Further, the Pollution load index also reflected "very high polluted" with PLIX>5 in the study area. Overall, the decreasing trend of Contamination factor (CFR) for the Bollaram lentic aquatic system through various heavy metals was observed as Mn>Fe>Pb>Hg>Zn>Cr>Cd. Overall, the potential ecological risk index (PERI) value for all sampling sites varied from 1629.41 to 2505.37, reflecting very high ecological risks. Fe and Mn predominantly contribute a significant amount to the potential ecological risk in the Bollaram pond. A pollution load index (PLIX) value equal to zero signifies perfection, 1 represents only the occurrence of the baseline level of the various contaminants, whereas a value of more than 1 reflects progressive water contamination by the trace metals. The highest PLIX value was perceived at SP6 (10.23), followed by SP2 (8.79). The lowest PLIX value was obtained at SP4 (5.22), followed by SP7 (5.91). An elevated level of PLIX in the sampling sites reported that the

disposal of municipal waste and sludge from small-scale industries illegally might have caused ecological risk to the water ecosystems.

Key Words: Pollution indices, Ecological risk, Heavy Metal pollution, Contamination



Graphical Abstract

Introduction

Industrialization and urbanization have significantly impacted the ecosystem, causing deterioration and depletion of freshwater resources. Wastewater can come from various sources, including industrial, residential, commercial, agricultural, and storm water runoff in urban areas. Pollutants of concern include organic matter, nutrients, pharmaceuticals, personal care products, polyfluoroalkyl compounds, biocides, heavy metals, dyes, radionuclides, plastics, nanoparticles, and pathogens (Venkataraman *et al.* 2024). According to Adimalla (2020), heavy metal ions are a major source of pollution and should be taken seriously. Studies suggest that live organisms accumulate these compounds gradually over time (Saha *et al.* 2021). They have long-lasting effects and can impact several organisms through biomagnification. Pollution indices (PIS) are widely recognized emphatic tools for comprehensively evaluating and assessing environmental

quality, degree of contamination, and forecasting future environmental sustainability (Kowalska *et al.* 2018). Basically, pollution indices used for the evaluation of heavy metals are classified into two categories: Single (individual metal) and multi or integrated (sum of all studied metals together) pollution index (Cheng *et al.* 2022). Single metal PI includes contamination factor (CNR), ecological risk factor (ERFR), and integrated PIS involves contamination degree (CDR), modified contamination degree (mCDR), pollution load index (PLIX), and potential ecological risk index (PERI) (He *et al.* 2022, Selvanarayanan *et al.* 2024). The research location is the Bollaram industrial zone situated in Hyderabad, India. The pollution issue is exacerbated by the direct release of runoff water from numerous iron, pharmaceutical, and cement businesses into nearby water bodies (Su *et al.* 2023). Consequently, this study was conducted to evaluate pollution trends in the adjacent pond in Bollaram by ecological risk assessments.

Methodology

The study area Bollaram pond is also known as IDA Bollaram area, present in the Hyderabad metropolitan region of India. Many industries are located here, like small and medium-scale iron industries, cement industries, pharma companies, etc. The gathered water samples from study area were examined for their heavy metal composition, including chromium (Cr), manganese (Mn), lead (Pb), iron (Fe), zinc (Zn), cadmium (Cd), and mercury (Hg) via the ICP-MS technique (Chen *et al.* 2020). The concentrations of several heavy metal ions were quantified using the ICP-MS technique using an AGILENT ICPMS (Model - 7800) (Kumar *et al.* 2022). Heavy metal contamination levels are assessed using single metal and multiple metal pollution indices (Islam *et al.* 2015b). The single metal pollution indices (SPI) include contamination

factor (CFR) and ecological risk factor (ERFR). The multi-metal pollution indices (MMPI) include contamination degree (CDR) and modified contamination degree (mCDR), pollution load index (PLIX), and potential ecological risk index (PERI).

The contamination of aquatic ecosystems can be assessed using the concentration of metal (C_m) and the degree of metal contamination (D_m). It was calculated using the formula given by Hakanson (Dipti *et al.* 2023). The $CFR < 1$ indicates less contamination, where as $CFR > 6$ indicates very high contamination. In this, CFR value in between 1 to 3 indicates moderate contamination and 3 to 6 indicates considerable contamination. The heavy metal ecological risk factor (HER) is the value of the contamination factor multiplied by its "toxic-response" factor (TRF). The $HER < 40$ indicates low potential ecological risk where as $HER > 320$ resembles very high ecological risk. HER in between 40 to 50 indicates moderate ecological risk, 80 to 160 shows considerable ecological risk, and 160 to 320 resembles high ecological risk. Contamination degree (CDR) of aquatic ecosystem can be calculated by the addition of all metal contamination factors in all the study locations. Modified contamination degree (mCDR) of aquatic ecosystem was calculated by dividing the contamination degree with the 'n' number of analyzed elements (Villarin and Merel, 2020). Classification criteria for multi-metal contamination degree and modified contamination degree is presented in Table 1.

Table 1 Classification criteria of CDR and mCDR

Degree of Contamination	CDR value	Modified degree of contamination	mCDR value
Low concentration	$CDR < 6$	Unpollution	$mCDR < 1.5$
Moderate concentration	$6 \leq CDR < 12$	Slightly pollution	$1.5 \leq mCDR < 2$
Considerable concentration	$12 \leq CDR < 24$	Moderately pollution	$2 \leq mCDR < 4$
Very high contamination	$CDR \geq 24$	Considerably pollution	$4 \leq mCDR < 8$
		Highly pollution	$8 \leq mCDR < 16$
		Strongly pollution	$16 \leq mCDR < 32$
		Extremely pollution	$mCDR \geq 32$

85

86 **Pollution load index (PLIX):** PLIX explicit the overall toxicity level in the water body and it
87 can be calculated by the n^{th} root of multiplied contamination factor (CFR). The PLIX value in
88 between 2 to 3 indicates moderate pollution, 3 to 4 indicates moderate to high, 4 to 5 indicates
89 high pollution and >5 resembles extremely pollution. Potential ecological risk index (PERI) is
90 extensively applied to calculate the potential ecological deficit which was caused by the heavy
91 metals (Wan *et al.* 2020). This process integrates various interdisciplinary fields such as bio-
92 toxicology, and ecology to reflect the impacts of different toxic elements on the various
93 ecosystems and their comprehensive effects too. PERI is the sum of all HER which were
94 calculated at every sampling location. $PERI < 65$ indicates low risk, 130-260 indicates
95 considerable risk, and >260 indicates high risk.

96 **Results and Discussion**

97 Various pollution indices were calculated with respect to Cr, Mn, Pb, Fe, Zn, Cd, and Hg⁵. The
98 local background values and toxic factor values for the reported heavy metals are available in the
99 study area (Monteiro *et al.* 2021). Overall, nine sampling points (SP 1, SP 2, SP3, SP 4, SP 5, SP
100 6, SP 7, SP 8, AND SP 9) were selected to collect the water samples from the study area, which

covers the entire aquatic body. Many heavy metals in the Bollaram pond were observed at different study sites (Maruthai *et al.* 2025). The mean values of the heavy metal concentrations in the nine sampling sites are presented in Figure 1.

The Chromium (Cr) concentration in the study area is 4.3 ppm, which exceeds the prescribed limit of BIS (0.1 ppm) at all the sampling points (SP 1 to SP 9). The concentration of Mercury (Hg) is identified as 0.4 ppm, which exceeds the prescribed limit of BIS (0.001 ppm) at all the sampling points (SP 1 to SP 9). The Cadmium (Cd) concentration in the study area is 0.6 ppm, and Zinc (Zn) is 1.2 ppm, which are under the permissible limits of BIS, i.e., 3 ppm and 5 ppm, respectively in all the sampling points (SP 1 to SP 9). The concentration of Lead (Pb) is reported as 2.5 ppm, which exceeds the prescribed limit of BIS (0.1 ppm) at all sampling points (SP 1 to SP 9). Iron (Fe) concentration is reported as 426.2 ppm, and Manganese (Mn) is reported as 326 ppm, which exceeded the permissible limit of BIS, i.e., 5 ppm in all the sampling points (SP 1 to SP 9). These results showed that this lentic water system is highly contaminated with heavy metals. Fe, Mn, and Cr metals are present at high levels. Hg and Pb heavy metals are present at moderate contamination levels. Zn and Cd levels are reported below the BIS permissible limits in the study area. The surrounding area was surveyed manually to identify the main reasons for these heavy metal concentrations in the Bollaram lentic aquatic system. The survey identified that the runoff water from cement and iron-casting-based small-scale industries and wastewater discharge from the surrounding settlements through open drainage channels are reaching the study area. These water channels are active in the monsoon season and dry in the summer season (Xiong *et al.* 2016).




Fig 1: Concentrations of evaluated Heavy metals in the study area

The average contamination factor (CFR) of Fe, i.e., 236.79, indicates very high contamination (CFR>6). The CFR for Fe is reported in the range of 146.67 to 313.33 in the study area. The average CFR of Cr, i.e. 0.9, indicated low contamination (CFR<1), and the range of CFR reported in all the sampling points is between 0.73 to 1.10. The average CFR for Pb is 10.32, which indicates very high contamination (CFR>6). The reported Pb for CFR is in the range of 6.67 to 12.08. The average CFR for Cd is reported as 0.26, which indicates low concentration (CFR<1). Its CFR range is between 0.04 to 0.49 for all the sampling points. The average CFR for Mn is 1811.11, which indicates a very high CFR (CFR>6), and its range is between 1577.78 and 2122.22. The average CFR for Zn is 1.23, which indicates moderate contamination ($1 \leq \text{CFR} < 3$), and the CFR range is between 0.42 to 2.00. The average contamination factor for Hg is 1.89, which indicates moderate contamination ($1 \leq \text{CFR} < 3$), and the CFR range is reported between 0.5 to 4.00 in all sampling sites. Overall, the decreasing trend of CFR through various heavy metals was observed as: $\text{Mn} > \text{Fe} > \text{Pb} > \text{Hg} > \text{Zn} > \text{Cr} > \text{Cd}$ (Figure 2). Among all heavy metals, Mn and Fe showed extreme dominance in all the sampling sites with very high CFR. These results signify the heterogeneous anthropogenic pollution sources, exhibiting that these toxic heavy metals can

cause potential carcinogenic and non-carcinogenic health risks to the surrounding living organisms (Zzaman, 2015; Li *et al.* 2020)).




Fig 2: Contamination factor, mCDR, CDR of identified heavy metals

The heavy metal ecological risk factor (HER) in the study area was in the following descending order: $Mn > Fe > Pb > Hg > Cd > Cr > Zn$. Conceivable variations were noticed for HER of individual elements, signifying that the ecological risk of these heavy metals varied from one sampling point to the other (Liu *et al.* 2020 and Ma *et al.* 2020). Fe showed considerable variations in HER. It ranged from 146.67 to 313.33, which resembled considerable ecological risk to the high ecological risk. Cr, Hg, Zn, and Cd showed low potential ecological risk, which is less than 40 in all the sampling points. HER for Pb varied from 33.33 to 85.42, which signifies low potential ecological risk in four sampling points, moderate ecological risk in another four sampling points, and considerable ecological risk in one sampling point. In all the sampling points, Mn showed a very high ecological risk, which is above 320. It varies between 1360.67 to 2144.44.

The contamination assessment for all the sampling sites was done based on the Contamination degree (CDR) and mCDR (Fig 2). The results clearly showed that the contamination degree

(CDR) is very high ($CDR \geq 24$) in all the sampling sites. In this particular, the sampling sites 4 and 6 showed the highest CDR values (2449.34 and 2437.48, respectively). The minimum CDR values reported in sampling site (SP) 5 as 1572.15. The results depicted the mCDR with extreme pollution in all sampling sites ($mCDR > 32$) (Hafizure *et al.* 2012; Yadav and Yadav, 2018). The lowest mCDR value is identified as 224.59 at sampling site 8, and the highest mCDR values are reported as 348.21 and 349.91 at sampling points 6 and 4, respectively.

Pollution Load Index (PLIX) and Potential ecological risk index (PERI)

A pollution load index (PLIX) value equal to zero signifies perfection, 1 represents only the occurrence of the baseline level of the various contaminants, whereas a value of more than 1 reflects progressive water contamination by the trace metals. As per these grades in this study, the water samples collected from various sampling points were considerably contaminated with heavy metals (Figure 3) as the PLIX value was more significant than 1 (13). The highest PLIX value was perceived at SP6 (10.23), followed by SP2 (8.79). The lowest PLIX value was obtained at SP4 (5.22), followed by SP7 (5.91). In the remaining sampling points, PLIX values were perceived to be around 7. All these sampling points showed very high pollution with the heavy metals respectively. An elevated level of PLIX in the sampling sites reported that the disposal of municipal waste and sludge from small-scale industries illegally might have caused ecological risk to the water ecosystems.




Fig 3 Pollution Load Index (PLIX) and Potential ecological risk index (PERI)

The potential ecological risk index (PERI) indicates the sensitivity of the biological communities to numerous toxic substances and demonstrates the probable ecological risks resulting from trace metals. The PERI of the study area was in the following order: SP6> SP4> SP2> SP5> SP1> SP3> SP7> SP9> SP8. Overall, the PERI value for all sampling sites varied from 1629.41 to 2505.37, reflecting very high ecological risks.

Conclusion

This study confirmed that heavy metals from the Bollaram aquatic body showed a high degree of contamination based on calculated values of different pollution indices. However, the dumping of pharma sludge at the boundaries of Bollaram Pond and runoff water from the nearby iron-based small-scale industries highly contaminated this study area. The toxic situation in the nine studied sampling points showed extreme dominance by Fe and Mn. Cr and Cd showed less contamination factor. Similarly, the pollution load index also signifies "very high Polluted" in all the study sites. Seasonal variations are the main limitations for this study. A pollution load index (PLIX) value equal to zero signifies perfection, 1 represents only the occurrence of the baseline

level of the various contaminants, whereas a value of more than 1 reflects progressive water contamination by the trace metals. The highest PLIX value was perceived at SP6 (10.23), followed by SP2 (8.79). The lowest PLIX value was obtained at SP4 (5.22), followed by SP7 (5.91). Hence, continuous monitoring of contaminants at a regular time interval is necessary after restricting the waste disposal at the boundaries of Bollaram pond. During the summer season, when the water quantity is less, the local authority has to take action to remove the sludge accumulated at the bottom of the water body. Also, it is necessary to provide artificial aeration to reduce the degree of contamination, which helps to restore the pond water quality.

References:

1. Adimalla N. (2020), Heavy metals contamination in urban surface soils of Medak province, India and its risk assessment and spatial distribution, *Environmental Geochemistry and Health*, 42(1), 59-75.
2. Chen X.X., Liu Y.M., Zhao Q.Y., Cao W.Q. and Chen X.P. (2020), Health risk assessment associated with heavy metal accumulation in wheat after long term phosphorus fertilizer application, *Environmental Pollution*, 262, 1143-48.
3. Cheng, X., Wei, C., Ke, X., Pan, J., Wei, G., Chen, Y., Wei, C., Li, F. and Preis, S. (2022), Nationwide review of heavy metals in municipal sludge wastewater treatment plants in china: sources, composition, accumulation and risk assessment, *Journal of Hazardous Materials*, 437, 129267.
4. Dipti, Kumar P., Dwivedi S. and Singh R.P. (2023), Assessment of Heavy metal contaminations in agro-ecosystems around distinct pollution sources, its Spatial distribution and Principal component analysis, *Res. J. Chem. Environ*, 27(7), 74-83.
5. Hafizur Rahman S., Khanam D., Mehedi Adyel T., Shahidul Islam M., Aminul Ahsan M. and Akbor A. (2012), Assessment of heavy metal contamination of agricultural soil around Dhaka export processing zone (DEPZ), Bangladesh: implication of seasonal variation and indices, *Appl Sci.*, 2, 584–601.

6. He, H., Peng, M., Ru, S., Hou, Z. and Li J. (2022), A suitable organic fertilizer substitution ratio could improve maize yield and soil fertility with low pollution risk, *Front. Plant Sci.*, 13, 988663.
7. Islam M.S., Ahmed M.K., Al-Mamun M.H. and Hoque M.F. (2015b), Preliminary assessment of heavy metal contamination in surface sediments from a river in Bangladesh, *Environmental Earth Science*, 73, 1837–1848.
8. Kowalska J.B., Mazurek R., Gąsiorrek M. and Zaleski T. (2018), Pollution indices as useful tools for the comprehensive evaluation of the degree of soil contamination—A review, *Environmental Geochemistry and Health*, 40(6), 2395-2420.
9. Kumar P., Dipti, Kumar S. and Singh R.P. (2022), Severe contamination of carcinogenic heavy metals and metalloid in agroecosystems and their associated health risk assessment, *Environ Pollut*, 301, 118953.
10. Li X., Zhang J., Gong Y., Liu Q., Yang S., Ma J., Zhao L. and Hou H. (2020), Status of copper accumulation in agricultural soils across China (1985–2016), *Chemosphere*, 244, 125516.
11. Liu, Y., Liu, D., Zhang, W., Chen, X., Zhao, Q., Chen, X. and Zou, C. (2020), Health risk assessment of heavy metals (Zn, Cu, Cd, Pb, As and Cr) in wheat grain receiving repeated Zn fertilizers, *Environ. Pollut.*, 257.
12. Ma, J., Chen, Y., Antoniadis, Vasileios. Wang, K., Huang, Y. and Tian, H. (2020), Assessment of heavy metal(loid)s contamination risk and grain nutritional quality in organic waste amended soil, *J. Hazard. Mater.*, 399.
13. Maruthai, S. *et al.* (2025), Wastewater Recycling Integration with IoT Sensor Vision for Real-time Monitoring and Transforming Polluted Ponds into Clean Ponds using HG-RNN, *Global NEST Journal* [Preprint]. Available at: <https://doi.org/10.30955/gnj.06758>.
14. Monteiro, L., Cristina, R., and Covas D. (2021), Water and energy efficiency assessment in urban green spaces, *Energies*, 14(17) , 5490.
15. Saha M.K., Sarkar R.R., Ahmed S.J., Sheikh A.H., and Mostafa M.G. (2021), Impacts of brick kiln emission on agricultural soil around brick kiln areas, *Nepal Journal of Environmental Science*, 9(1), 1-10.

- 245 16. Selvanarayanan R., Rajendran S., Pappa C. K. and Thomas B. (2024), Wastewater
246 recycling to enhance environmental quality using fuzzy embedded with rnn-iot for
247 sustainable coffee farming, *Global NEST Journal*, 26(8), 06346.
- 248 17. Su, C., Wang, J., Chen, Z., Meng, J., Yin, G., Zhou, Y. and Wang, T. (2023), Sources and
249 health risks of heavy metals in soils and vegetables from intensive human intervention
250 areas in South China, *Sci. Total Environ*, 857, 159389.
- 251 18. Venkatraman M., Surendran R., Srinivasulu S. and Vijayakumar K. (2024), Water quality
252 prediction and classification using attention based deep differential recurflownet with
253 logistic giant armadillo optimization, *Global NEST Journal*, 27(1), 06799'
- 254 19. Villarín MC. and Merel S. (2020), Paradigm shifts and current challenges in wastewater
255 management, *J Hazard Mater.*, 15, 390, 122139.
- 256 20. Wan, Y., Huang, Q., Wang, Q., Yu, Y., Su, D., Qiao, Y. and Li H. (2020), Accumulation
257 and bioavailability of heavy metals in an acid soil and their uptake by paddy rice under
258 continuous application of chicken and swine manure, *J. Hazard. Mater.*, 384, 121293.
- 259 21. Xiong T., Austruy A., Pierart A. and Shahid M. (2016), Kinetic study of phytotoxicity
260 induced by foliar lead uptake for vegetables exposed to fine particles and implications for
261 sustainable urban agriculture, *Journal of Environmental Science*, V(1), 12.
- 262 22. Yadav A. and Yadav P.K. (2018), Pollution Load Index (PLI) of Field Irrigated with
263 Wastewater of Mawaiya Drain in Naini Suburbs of Allahabad District, *Current World*
264 *Environment*, 13(1), 159.
- 265 23. zzaman M. (2015), Trace elements in different land use soils of Bangladesh and potential
266 ecological risk, *Environmental Monitoring and Assessment*, 187(9), 587.