

# Application of Selenium and Silica Nanoparticles on Pepper Seedlings for Controlling Root Rot Disease caused by *Rhizoctonia solani*

Naglaa A. Taha<sup>1</sup>, Yousry A. Bayoumi<sup>2</sup>, Hassan El-Ramady<sup>3\*</sup>, Tarek A. Shalaby<sup>4\*</sup>, Hossam S. El-Beltagi<sup>5</sup>, Mohamed S. AL-Saikhan<sup>4</sup>, Saleh M. Alturki<sup>4</sup> and Amr A. Khedr<sup>6</sup>

<sup>1</sup> Plant Pathology Research Institute, Agriculture Research Center, 12619 Giza, Egypt (N.A.T) ([naglaa\\_abdelbaset@yahoo.com](mailto:naglaa_abdelbaset@yahoo.com))

<sup>2</sup> Horticulture Department, Faculty of Agriculture, University of Kafrelsheikh, Kafr El-Sheikh 33516, Egypt (Y.B.)

([yousry.bayoumi@agr.kfs.edu.eg](mailto:yousry.bayoumi@agr.kfs.edu.eg))

<sup>3</sup> Soil and Water Department, Faculty of Agriculture, University of Kafrelsheikh, Kafr El-Sheikh 33516, Egypt (H.E.R.)

([hassan.elramady@agr.kfs.edu.eg](mailto:hassan.elramady@agr.kfs.edu.eg))

<sup>4</sup> Department of Arid Land Agriculture, College of Agricultural and Food Sciences, King Faisal University, Al-Ahsa 31982, Saudi Arabia ([tshalaby@kfu.edu.sa](mailto:tshalaby@kfu.edu.sa), [msaikhan@kfu.edu.sa](mailto:msaikhan@kfu.edu.sa), [smturki@kfu.edu.sa](mailto:smturki@kfu.edu.sa)) (T.A.S, M.S.A, S.A)

<sup>5</sup> Agricultural Biotechnology Department, College of Agricultural and Food Sciences, King Faisal University, Al-Ahsa 31982, Saudi Arabia ([helbeltagi@kfu.edu.sa](mailto:helbeltagi@kfu.edu.sa)) (H.S.E)

<sup>6</sup> Agricultural Botany Department, Faculty of Agriculture, University of Kafrelsheikh, Kafr El-Sheikh 33516, Egypt ([amr3400@yahoo.com](mailto:amr3400@yahoo.com); A.A.K)

\*Corresponding author: Tarek Shalaby ([tshalaby@kfu.edu.sa](mailto:tshalaby@kfu.edu.sa)); Hassan El-Ramady ([hassan.elramady@agr.kfs.edu.eg](mailto:hassan.elramady@agr.kfs.edu.eg))

## Abstract

There is an urgent need to find sustainable alternatives to chemical pesticides in order to protect the environment, ensure healthy food production and improve seedling quality. Therefore, the current study is an attempt to find nano-alternatives to chemical fungicides for protecting pepper seedlings (*Capsicum annuum* L.) production from the phytopathogen of *Rhizoctonia solani* as well as to enhance their growth and quality by individually applying nanosilica and nano-selenium (n-SiO<sub>2</sub>, n-Se), and their combination under nursery conditions. Single and combined applied doses of n-SiO<sub>2</sub> (200 mg L<sup>-1</sup>) and n-Se (100 mg L<sup>-1</sup>) were treated pepper seedlings for controlling the root rot disease of *R. solani*. Seedlings treated with nanosilica (200 mg L<sup>-1</sup>) produced the highest values in the studied parameters (vegetative growth of pepper seedlings, enzymatic antioxidants and the photosynthetic parameters under the fungal infection). The proposed mechanism of nano silica in promoting the growth of pepper seedlings against *R. solani* was supported by the Scanning Electronic Microscope (SEM) images, which compared infected seedlings (control) with nanosilica-treated ones. A similar trend was observed for enzymatic antioxidants responses after nanosilica treatment, indicated enhanced resistance of pepper seedlings against studied fungal pathogen. This superior performance was followed by nano-Se treatment and then by combined application of both nanoparticles. This study provides a novel insight into the potential use of nanofungicides against the studied fungal pathogen (*R. solani*) to reduce the chemical of fungicides application and improve the growth and quality of pepper seedlings. Further research is recommended to investigate additional nanoparticles and new phytopathogens.

**Keywords:** Antioxidants, Disease severity, Pathogenicity, Phytopathogen, Scanning Electronic Microscope

## 1. Introduction

Pepper (*Capsicum annuum* L.) is as an important vegetable crop worldwide cultivated under both greenhouse and open field. In Egypt, the cultivated area of pepper covered approximately 57,683 ha with a total production of about 18.47 Mg ha<sup>-1</sup> (FAOSTAT, 2025). The use of high-quality transplants is a common commercial practice in widespread production of peppers and most of vegetables for fast-sustained establishment joined with improvement of identical maturity, early and total yields beside quality properties (Hoffmann and Poorter, 2002). Pepper seedlings production faces several challenges due to climate change including abiotic and biotic stresses. Several studies have confirmed abiotic stresses can cause drastic, rapid and negative impacts on seedling growth and productivity such as metal stress (Altaf et al. 2022; Yu et al. 2025), low temperature (Li et al. 2023; Song X et al. 2023; Pu et al. 2025), heat stress (Jang et al. 2025), drought (Lu et al. 2023), and salinity stress (Yuce et al. 2025). Regarding biotic stresses, various microorganisms including fungi, bacteria, fungi-like viruses, and nematodes are able to cause diseases during the growth of pepper seedlings alongside soil borne pathogens (Wu et al. 2019; González-Hernández et al. 2021; El-Kazzaz et al. 2022; Hernández-Huerta et al. 2023; Xiao et al. 2025).

Soil-borne fungal diseases caused by many microbial genera such as *Rhizoctonia*, *Fusarium*, *Pythium*, and *Verticillium*), are among the most destructive pathogens affecting vegetable seedlings (Shalaby et al. 2022). Several studies reported that the controlling of such phyto-pathogen using environmentally friendly alternatives to chemical pesticides to protect both the environment and human health (He et al. 2023; Paz-Trejo et al. 2024; Yadav et al. 2025). In general, these alternatives are considered biological control agents having sustainable approaches, and eco-friendly approaches such as microbial inoculants (Hernández-Huerta et al. 2023; Xiao et al. 2025), nano-pesticides (Qiao et al. 2024; Zainab et al. 2024), nanomaterials (Kashyap et al. 2024; Vijayreddy et al. 2025), and plant growth regulators like paclobutrazol (Shalaby et al. 2022a,b). Among soil-borne fungi *Rhizoctonia solani* is one of the most an important pathogen, posing serious threats and causing yield losses in various crops (Huang et al. 2025) such as rice (Senapati et al. 2022), quinoa (Chethan et al. 2025), tomato (Kirupanatha-Rajan et al. 2025), faba bean (Osman et al. 2025), pigeon pea (Gao et al. 2025), and pepper (Rodríguez et al. 2023).

Nano selenium (n-Se) and nano-silica (n-SiO<sub>2</sub>) are considered promising nanoparticles (NPs) due to their anti-stress properties potential applications in agriculture (Gao and Tuda 2024). These NPs act as nano-fungicides capable of supporting plant growth under stress by stimulating biochemical, morphological, and molecular defenses mechanisms (Taha et al. 2023). The significance of such NPs was reported in numerous studies according to their low dose requirement; targeted supply improved bioavailability and organized release; low dose-dependent toxicity; more solubility and penetrability (Periakaruppan et al. 2023). The individual effects of n-Se and n-SiO<sub>2</sub> under biotic stress have been discussed in several studies (Goswami et al. 2022; Kang et al. 2023; Sulaiman et al. 2023; Shahbaz et al. 2023; Song J et al. 2023; Batista et al. 2025; Choudhary et al. 2025; Imran et al. 2025). However, few studies have examined the combined use of nano-Se

and nano-silica to alleviate stress such as drought in strawberry (Zahedi et al. 2020), salinity in rice (Badawy et al. 2021), and *Alternaria* leaf spot disease in common bean (Taha et al. 2023).

Therefore, the present study was designated to investigate the individual and combined roles of n-Se and n-SiO<sub>2</sub> in controlling damping-off disease caused by *Rhizoctonia solani* in pepper seedlings. The effects of these nano-fungicides on the growth and quality of pepper seedlings were comparing with the commercial fungicide and control treatments. This study also highlights the isolation and purification of the studied pathogen, *in vitro* and greenhouse experiments to produce pepper seedlings under disease stress, including chlorophyll content, and antioxidant enzymatic activities in pepper seedling leaves.


## 2. Materials and Methods

### 2.1 Pathogen isolation, purification and pathogenicity tests

Three isolates of *Rhizoctonia solani* were collected from infected pepper seedlings showing damping-off symptoms at commercial nurseries in the Delta region, Egypt (Latitude 31°06'25.20" N Longitude: 30°56'26.99" E), (**Figure 1**). In brief, the infected pepper seedlings with damping off disease were collected and the pathogen was isolated, and purified following the method described by Dhingra and Sinclair (1985). The morphological traits and microscopic features of the isolates were confirmed as *R. solani* according to Leslie and Summerell (2006). For the pathogenicity test, the three isolates were evaluated for their ability to cause disease on pepper seedlings (California Wonder, cv.), which were obtained from the common vegetable nurseries under greenhouse conditions. Detailed procedures are described in Shalaby et al. (2022). The trials were organized in four replications, each one included three Styrofoam trays for each isolate alongside the non-infected control without the pathogen. Pre- and post-emergence damping-off was recorded 20 and 40 days after sowing, respectively, using pepper seedlings grown in Styrofoam trays (209 cells). The following equation describes the calculation of the pre- and post-emergence damping-off:

$$\text{Damping-off (\%)} = (\text{No. of infected plants} / \text{total No. of plants}) \times 100$$

Among the three isolates, the isolate Rs2 exhibited the highest pathogenicity based on symptoms severity and damping-off incidence. The identified isolates fungus was known as the pathogen of *R. solani* at the Mycology and Disease Survey Research Department, Plant Pathology Research Institute, Agricultural Research Center (ARC), Giza, Egypt.



**Figure 1.** An outline on the design of the present study including various steps: photo (1) the isolation and purification of the studied pathogen, (2) *in vitro* experiment, (3) greenhouse experiment, (4) disease assessments, (5) production of pepper seedlings under disease stress, and (6) the main target of this study “healthy pepper seedlings”.

## 2.2 Preparing nanoparticles

The nanoparticles were provided by from Department of Agricultural Microbiology Research, A.R.C., Giza, Egypt. Nano selenium (n-Se) was synthesized biologically using *Bacillus cereus* TAH strain, with particle sizes ranged from 41 to 102 nm (Ghazi et al. 2022). The transmission electron microscopy (TEM) was used to measure the magnitude of n-Se, with high-resolution (HR-TEM, Tecnai G20, FEI, The Netherlands), at the Central Laboratory of Nanotechnology and Advanced Material, ARC, whereas n-SiO<sub>2</sub> was prepared by a physical method. Its main characterization includes an average diameter of 10 nm for, pH 4–4.5, and specific surface area ranges from 260 to 320 m<sup>2</sup> g<sup>-1</sup>.

## 2.3 Antifungal activity of studied nanoparticles

Three isolates of *Rhizoctonia solani* with different degree of pathogenicity were evaluated. The most aggressive isolate (Rs2) was used for subsequent *in vitro* tests. The applied dose of n-Se was 100 mg L<sup>-1</sup>, whereas n-SiO<sub>2</sub> was 200 mg L<sup>-1</sup>. The combined treatment of both included using ½ of the used doses (i.e., 50 mg L<sup>-1</sup> n-Se + 100 mg L<sup>-1</sup> n-SiO<sub>2</sub>). All used concentrations of the NPs were mixed with the PDA media (100 ml). Every dosage had five replications in Petri plates. 7 days after fungal growth, 5 mm in diameter agar

masses from the actively growing edge of *R. solani* were placed at the center of Petri dishes containing the doses of nanoparticles. Plates were incubated at 25°C for 7 days. Colony diameters were measured, and the reduction in mycelial growth was calculated using the formula described by Ferreira et al. (1991):

$$\text{Reduction rate (\%)} = \frac{R - r}{R} \times 100$$

Where (R) is the radial growth of fungi in control and (r) is radial growth of fungi in treated plates.

## 2.4 Greenhouse experiment

The greenhouse experiments were conducted at the Faculty of Agriculture, Kafrelsheikh University, during the two successive seasons of 2023 and 2024. This study aimed to evaluate the effectiveness of n-SiO<sub>2</sub>, n-Se and their combinations in controlling root rot caused by *R. solani* and improving pepper seedlings growth and quality under greenhouse conditions. The root rot-susceptible pepper cultivar (California Wonder) was used in all experiments. Seeds were sown in 209 cells trays containing 1:1 v/v mixture of coco peat and vermiculite. The trays were irrigated daily with equal amount of water to maintain field capacity and seedling were fertilized weekly after full germination with a compound fertilizer including 400 N, 400 P<sub>2</sub>O<sub>5</sub>, 400 K<sub>2</sub>O at the rate of 1 g L<sup>-1</sup>. The treatments are presented in **Table 1**.

**Table 1:** Details of applied treatments on pepper seedlings during both growing seasons

| Code | Treatments                        | Description of the treatment(s)                                                                                                                        |
|------|-----------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------|
| T1   | Control (negative control)        | Healthy seedlings treatment without any infection                                                                                                      |
| T2   | Control (positive control)        | Seedlings infected with <i>Rhizoctonia solani</i>                                                                                                      |
| T3   | Fungicide                         | Seedlings infected by <i>R. solani</i> and applied commercial fungicide Rizolex 50% WP                                                                 |
| T4   | nano-silica (n-SiO <sub>2</sub> ) | Seedlings infected by <i>R. solani</i> and applied 200 mg L <sup>-1</sup>                                                                              |
| T5   | nano-Se (n-Se)                    | Seedlings infected by <i>R. solani</i> and applied 100 mg L <sup>-1</sup>                                                                              |
| T6   | ½ (n-Se + n-SiO <sub>2</sub> )    | Seedlings infected by <i>R. solani</i> and applied 50% of T4 and T5 (i.e., n-SiO <sub>2</sub> = 100 mg L <sup>-1</sup> + n-Se= 50 mg L <sup>-1</sup> ) |

Healthy pepper transplantings (uninfected by the pathogen of *R. solani*) were utilized as a negative control, while transplants infected by *R. solani* represented the positive control. To prepare the inoculated growing media, barley grains infected with *R. solani* pathogen were mixed individually with growing substrate at 1:1 (w/w) ratio, then placed directly into trays. Based on a preliminary trial, concentration of 100 mg L<sup>-1</sup> for n-Se and 200 mg L<sup>-1</sup> for n-SiO<sub>2</sub> were selected for this study. The combined treatments, n-Se and n-SiO<sub>2</sub> were applied at 50 and 100 mg L<sup>-1</sup>, respectively. The fungicide (Rizolex 50% WP, which imported from Japan) produced in Japan by Company of Sumitomo Chemical was applied at the recommended rate of 2 g/L to the growing media under the current pathogen (*R. solani*). Six treatments were applied by adding solutions directly prior to seeds planting. After planting, the trays were maintained under greenhouse conditions at 25 ± 2

°C for 40 days. The experiment arranged in four replications; each replication containing three trays per treatment besides untreated control treatment, which was trays with pathogen-free (non-diseased). All treatments were repeated again after 15 days from sowing, once seeds had fully germinated by sprinkling the solutions onto trays until the drops penetrated the growing media and reached the roots.

## 2.5 Disease assessments

The pre- and post- emergence percentages of pepper transplants were recorded after 20 and 40 days from sowing, as reported by Chandler and Santelman (1968). The disease severity was assessed after 40 days using the scale of 5-rating (Altier and Thies, 1995) (1 = vigorous and non-diseased, 2 = principal root end necrotic, 3= principal root end spineless and rotten, 4 = dead transplantings, emergence seeds with rotten essential roots and 5 = dead seed, un-germinated seed). The following equation was applied to calculate disease severity percent:

$$\text{Disease severity, \%} = \frac{(\text{rating no.} \times \text{no.of seedlings in rating})}{(\text{Total no.of seedlings} \times \text{highest rating})} \times 100$$

## 2.6 Growth Assessments of Transplantings

After 40 days of sowing, vegetative growth parameters were measured; including seedling height (cm), stem diameter (mm), fresh and dry weights of seedlings (g/seedling), and dry weight was registered after drying at 70 °C for 48 h.

## 2.7 Chlorophyll determination

Leaf chlorophyll content was determined according to Nagata and Yamashita (1992) using a 645 and 663 nm wave length at spectrophotometer and calculating by relieving the analysis as follows:

$$\text{Chl. a} = 0.999 \times A_{663} - 0.0989 \times A_{645}$$

$$\text{Chl. b} = -0.328 \times A_{663} + 1.77 \times A_{645}$$

As reported by Maxwell and Johnson (2000), the highest efficacy of the photosystem (Fv/Fm) was assessed with a movable Optic-Science OS-30p + Fluorometer (Inc., Hudson NH, USA).

## 2.8 Antioxidant enzyme activities and electrolyte leakage

Fresh leaf tissue (0.5 g) was homogenized in 3 ml of 0.05 M Tris buffer (pH 7.8), comprising 0.001 M EDTA–Na<sub>2</sub> and 7.5% Polyvinylpyrrolidone at 0–4°C for enzymes activity. Homogenates were recorded by spectrophotometer (Shi-madzu, Japan). Catalase (CAT), peroxidase (POX), and polyphenol oxidase enzyme (PPO) activities were determined according to Aebi (1984), Rathmell and Sequeira (1974), and Malik and

Singh (1980), respectively. Electrolyte leakage was measured using EC meter or electrical conductivity meter (Lutts et al. 1996). All vegetative growth traits of pepper seedlings, chlorophyll content, chlorophyll florescence, electrolyte leakage, and antioxidant enzymes were measured 40 days after sowing in both seasons.

### 2.9 Microscopic investigation

The microscopic measurements were achieved using the fungal growth of the pathogen on the PDA media accompanied by n-SiO<sub>2</sub> application (microscope power of 200X by tacky bar touch process after 7 days' incubation). Scanning electron microscope (Model: SEM, JEOL JSM 6510 Iv, Japan) was performed at Nanotechnology Institute (Kafrelsheikh University, Egypt) to observe) the influence of the n-SiO<sub>2</sub> treatment at 200 mg L<sup>-1</sup> on growth performance changes of *Rhizoctonia solani*.

### 2.10 Statistical analyses

The experiment was arranged in a completely randomized design (CRD) with four replications. Statistical tests were achieved by the CoStat program package (CoHort Software, CA, USA; Version 6.303) by use of ANOVA and then Duncan's multiple range tests at 5% level of possibility used to match significant variances among averages of the treatments (Snedecor and Cochran, 1989).

## 3. Results


### 3.1. Pathogenicity evaluation

To evaluate the pathogenic aptitude of *Rhizoctonia solani* isolates, three isolates of the pathogen were tested in a greenhouse experiment. The most aggressive isolate was identified as Rs2 (*Rhizoctonia solani*), while Rs3 exhibited the lowest one (**Table 2 and Figure 2**). Rs2 isolate was recorded the highest disease severity (78%) as well as the highest pre- and post-emergence damping-off rates (67.13 and 15.51 %, respectively). On the other hand, the lowest survival seedlings rate (17.63%) was belonged to the isolate Rs2. Therefore, the Rs2 isolate of *Rhizoctonia solani* was selected for subsequent experiments.

**Table 2.** Pathogenicity test of three isolates obtained from pepper seedlings under

| Pathogen isolates               | Pre-emergence (%) | Post-emergence (%) | Survival seedlings (%) | Disease severity (%) |
|---------------------------------|-------------------|--------------------|------------------------|----------------------|
| Control (healthy)               | 2.50 d            | 0.001 c            | 97.50 a                | 0.001 d              |
| <i>Rhizoctonia solani</i> (Rs1) | 55.40 b           | 13.10 a            | 31.50 c                | 57.70 b              |
| <i>Rhizoctonia solani</i> (Rs2) | 67.13 a           | 15.51 a            | 17.63 d                | 78.00 a              |
| <i>Rhizoctonia solani</i> (Rs3) | 35.77 c           | 7.88 b             | 56.35 b                | 45.87 c              |
| F. test                         | **                | **                 | **                     | **                   |

\*\* indicates highly significant treatment and values of means in each column followed by the same letter are not significantly at level of  $p < 0.05$



220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231

**Figure 2.** The three isolates (Rs1, Rs2, Rs3) of *Rhizoctonia solani* as investigated using Petri dishes and their morphological and microscopic parameters


**3.2. In Vitro antifungal activity of selected nanoparticles**

Individual and combined antifungal effects of n-Se and n-SiO<sub>2</sub> were examined *in vitro* experiment using isolate Rs2 of *R. solani*. The highest antagonistic result was found with fungicide application (i.e., synthetic control, which recorded reduction rate 94.44%), followed by n-SiO<sub>2</sub>, and n-Se recording reduction rate of 81.11% and 66.67%, respectively. All used nanoparticles statistically reduced mycelia growth of *R. solani* compared with the control treatment (**Table 3; Figure 3**).

**Table 3.** Responses of mycelial growth of *Rhizoctonia solani* to different nanoparticles under *in vitro* conditions and the corresponding reduction of the pathogen

| Treatments                                                                      | Mycelial growth (cm) | Reduction (%) |
|---------------------------------------------------------------------------------|----------------------|---------------|
| (T2) Negative control                                                           | 9.0 a                | 0.001 f       |
| (T3) Fungicide (Rizolex 50% WP)                                                 | 0.5 e                | 94.44 a       |
| (T4) n-SiO <sub>2</sub> (200 mg L <sup>-1</sup> )                               | 1.7 b                | 81.11 e       |
| (T5) n-Se (100 mg L <sup>-1</sup> )                                             | 3.0 c                | 66.67 cd      |
| (T6) 1/2 (n-SiO <sub>2</sub> + n-Se) (100+50 mg L <sup>-1</sup> , respectively) | 3.6 d                | 60.00 bc      |
| F. test                                                                         | **                   | **            |


234 \*\* indicates highly significant treatment and values of means in each column followed by the same letter are  
 235 not significantly at level of  $p < 0.05$  ( $\pm$ , SD =standard deviation; N= 5).  
 236  
 237



238  
 239 **Figure 3.** Images illustrating of the inhibition of *Rhizoctonia solani* mycelial growth due to applying of the treatments in the *in vitro*  
 240 experiments. Where, T2: control (positive control), T3: fungicide, T4: n-SiO<sub>2</sub> (200 mg L<sup>-1</sup>), T5: n-Se (100 mg L<sup>-1</sup>), and T6: ½ (n-Se  
 241 + n-SiO<sub>2</sub>) (50 + 100 mg L<sup>-1</sup>, respectively)  
 242  
 243

### 244 3.3. Microscopic Investigation of the *R. solani*

245 The *Rhizoctonia* mycelium was examined using scanning electronic microscope (SEM) under *in vitro* con-  
 246 ditions. The obtained images were belonged the control and n-SiO<sub>2</sub> treatments, where a clear variation in the  
 247 mycelium growth was observed for n-SiO<sub>2</sub> compared with the negative control treatment (**Figure 4**). More-  
 248 over, an occurrence of alterations, along with the variation in mycelia development was observed seven days  
 249 after inoculation as observed by SEM. The application of n-SiO<sub>2</sub> (200 mg L<sup>-1</sup>) induced notable morphological  
 250 modifications in hyphae and mycelia growth. Mycelium and hyphae of the *R. solani* was severely damaged  
 251 and shriveled in the presence of n-SiO<sub>2</sub> (**Figure 4B**) with the secondary mycelium appearing folded and  
 252 damaged, as compared to negative control that was un-treated and showed typical mycelial structures with a  
 253 smooth surface, while the growth of secondary mycelium was complete as typical fungus morphological  
 254 features (**Figure 4A**).  
 255



256  
 257 **Figure 4.** Scanning electron microscope (SEM) investigation of *Rhizoctonia solani* comprising with applied control (A) and with  
 258 applied n-SiO<sub>2</sub> at dose of 200 ppm (B) on PDA media. Scale bars in both A and B are 20  $\mu$ m with x 950  
 259

260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282

**3.4. Disease Severity of pepper seedlings under Greenhouse Conditions**

Under the greenhouse experiments, survival seedlings, disease severity and its efficacy along with pre-, and post-emergence percentage of pepper transplanting's infected with *R. solani* were studied compared to the control and healthy ones (Table 4 and Figure 5). Disease severity was significantly affected by all applied treatments during both seasons. The highest disease severity values were recorded in the positive control (T2) treatment during the two growing seasons (76.70 and 80.06 %, respectively). The seedlings treated by nano-silica (T4) exhibited the lowest disease severity (15.93 and 13.93 %) in both seasons, respectively, which was nearly comparable to the results obtained with fungicide application. The pre- and post-emergence as well as the survival rate of seedlings, were measured as shown in Table 4 and showed the same tendency in both seasons. On the other hand, the combined application of nanoparticles (n-Se+n-SiO<sub>2</sub>) recorded a similar result of survival seedlings comparable to fungicide treatment under infection by tested fungus pathogen with no significant differences between them in most cases. The feature of pepper seedlings after 40 days of infection with *R. solani* under greenhouse conditions for each treatment is shown in Figure 5. It is clear that seedlings treated with nano-silica (T4) showed growth performance closely similar to the control under infection conditions.

**Table 4.** Survival seedlings, disease severity, pre-, and post-emergence of pepper seedlings infected with *R. solani* compared to the healthy ones and control in both seasons

| Treatments           | Pre-emergence (%) | Post-emergence (%) | Survival seedlings (%) | Disease severity (%) | Efficacy (%) |
|----------------------|-------------------|--------------------|------------------------|----------------------|--------------|
| First season (2023)  |                   |                    |                        |                      |              |
| T1                   | 3.71 d            | 00.001 e           | 96.29 a                | 00.001 c             | 00.001 d     |
| T2                   | 75.82 a           | 10.46 a            | 13.72 d                | 76.70 a              | 00.001 d     |
| T3                   | 16.99 c           | 3.92 d             | 79.08 b                | 10.24 d              | 86.65 a      |
| T4                   | 28.10 b           | 4.58 cd            | 75.82 b                | 15.93 c              | 79.23 b      |
| T5                   | 19.60 c           | 5.88 c             | 66.03 c                | 23.83 b              | 68.93 c      |
| T6                   | 31.37 b           | 7.84 b             | 60.79 c                | 26.70 b              | 65.19 c      |
| F-test               | **                | *                  | **                     | **                   | **           |
| Second season (2024) |                   |                    |                        |                      |              |
| T1                   | 5.23 e            | 0.001 e            | 94.78 a                | 0.001 d              | 0.001 c      |
| T2                   | 77.78 a           | 9.80 a             | 12.42 d                | 80.06 a              | 0.001 c      |
| T3                   | 15.03 d           | 2.61 b             | 82.35 b                | 11.81 c              | 85.25 a      |
| T4                   | 24.84 c           | 5.23 d             | 79.08 b                | 13.93 c              | 82.60 a      |
| T5                   | 15.69 d           | 6.54 c             | 68.62 c                | 26.93 b              | 66.36 b      |
| T6                   | 30.29 b           | 8.49 b             | 61.22 c                | 28.59 b              | 64.29 b      |
| F-test               | **                | *                  | *                      | ***                  | **           |

Note: \* and \*\* indicate significant and highly significant, respectively  
Values of means in each column followed by the same letter are not significantly at  $p < 0.05$   
Where, T1: negative control (healthy seedlings), T2: control (positive control), T3: fungicide, T4: n-SiO<sub>2</sub> (200 mg L<sup>-1</sup>), T5: n-Se (100 mg L<sup>-1</sup>), and T6: ½ (n-Se + n-SiO<sub>2</sub>) (50 + 100 mg L<sup>-1</sup>, respectively)



**Figure 5.** Pepper seedlings after 40 days from infection with *R. solani* at greenhouse and applying the treatments, where T1: negative control (healthy seedlings), T2: control (positive control), T3: fungicide, T4: n-SiO<sub>2</sub> (200 mg L<sup>-1</sup>), T5: n-Se (100 mg L<sup>-1</sup>), and T6: ½ (n-Se + n-SiO<sub>2</sub>) (50 + 100 mg L<sup>-1</sup>, respectively)

### 3.5. Vegetative Growth and Photosynthetic Traits of pepper seedlings under biotic stress

After 40 days from seeds sowing, seedlings growth parameters, chlorophyll fluorescence and chlorophyll contents were measured (**Table 5**). In general, application of nano-silica recorded the highest value for almost all measured growth traits (seedling height, stem diameter, fresh and dry weights) compared with both fungicide treatment and untreated plants. a similar trend was observed during both growing seasons for the measuring photosynthetic parameters (chlorophyll fluorescence and content). The highest values of chlorophyll *a* and *b* were achieved by using single n-Se application or the combined n-Se and nano-silica treatment in both seasons. However, pepper seedlings infected only with *R. solani* exhibited the lowest values of all vegetative traits. Healthy (non-infected) pepper seedlings recorded the greatest height, diameter, fresh and dry weights of seedlings compared to all infected treatments with no-significant differences between T4 (nano-silica) and T3 (fungicide treatment) in both seasons. Therefore, the favorable seedlings quality traits as moderate height, thicker stem and greater dry mass and higher chlorophyll content was resulted from of individual application of nano-silica (T4) under the infection by *R. solani* compared with the other treatments.

**Table 5.** Chlorophyll content and fluorescence along with vegetative growth traits of pepper seedlings after 40 days after sowing during both seasons

| Treatments          | Seedling height (cm) | Stem diameter (mm) | Fresh weight (g/seedling) | Dry weight (g/seedling) | Chlorophyll Fluorescence (F <sub>v</sub> /F <sub>m</sub> ) | Chl. a (mg g <sup>-1</sup> FW) | Chl. b |
|---------------------|----------------------|--------------------|---------------------------|-------------------------|------------------------------------------------------------|--------------------------------|--------|
| First season (2023) |                      |                    |                           |                         |                                                            |                                |        |
| T1                  | 21.36 a              | 3.18 a             | 1.14 a                    | 0.118 a                 | 0.723 a                                                    | 4.78 b                         | 1.5 a  |
| T2                  | 5.18 e               | 1.85 c             | 0.46 d                    | 0.056 d                 | 0.633 b                                                    | 3.28 c                         | 1.03 c |

|                      |          |        |        |          |         |        |        |
|----------------------|----------|--------|--------|----------|---------|--------|--------|
| T3                   | 19.16 b  | 2.99 a | 1.12 a | 0.097 b  | 0.719 a | 4.55 b | 1.5 a  |
| T4                   | 19.97 b  | 3.04 a | 1.15 a | 0.118 a  | 0.726 a | 5.85 a | 1.5 a  |
| T5                   | 14.74 c  | 2.85 a | 0.85 b | 0.079 c  | 0.725 a | 4.50 b | 1.3 b  |
| T6                   | 10.94 d  | 2.64 b | 0.69 c | 0.066 cd | 0.731 a | 4.58 b | 1.5 a  |
| F-test               | **       | **     | **     | **       | *       | *      | *      |
| Second season (2024) |          |        |        |          |         |        |        |
| T1                   | 23.14 a  | 3.34 a | 1.17 a | 0.125 a  | 0.742 a | 4.66 c | 1.5 b  |
| T2                   | 7.10 e   | 1.79 b | 0.45 d | 0.049 d  | 0.642 b | 2.77 e | 1.1 c  |
| T3                   | 18.55 bc | 3.21 a | 1.15 a | 0.119 a  | 0.734 a | 4.87 b | 1.6 ab |
| T4                   | 21.15 a  | 3.25 a | 1.24 a | 0.122 a  | 0.735 a | 5.64 a | 1.5 a  |
| T5                   | 16.45 c  | 3.06 a | 1.02 b | 0.085 b  | 0.731 a | 5.15 a | 1.4 a  |
| T6                   | 13.61 d  | 3.01 a | 0.86 c | 0.074 c  | 0.728 a | 4.82 d | 1.4 c  |
| F-test               | **       | *      | *      | **       | *       | *      | **     |

307 \* and \*\* indicate significant and highly significant, respectively

308 Values of means in each column followed by the same letter are not significantly at  $p < 0.05$

309 Where, T1: negative control (healthy seedlings), T2: control (positive control), T3: fungicide, T4: n-SiO<sub>2</sub> (200 mg L<sup>-1</sup>), T5: n-Se (100 mg L<sup>-1</sup>), and  
310 T6: ½ (n-Se + n-SiO<sub>2</sub>) (50 + 100 mg L<sup>-1</sup>, respectively)




311


312

313

### 314 3.6. Antioxidant Activity and Electrolyte Leakage of pepper seedlings under biotic stress

315 The activities of three antioxidant enzymes including CAT, POX, and PPO were measured in pepper seedlings  
316 leaves 40 days after sowing, along with the percentage of electrolyte leakage under different treatments in both  
317 seasons (**Figure 6**). Generally, the highest enzyme activities were recorded with the application of n-SiO<sub>2</sub>  
318 (200 mg L<sup>-1</sup>), followed by the combined n-Se (50 mg L<sup>-1</sup>) and n-SiO<sub>2</sub> (100 mg L<sup>-1</sup>), and both n-Se (100 mg L<sup>-1</sup>)  
319 and chemical fungicide applications, whereas the lowest enzyme activities were recorded in the positive  
320 control (T2). In contrast, the infected seedlings (T2) had the highest electrolyte leakage values followed by  
321 treatments T5, T6 and T3, The n-SiO<sub>2</sub> treatment at 200 mg L<sup>-1</sup> showed the lowest electrolyte leakage in both  
322 seasons. Best (lowest) values of electrolyte leakage and antioxidants were obtained with n-SiO<sub>2</sub> (200 mg L<sup>-1</sup>)  
323 as confirmed by the earlier assessing enzymes, followed by the negative control and the combined n-Se (50  
324 mg L<sup>-1</sup>) and n-SiO<sub>2</sub> (100 mg L<sup>-1</sup>). It is obvious that n-silica can enhance the producing of enzymes in 3-5 folds  
325 compared to the controls.





**Figure 6.** Total antioxidant activity including (A) peroxidase (POX), (B) catalase (CAT), (C) polyphenol oxidase (PPO), and (D) electrolyte leakage percentage measured 40 days after sowing during the both growing seasons. Results are given as mean values  $\pm$  standard error bars

Where, T1: negative control (healthy seedlings), T2: control (positive control), T3: fungicide, T4: n-SiO<sub>2</sub> (200 mg L<sup>-1</sup>), T5: n-Se (100 mg L<sup>-1</sup>), and T6:  $\frac{1}{2}$  (n-Se + n-SiO<sub>2</sub>) (50 + 100 mg L<sup>-1</sup>, respectively)

#### 4. Discussion

The quality of vegetable seedlings a critical factor in vegetable production, which mainly depends on the growing conditions. Effective management of seedling production requires several parameters should be kept in mind including the growing conditions, abiotic/biotic stress and cultivation facilities. Therefore, the present study was carried out to examine the potential of applied nanoparticles (nano selenium and nano silica) to boost pepper seedlings production and to control the root rot disease caused by *Rhizoctonia solani*. This work also explores the individual and combined effects of these nanoparticles in combating the pathogen during the production of pepper seedlings. To what extent can the applied NPs be effective for inhibiting studied pathogen, as compared to traditional fungicides? Concerning the most effective applied NPs (individually or in a combination), *in vitro* and under greenhouse conditions were achieved.

Seedling growth and quality need a specific handling in the nursery of vegetables. There is an urgent need concerning the role of environmental factors and their impact the growth of pepper seedlings. To produce high-quality seedlings, various factors must be considered, particularly environmental stresses such as light intensity, temperature, nutrients availability and water), and biotic stress (Islam et al. 2024). Several fungal diseases can cause root rots, damping-off, and wilting in pepper seedlings. Numerous reports discussed different fungal phytopathogens causing root rot in pepper seedlings including *Rhizoctonia* spp. (Mannai et al. 2018; El-Kazzaz et al. 2022), and Oomycetes such as *Pythium* spp. (Ao et al. 2024), and *Fusarium* spp. (Hafez et al. 2025). **Table 6** summarizes important studies worldwide on fungal, bacterial, and viral infections af-

fecting pepper plants. Due to susceptibility of pepper to many diseases; the control strategy of such pathogens mainly depends on the pathogen species (e.g., virus, bacteria, fungi, and nematode). Among various control strategies, nanomaterials are considered a promising agent against biotic stress besides the biological controls such as beneficial microorganisms (mainly Tricoderma, Arbuscular mycorrhizal fungi species), and chitosan (Table 6).

**Table 6.** A survey on the published studied on pepper diseases infected by bacterial, fungal, and viral pathogens along with the possible control methods

| Country  | Pepper variety                                   | Studied pathogen                                                                                          | The common disease                                                | Control strategy                                                                  | Refs.                           |
|----------|--------------------------------------------------|-----------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------|-----------------------------------------------------------------------------------|---------------------------------|
| Egypt    | Pepper seedlings<br>( <i>Capsicum annuum</i> L.) | <i>Rhizoctonia solani</i> and<br><i>Fusarium oxysporum</i>                                                | Pepper root rot and<br>wilt diseases                              | <i>Paenibacillus polymyxa</i><br>and <i>Trichoderma</i><br><i>longibrachiatum</i> | El-Kazzaz et al.<br>(2022)      |
| Egypt    | Sweet pepper (cv.<br>Golden plants)              | Fungus ( <i>Fusarium</i><br><i>oxysporum</i> ) and<br>nematode ( <i>Meloidogyne</i><br><i>incognita</i> ) | Root-knot nematodes<br>& root rot fungus                          | Bio-ZnO-NPs &<br><i>Trichoderma</i><br><i>longibrachiatum</i>                     | Ghareeb et al.<br>(2024)        |
| Egypt    | Pepper seedlings                                 | Fungus ( <i>Fusarium</i><br><i>Oxysporum</i> )                                                            | Wilt disease of pepper<br>plants                                  | Green Ag-NPs by<br><i>Aspergillus Ustus</i><br>ON076464                           | Yahia et al.<br>(2023)          |
| India    | Pepper chilli (var.<br>Arka Lohith)              | Fungus ( <i>Colletotrichum</i><br><i>truncatum</i> )                                                      | Chilli anthracnose<br>disease                                     | Chitosan-silver<br>nanocomposite<br>(CS-AgNPs)                                    | Gowda et al.<br>(2023)          |
| India    | Pepper chilli                                    | Fungus ( <i>Colletotrichum</i><br><i>capsici</i> )                                                        | Chilli fruit rot disease                                          | Green copper<br>nanoparticles                                                     | Iliger et al.<br>(2021)         |
| Pakistan | Pepper chili<br>( <i>Capsicum annuum</i><br>L.)  | Fungus ( <i>Colletotrichum</i><br><i>capsici</i> )                                                        | Chilli fruit rot disease                                          | Green silver nanoparticles                                                        | Nawaz et al.<br>(2024)          |
| India    | Pepper chili<br>( <i>Capsicum annuum</i><br>L.)  | Fungi ( <i>Phomopsis</i><br><i>vexans</i> and<br><i>Colletotrichum capsici</i> )                          | Blight and fruit rot in<br>chili                                  | <i>Trichoderma</i> -mediated<br>ZnO-NPs                                           | Kumar et al.<br>(2024)          |
| Mexico   | Varieties of<br>jalapeño pepper                  | Begomovirus                                                                                               | Pepper huasteco<br>yellow vein virus                              | Chemical ZnO-NPs                                                                  | Rivero-Montejo<br>et al. (2023) |
| Mexico   | Pepper ( <i>Capsicum</i><br><i>annuum</i> L.)    | Virus disease                                                                                             | Tobacco mosaic virus                                              | Titanium dioxide<br>nanoparticles (TiO <sub>2</sub> -NPs)                         | Acuña-Fuentes<br>et al. (2022)  |
| Egypt    | Sweet pepper (cv.<br>Sirtaki)                    | Tobamovirus                                                                                               | Pepper mild mottle<br>virus                                       | Chemical Ag-NPs                                                                   | Elbeshehy et al.<br>(2023)      |
| Pakistan | Bell pepper<br>( <i>Capsicum annuum</i><br>L.)   | Bacterial leaf spot                                                                                       | <i>Xanthomonas</i><br><i>campestris</i> pv.<br><i>vesicatoria</i> | Green Cu-Zn hybrid-NPs                                                            | Ali et al. (2024)               |
| Pakistan | Chilli pepper<br>( <i>Capsicum annuum</i><br>L.) | Bacterial leaf spot                                                                                       | <i>Xanthomonas</i><br><i>campestris</i> pv.<br><i>vesicatoria</i> | Green Ag-NPs and<br>Zn-NPs                                                        | Atiq et al.<br>(2023)           |
| Turkey   | Pepper variety                                   | <i>Fusarium solani</i> and <i>F.</i>                                                                      | Root and crown rot                                                | Arbuscular mycorrhizal                                                            | Bilgili (2025)                  |

|       |                  |                                                                                               |                 |                                                  |                    |
|-------|------------------|-----------------------------------------------------------------------------------------------|-----------------|--------------------------------------------------|--------------------|
|       | INAN-3363 F1     | mix ( <i>F. solani</i> , <i>F. oxysporum</i> ; <i>F. oxysporum</i> f.sp. <i>vasinfectum</i> ) | pathogens       | fungi species                                    |                    |
| China | Pepper seedlings | Fungus ( <i>Fusarium solani</i> )                                                             | Pepper root rot | <i>Bacillus subtilis</i> PTS-394 wettable powder | Qiao et al. (2023) |
| China | Pepper seedlings | Fungus ( <i>Fusarium solani</i> )                                                             | Pepper root rot | <i>Bacillus velezensis</i> Yao                   | Pei et al. (2023)  |

As far as we know, this study is first report about using nanoparticles of selenium and silica to control root rot disease of pepper seedlings caused by *Rhizoctonia solani*. This section explains our choice of nanoparticles and the proposed mechanism. A variety of nanomaterials have previously been used against phytopathogens of pepper seedlings such as ZnO-NPs, Ag-NPs, CuO-NPs, TiO<sub>2</sub>-NPs, Cu-Zn hybrid-NPs, and Zn-NPs. On the other hand, many pepper diseases were reported to be controlled by applying such nanomaterials or nanoparticles depending on microbial species such as virus, bacteria, fungi, and nematode (Table 6). Among fungal phytopathogens, *Rhizoctonia solani* is a devastating phytopathogen causing pepper root rot disease (Ao et al., 2024).

The selected NPs (nano selenium and nano silica) have well-documented applications under abiotic and biotic stress due to their unique chemical, physical, and biological properties (Díaz-Parra et al. 2025). Numerous studies confirmed the potential of selected Se-NPs and SiO<sub>2</sub>-NPs under biotic stress individually (e.g., Bhat et al. 2021; Goswami et al. 2022; Naidu et al. 2023; Mahawar et al. 2023; Yan et al. 2024; Qin et al. 2025), whereas a few in combined application (Taha et al. 2023). The reasons behind selecting nanoparticles of selenium and silica because their well-known role in controlling biotic stress on plants through their inducing plant defense system and systemic resistance, activating the antioxidants system, and enhancing the plant growth of stressful plants (Qin et al. 2025; Díaz-Parra et al. 2025). As far as we know, this is first report on using nano—silica and nano-Se to control the pathogen *R. solani* in pepper seedlings.

Our current findings confirmed that nano-silica (n-SiO<sub>2</sub>) was the most effective treatment against *Rhizoctonia solani* (Table 3). *In vitro*, n-SiO<sub>2</sub> achieved an 81.1% reduction in mycelial growth compared to other NPs. Concerning the survival seedlings percentage under greenhouse conditions, there is no significant difference between chemical fungicide (82.3 and 79.4%) and n-SiO<sub>2</sub> (79.0 and 75.8%) in both seasons, respectively. These results suggested that n-SiO<sub>2</sub> at 200 mg L<sup>-1</sup> is a promising alternative to chemical fungicide (Rizolex 50% WP) for controlling *Rhizoctonia solani* in pepper seedlings. Similar results were reported by Awad-Allah et al. (2021) who recognized the effect of n-SiO<sub>2</sub> at 150 mg L<sup>-1</sup> against bacterial leaf spot (*Xanthomonas vesicatoria*) due to the stimulating of n-SiO<sub>2</sub> to release salicylic acid which activates the plant defense and immune system along with protecting plants against pathogen attacks. In this concern, it is re-

ported that n-silica improved the biochemical, physiological response and growth in plants against stress conditions (Hasan et al. 2024).

Furthermore, n-silica (200 mg L<sup>-1</sup>) treatment induced the highest values of vegetative growth parameters and the enzymatic antioxidants (CAT, POX, PPO) in pepper seedlings. It is found that applied n-silica (150 mg L<sup>-1</sup>) induced pepper resistance to bacterial pathogen by stimulating activity of PPO (Awad-Allah et al. 2021). These results were also confirmed on infected plants with *R. solani* after application of n-silica (100 mg L<sup>-1</sup>) on wheat plants (*Triticum aestivum* L.) by Abdelrhim et al. (2021), on carrot (*Daucus carota* L.) by Ahamad and Siddiqui (2021), on oat (*Avena sativa* L.) by Ahmad et al. (2023), and on rice (*Oryza sativa* L.) by Tan et al. (2025). Why applied n-silica record the best results in controlling *R. solani* in the current study? To answer the previous question, three levels of the explanations are being needed including the vegetative growth of pepper seedlings and the biochemical measurements of enzymatic antioxidants along with the photosynthetic parameters.

Concerning the vegetative growth parameters of pepper seedlings, all studied vegetative parameters Stem diameter, seedling height, fresh and dry weights of seedlings were produced from n-silica treatment (Table 5). The suggested mechanism of n-silica to strengthen the vegetative parameters against the pathogen may go back to the deposition of n-SiO<sub>2</sub> in pepper root tissues, which improved plant defense against the studied pathogen as confirmed by the SEM investigation (**Figure 4**). The suggested defense mechanisms may go back to (1) forming physical barriers by accumulation of Si in cell walls and below the cuticle, preventing the infection pegs of *R. solani* appressoria (Ahammed and Yang 2021), and (2) forming Si-enriched layer through uniform distribution of Si-aggregates and preventing the fungal ingress (Abdelrhim et al. 2021).

Along with the vegetative parameters of pepper seedlings, the highest values of chlorophyll fluorescence, Chl. a, and Chl. b were achieved with n-silica application. Under *R. solani* stress, it is expected reducing the photosynthesis process through inhibiting photosynthetic activities by causing damage to the structural along with stomatal closures of the seedlings (Taha et al. 2023). N-silica is known to protect chloroplasts from oxidative damage, increase chlorophyll content, and act as a cofactor in many enzymatic reactions (Zahedi et al. 2020; Naidu et al. 2023). This enhanced efficiency of photosynthesis likely contributes to strong seedlings (Ahamad and Siddiqui 2021).

Regarding the enzymatic antioxidants, pepper seedlings treated with n-silica showed the highest activities of antioxidant enzymes (CAT, POX, and PPO) compared to other treatments. This result is in agreement with the suggested defense mechanism, which confirm the role of Si in producing defense non-enzymatic compounds (i.e., anthocyanins, flavonoids and phenolics) and enzymatic activities leading to the development of physical barriers against phyto-pathogens (Ahanger et al. 2020). The opposite behavior can be noticed for the electrolyte leakage, which recorded the lowest values upon n-silica compared to control and other treatments.

## 5. Conclusions

This study presents noval insight into alternatives to chemical fungicides through the use of nanofungicides. As far as we know this is first report on applying n-silica and n-Se in controlling *R. solani* on pepper seedlings. Among the treatments n-silica at 200 mg L<sup>-1</sup> recorded the highest values in all studied vegetative parameters, photosynthetic attributes, and antioxidants enzyme activities (CAT, POX, PPO), as compared to other treatments. It could be recommended that application of n-silica (200 mg L<sup>-1</sup>) is a promising and effective alternative to the chemical fungicide (Rizolex 50% WP) against root rot disease caused by *R. solani*. Moreover, nano-silica is better than n-Se and/or the combined of n-Se + n-silica. While current findings enhance our understanding of the antifungal potential of n-Se and/or n-silica nanofungicides, further research is warranted to explore their interactions and efficacy across different crops before broad agriculture application.

**CRedit authorship contribution statement:** Conceptualization and visualization, Y.A.B. and N.A.T.; methodology, N.A.T. and Y.A.B.; software, M.S.A, A.A.K.; validation, M.S.A. and S. M. A.; formal analysis, T.A.S., H.S.E. and Y.A.B.; investigation, N.A.T.; resources, H.E.R.; data curation, Y.A.B.; writing—original draft preparation, H.E.R. and T.A.S.; writing, review and editing, all authors; visualization, A.A.K, S.H. and H.E.R.; supervision, Y.A.B. and T.A.S.; project administration, H.E.R. All authors have read and agreed to the published version of the manuscript.

**Acknowledgments:** Authors extend their gratefulness to Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research, King Faisal University, Saudi Arabia, for supporting this research work through the grant number (KFU252104).

**Funding:** Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research, King Faisal University, Saudi Arabia, Grant number (KFU252104)

**Declaration of competing interest:** Authors declare no conflict of interest.

**Data Availability Statement:** Available on request.

## References

- Abdelrhim, A.S., Mazrou, Y.S.A., Nehela, Y., Atallah, O.O., El-Ashmony, R.M., Dawood, M.F.A. (2021). Silicon Dioxide Nanoparticles Induce Innate Immune Responses and Activate Antioxidant Machinery in Wheat Against *Rhizoctonia solani*. *Plants* 10, 2758. <https://doi.org/10.3390/plants10122758>
- Acuña-Fuentes, N.L., Vargas-Hernandez, M., Rivero-Montejo, S.d.J., Rivas-Ramirez, L.K., Macias-Bobadilla, I., Palos-Barba, V., Rivera-Muñoz, E.M., Guevara-Gonzalez, R.G., Torres-Pacheco, I. (2022). Antiviral Activity of TiO<sub>2</sub> NPs against Tobacco Mosaic Virus in Chili Pepper (*Capsicum annuum* L.). *Agriculture* 12, 2101. <https://doi.org/10.3390/agriculture12122101>
- Aebi, H. (1984). Catalase *in vitro*. In *Methods in Enzymology*; Packer, L., Ed.; Academic Press: San Diego, CA, USA, Volume 105, pp. 121–126.

465 Ahamad, L., Siddiqui, Z.A. (2021). Effects of Silicon Dioxide, Zinc Oxide and Titanium Dioxide Nanopar-  
 466 ticles on *Meloidogyne incognita*, *Alternaria dauci* and *Rhizoctonia solani* Disease Complex of Carrot.  
 467 *Experimental Parasitology* 230, 108176. <https://doi.org/10.1016/j.exppara.2021.108176>

468 Ahammed, G.J., Yang, Y. (2021). Mechanisms of Silicon-Induced Fungal Disease Resistance in Plants.  
 469 *Plant Physiology and Biochemistry* 165, 200–206. <https://doi.org/10.1016/j.plaphy.2021.05.031>

470 Ahanger, M.A., Bhat, J.A., Siddiqui, M.H., Rinklebe, J., Ahmad, P. (2020). Integration of Silicon and Sec-  
 471 ondary Metabolites in Plants: A Significant Association in Stress Tolerance. *Journal of Experimental*  
 472 *Botany* 71, 6758–6774. <https://doi.org/10.1093/jxb/eraa291>

473 Ahmad, F., Jabeen, K., Iqbal, S., Umar, A., Ameen, F., Gancarz, M., Darwish, D.B.E (2023). Influence of  
 474 Silicon Nanoparticles on *Avena sativa* L. to Alleviate the Biotic Stress of *Rhizoctonia solani*. *Scientific*  
 475 *Reports* 13, 15191. <https://doi.org/10.1038/s41598-023-41699-w>

476 Ali, F., Atiq, M., Rajput, N.A (2024). Intervention of Bacterial Leaf Spot of Bell Pepper through Neem  
 477 Mediated Copper and Zinc Hybrid Nanoparticles. *Phytopathogenomics and Disease Control* 3, 251–  
 478 259. <https://doi.org/10.22194/Pdc/3.1043>

479 Altaf, M.A., Hao, Y., He, C., Mumtaz, M.A., Shu, H., Fu, H., Wang, Z (2022). Physiological and Bio-  
 480 chemical Responses of Pepper (*Capsicum annuum* L.) Seedlings to Nickel Toxicity. *Frontiers in Plant*  
 481 *Science* 13, 950392. <https://doi.org/10.3389/fpls.2022.950392>

482 Altier, N.A., Judy, A.T (1995). Identification of Resistance to *Pythium* Seedling Diseases in Alfalfa Using a  
 483 Culture Plate Method. *Plant Disease* 79, 341–346.

484 Ao, N., Zou, H., Li, J., Shao, H., Kageyama, K., Feng, W (2024). First Report of *Pythium aphanidermatum*  
 485 and *Pythium myriotylum* Causing Root Rot on Chili Pepper (*Capsicum annuum* L.) in Guizhou, China.  
 486 *Crop Protection* 181, 106704. <https://doi.org/10.1016/j.cropro.2024.106704>

487 Atiq, M.; Fatima, T., Rajput, N.A., Usman, M., Kachelo, G.A., Ahmad, U., Arif, A.M., Nawaz, A., Kashif,  
 488 M., Ashraf, M (2023). *Moringa oleifera* Plant Based Copper and Silver Nanoparticles and Its Anti-  
 489 bacterial Activity towards Leaf Spot of Chilli Caused by *Xanthomonas campestris* pv. *vesicatoria*.  
 490 *International Journal of Phytopathology* 12, 63–72.

491 Awad-Allah, E.F.A., Shams, A.H.M., Helaly, A.A (2021). Suppression of Bacterial Leaf Spot by Green  
 492 Synthesized Silica Nanoparticles and Antagonistic Yeast Improves Growth, Productivity and Quality  
 493 of Sweet Pepper. *Plants* 10, 1689. <https://doi.org/10.3390/plants10081689>

494 Badawy, S.A., Zayed, B.A., Bassiouni, S.M.A., Mahdi, A.H.A., Majrashi, A., Ali, E.F., Seleiman, M.F  
 495 (2021). Influence of Nano Silicon and Nano Selenium on Root Characters, Growth, Ion Selectivity,  
 496 Yield, and Yield Components of Rice (*Oryza sativa* L.) under Salinity Conditions. *Plants* 10, 1657.  
 497 <https://doi.org/10.3390/plants10081657>

498 Batista, A., Mai, V.C., Sadowska, K., et al. (2025). Application of Silver and Selenium Nanoparticles to  
 499 Enhance Plant-Defense Response against Biotic Stressors. *Acta Physiologiae Plantarum* 47, 21.  
 500 <https://doi.org/10.1007/s11738-025-03768-7>

501 Bhat, J.A., Rajora, N., Raturi, G., Sharma, S., Dhiman, P., Sanand, S., Shivaraj, S.M., Sonah, H., Deshmukh,  
 502 R (2021). Silicon Nanoparticles (SiNPs) in Sustainable Agriculture: Major Emphasis on the Practi-  
 503 cality, Efficacy and Concerns. *Nanoscale Advances* 3, 4019–4028.  
 504 <https://doi.org/10.1039/d1na00233c>

505 Bilgili, A (2025). The Effectiveness of Arbuscular Mycorrhizal Fungal Species (*Funnelformis mosseae*,  
 506 *Rhizophagus intraradices*, and *Claroideoglomus etunicatum*) in the Biocontrol of Root and Crown Rot  
 507 Pathogens, *Fusarium solani* and *Fusarium* Mixture in Pepper. *PeerJ* 13, e18438.  
 508 <https://doi.org/10.7717/peerj.18438>

509 Chandler, J.M.; Santelman, P.W (1968). Interaction of Four Herbicides with *Rhizoctonia solani* on Seedling  
 510 Cotton. *Weed Science* 16, 453–454.

511 Chethan, D., Chandana, R., Khan, F., Kavya, B.S., Poonacha, T.T., Karan, R., Bevanur, A., Kruthika, R.,  
 512 Ashwini, K.S., Vani, Y.; et al. (2025). A Novel Report of *Rhizoctonia solani* AG-4 HG-I Associated  
 513 with Damping-Off and Wilt of Quinoa (*Chenopodium quinoa*) in India. *Physiological and Molecular*  
 514 *Plant Pathology* 102725. <https://doi.org/10.1016/j.pmpp.2025.102725>

515 Choudhary, D., Deshmukh, S., Maheswari, G., Kumari, A., Ghormade, V (2025). Silica and Mesoporous  
 516 Silica Nanoparticles Display Effective Insecticidal Effect and Augment Plant Defense Respons-  
 517 es. *Pesticide Biochemistry and Physiology* 210, 106389. <https://doi.org/10.1016/j.pestbp.2025.106389>

518 Dhingra, O.D., Sinclair, J.B (1985). *Culture of Pathogens. Basic Plant Pathology Methods*; CRC Press Inc.:  
 519 Boca Raton, FL, USA, pp. 11–47.

520 Díaz-Parra, D.G., García-Casillas, L.A., Velasco-Ramírez, S.F., Guevara-Martínez, S.J., Zamudio-Ojeda, A.,  
 521 Zuñiga-Mayo, V.M., Rodríguez-Guzmán, E., Melchor-González, A., Lomelí-Rosales, D.A., Veláz-  
 522 quez-Juárez, G (2025). Role of Metal-Based Nanoparticles in *Capsicum* spp. Plants. *ACS Omega* 10,  
 523 10756–10768. <https://doi.org/10.1021/acsomega.4c11102>

524 Elbeshehy, E.K.F., Hassan, W.M., Baeshen, A.A (2023). Controlling Pepper Mild Mottle Virus (PMMoV)  
 525 Infection in Pepper Seedlings by Use of Chemically Synthetic Silver Nanoparticles. *Molecules* 28,  
 526 139. <https://doi.org/10.3390/molecules28010139>

527 El-Kazzaz, M.K., Ghoneim, K.E., Agha, M.K.M., Helmy, A., Behiry, S.I., Abdelkhalek, A., Saleem, M.H.,  
 528 Al-Askar, A.A., Arishi, A.A., Elsharkawy, M.M (2022). Suppression of Pepper Root Rot and Wilt  
 529 Diseases Caused by *Rhizoctonia solani* and *Fusarium oxysporum*. *Life* 12, 587.  
 530 <https://doi.org/10.3390/life12040587>

531 FAOSTAT (2025). *FAOSTAT Statistics Database*. Available online: <http://www.fao.org/faostat> (accessed on  
 532 20 April 2025).

533 Ferreira, J.H., Mathee, F.N., Thomas, A.C (1991). Biological Control of *Eutypalota* on Grapevine by an  
 534 Antagonistic Strain of *Bacillus subtilis*. *Phytopathology* 81, 283–287.

535 Gao, J., Fu, J.X., Jiao, J., Gai, Q.Y., Zhang, Z.Y., Wang, X.Q., Fu, Y.J (2025). An Endophytic Fun-  
 536 gus *Dichotomopilus funicola* J-219 for the Control of Pigeon Pea Root Rot Caused by *Rhizoctonia*  
 537 *solani* and Its Role in Regulating the Secondary Metabolic Defense Response. *Physiological and*  
 538 *Molecular Plant Pathology* 137, 102613. <https://doi.org/10.1016/j.pmpp.2025.102613>

539 Gao, S., Tuda, M (2024). Silica and Selenium Nanoparticles Attract or Repel Scale Insects by Altering  
 540 Physicochemical Leaf Traits. *Plants* 13, 952. <https://doi.org/10.3390/plants13070952>

541 Ghareeb, Y.R., Belal, E.B., El-Khateeb, N.M.M., Shreef, B.A (2024). Utilizing Bio-Synthesis of Nano-  
 542 materials as Biological Agents for Controlling Soil-Borne Diseases in Pepper Plants: Root-Knot  
 543 Nematodes and Root Rot Fungus. *BMC Plant Biology* 24, 110.  
 544 <https://doi.org/10.1186/s12870-024-04760-y>

545 Ghazi, A.A., El-Nahrawy, S., El-Ramady, H., Ling, W (2022). Biosynthesis of Nano-Selenium and Its Im-  
 546 pact on Germination of Wheat under Salt Stress for Sustainable Production. *Sustainability* 14, 1784.

547 González-Hernández, A.I., Suárez-Fernández, M.B., Pérez-Sánchez, R., Gómez-Sánchez, M.Á., Mo-  
 548 rales-Corts, M.R (2021). Compost Tea Induces Growth and Resistance against *Rhizoctonia solani* and  
 549 *Phytophthora capsici* in Pepper. *Agronomy* 11, 781. <https://doi.org/10.3390/agronomy11040781>

550 Goswami, P., Mathur, J., Srivastava, N (2022). Silica Nanoparticles as Novel Sustainable Approach for Plant  
 551 Growth and Crop Protection. *Heliyon* 8, e09908. <https://doi.org/10.1016/j.heliyon.2022.e09908>

552 Gowda, S., Sriram, S (2023). Green Synthesis of Chitosan Silver Nanocomposites and Their Antifungal  
 553 Activity against *Colletotrichum truncatum* Causing Anthracnose in Chillies. *Plant Nano Biology* 5,  
 554 100041. <https://doi.org/10.1016/j.plana.2023.100041>

555 Hafez, M., Rashad, Y.M., Ibrahim, O.M., El-Sharkawy, H.A., Abd-ElGawad, A.M., Bourouah, M., Abdalla,  
 556 S.A., Sleem, M.M., Madbouly, A.K (2025). Optimization of Antifungal and Nitrogen-Fixing Activi-  
 557 ties of *Niallia circulans* YRNF1 Using Response Surface Methodology to Control *Fusarium* Root Rot  
 558 of Pepper and Promote the Plant Growth. *Rhizosphere* 33, 101049.  
 559 <https://doi.org/10.1016/j.rhisph.2025.101049>

560 Hasan, M.K., Shopan, J., Jahan, I., Suravi, T.I (2024). Silicon Nanomaterials Enhance Seedling Growth and  
 561 Plant Adaptation to Acidic Soil by Promoting Photosynthesis and Antioxidant Activity in Mustard  
 562 (*Brassica campestris* L.). *International Journal of Molecular Sciences* 25, 10318.  
 563 <https://doi.org/10.3390/ijms251910318>

564 He, J., Li, J., Gao, Y., He, X., Hao, G (2023). Nano-based smart formulations: A potential solution to the  
 565 hazardous effects of pesticide on the environment. *J. Hazard. Mater.* 456, 131599.  
 566 <https://doi.org/10.1016/j.jhazmat.2023.131599>.

567 Hernández-Huerta, J., Tamez-Guerra, P., Gomez-Flores, R., Delgado-Gardea, M.C.E., Robles-Hernández,  
568 L., Gonzalez-Franco, A.C., Infante-Ramirez, R (2023). Pepper growth promotion and biocontrol  
569 against *Xanthomonas euvesicatoria* by *Bacillus cereus* and *Bacillus thuringiensis* formulations. *PeerJ*  
570 11, e14633. <https://doi.org/10.7717/peerj.14633>.

571 Hoffmann, A.B., Poorter, H (2002). Avoiding bias in calculations of relative growth rate. *Ann. Bot.* 80, 37–  
572 42. <https://doi.org/10.1093/aob/mcf140>.

573 Huang, S.; Yang, X.; Zhao, J.; Zeeshan, M.; Wang, C.; Yang, D.; Han, X.; Zhang, G. Antifungal activity and  
574 potential inhibition mechanism of 4-(diethylamino)salicylaldehyde against *Rhizoctonia solani*. *Pestic.*  
575 *Biochem. Physiol.* **2025**, \*212\*, 106444. <https://doi.org/10.1016/j.pestbp.2025.106444>.

576 Illiger, K.S., Sofi, T.A., Bhat, N.A., Ahanger, F.A., Sekhar, J.C., Elhendi, A.Z., Al-Huqail, A.A., Khan, F  
577 (2021). Copper nanoparticles: Green synthesis and managing fruit rot disease of chilli caused by  
578 *Colletotrichum capsici*. *Saudi J. Biol. Sci.* 28, 1477–1486. <https://doi.org/10.1016/j.sjbs.2020.12.003>.

579 Imran, M., Umer, M., Raja, N.I., Abasi, F., Sardar, N., Rahman, U., Naqvi, S.A.M., Baloch, M.Y.J., Alrefaei,  
580 A.F (2025). Antibacterial potential of silver–selenium nanocomposites in mitigating fire blight disease  
581 in *Pyrus communis* L. *Front. Plant Sci.* 16, 1541498. <https://doi.org/10.3389/fpls.2025.154149>.

582 Islam, S., Reza, M.N., Ahmed, S., Samsuzzaman; Cho, Y.J., Noh, D.H., Chung, S.-O (2024). Seedling  
583 Growth Stress Quantification Based on Environmental Factors Using Sensor Fusion and Image Pro-  
584 cessing. *Horticulturae* 10, 186. <https://doi.org/10.3390/horticulturae10020186>.

585 Jang, Y., Schafleitner, R., Barchenger, D.W., Lin, Y., Lee, J (2025). Evaluation of heat stress response in  
586 pepper (*Capsicum annuum* L.) seedlings under controlled environmental conditions using a  
587 high-throughput 3D multispectral phenotyping. *Sci. Hortic.* 345, 114136.  
588 <https://doi.org/10.1016/j.scienta.2025.114136>.

589 Kang, L., Wu, Y., Jia, Y., Zhang, X., Li, H., Wang, J., Sun, Y., Chen, R., Liu, M., Yang, Q (2023).  
590 Nano-Selenium Enhances Melon Resistance to *Podosphaera xanthii* by Enhancing the Antioxidant  
591 Capacity and Promoting Alterations in the Polyamine, Phenylpropanoid and Hormone Signaling  
592 Pathways. *Journal of Nanobiotechnology* 21, 377. <https://doi.org/10.1186/s12951-023-02148-y>.

593 Kashyap, A.S., Manzar, N., Vishwakarma, S.K., Mahajan, C., Dey, U (2024). Tiny but mighty: metal na-  
594 noparticles as effective antimicrobial agents for plant pathogen control. *World J. Microbiol. Bio-*  
595 *technol.* 40, 104. <https://doi.org/10.1007/s11274-024-03911-5>.

596 Kirupanatha-Rajan, P., Chanthini, K.M.P., Deepthi, R., Manju, R., Pavithra, G.S., Murugan, P., De-  
597 va-Andrews, A., Senthil-Nathan, S (2025). Enhanced secondary metabolite accumulation in tomato  
598 roots mediated by colonization of *Paenibacillus dendritiformis* confers crop protection against foot rot  
599 caused by *Rhizoctonia solani*. *Physiol. Mol. Plant Pathol.* 102721.  
600 <https://doi.org/10.1016/j.pmpp.2025.102721>.

601 Kumar, R., Singh, S.P., Ivanov, F., Maksimov, A.Y., Latsynnik, E., Minkina, T., Keswani, C (2024).  
602 *Trichoderma*-Mediated Synthesis of ZnONPs: Trend Efficient for Repressing Two Seed- and  
603 Soil-Borne Phytopathogens *Phomopsis vexans* and *Colletotrichum capsici*. *Bionanoscience* 14, 5363.  
604 <https://doi.org/10.1007/s12668-024-01495-w>.

605 Leslie, J.F., Summerell, B.A (2006). *The Fusarium Laboratory Manual*; Blackwell Publishing: Hoboken, NJ,  
606 USA, 1–2. <https://doi.org/10.1002/9780470278376>.

607 Li, N., Pu, K., Ding, D., Yang, Y., Niu, T., Li, J., Xie, J (2023). Foliar Spraying of Glycine Betaine Alleviated  
608 Growth Inhibition, Photoinhibition, and Oxidative Stress in Pepper (*Capsicum annuum* L.) Seedlings  
609 under Low Temperatures Combined with Low Light. *Plants* 12, 2563.  
610 <https://doi.org/10.3390/plants12132563>.

611 Lu, X., Wu, Q., Nie, K., Wu, H., Chen, G., Wang, J., Ma, Z (2023). Exogenous phthalanilic acid induces  
612 resistance to drought stress in pepper seedlings (*Capsicum annuum* L.). *Front. Plant Sci.* 14, 1156276.  
613 <https://doi.org/10.3389/fpls.2023.1156276>.

614 Lutts, S., Kinet, J.M., Bouharmont, J (1996). NaCl-induced senescence in leaves of rice (*Oryza sativa* L.)  
615 cultivars differing in salinity resistance. *Ann. Bot.* 78, 389–398.

616 Mahawar, L., Ramasamy, K.P., Suhel, M., Prasad, S.M., Živčák, M., Brestic, M., Rastogi, A., Skalický, M  
617 (2023). Silicon nanoparticles: Comprehensive review on biogenic synthesis and applications in agri-  
618 culture. *Environ. Res.* 232, 116292. <https://doi.org/10.1016/j.envres.2023.116292>.

619 Malik, C.P., Singh, M.B (1980). *Plant Enzymology and Histo-Enzymology: A Text Manual*; Kalyani Publi-  
620 cations: New Delhi, India, p. 434.

621 Mannai, S., Jabnoun-Khiareddine, H., Nasraoui, B., Daami-Remadi, M (2018). *Rhizoctonia* Root Rot of  
622 Pepper (*Capsicum annuum*): Comparative Pathogenicity of Causal Agent and Biocontrol Attempt  
623 using Fungal and Bacterial Agents. *J. Plant Pathol. Microbiol.* 9, 431.  
624 <https://doi.org/10.4172/2157-7471.1000431>.

625 Maxwell, K., Johnson, G.N (2000). Chlorophyll Fluorescence—A Practical Guide. *J. Exp. Bot.* 51, 659–668.  
626 <https://doi.org/10.1093/jexbot/51.345.659>.

627 Nagata, M., Yamashita, I (1992). Simple Method for Simultaneous Determinations of Chlorophyll and Ca-  
628 rotenoids in Tomato Fruit. *Nippon Shokuhin Kogyo Gakkaish* 39, 925–928.

629 Naidu, S., Pandey, J., Mishra, L.C., Chakraborty, A., Roy, A., Singh, I.K., Singh, A (2023). Silicon nano-  
630 particles: Synthesis, uptake and their role in mitigation of biotic stress. *Ecotoxicol. Environ. Saf.* 255,  
631 114783. <https://doi.org/10.1016/j.ecoenv.2023.114783>.

632 Nawaz, A.H., Mehmood, A., Khalid, A.u.R., Khan, M.A.R., Gul-e-lala; Ahmad, K.S., Nabi, A.G (2024).  
633 Green Synthesis of Silver Nanoparticles for Their Antifungal Activity against Anthracnose Disease  
634 Causing *Colletotrichum capsici*. *Biocatal. Agric. Biotechnol.* 58, 103178.  
635 <https://doi.org/10.1016/j.bcab.2024.103178>.

Osman, M.S., Sharaf, M., Mekky, A.E., Hassuba, M.M.M., Elballat, E.M., Fouda, H.M (2025). Enhancing of faba bean (*Vicia faba* L.) plant's physiological tolerance against *Rhizoctonia solani* damping-off diseases by application of silver nanoparticle and tebuconazole fungicide. *Physiol. Mol. Plant Pathol.* 136, 102483. <https://doi.org/10.1016/j.pmpp.2024.102483>.

Paz-Trejo, C., Arenas-Huertero, F., Gómez-Arroyo, S (2024). Nano fraction of pesticide induces genotoxicity and oxidative stress-dependent reticulum stress. *Environ. Toxicol.* 39, 1072–1085. <https://doi.org/10.1002/tox.23991>.

Pei, D., Zhang, Q., Zhu, X., Yao, X., Zhang, L (2023). The Complete Genome Sequence Resource of Rhizospheric Soil-Derived *Bacillus velezensis* Yao, with Biocontrol Potential Against *Fusarium solani*-Induced Pepper Root Rot. *Phytopathology* 113, 580–583. <https://doi.org/10.1094/PHYTO-03-22-0101-A>.

Periakaruppan, R., Palanimuthu, V., Abed, S.A., Danaraj, J (2023). New perception about the use of nano-fungicides in sustainable agriculture practices. *Arch. Microbiol.* 205, 4.

Pu, K., Li, N., Gao, Y., Wang, T., Zhang, M., Sun, W., Li, J., Xie, J (2025). Mitigating effects of Methyl Jasmonate on photosynthetic inhibition and oxidative stress of pepper (*Capsicum annuum* L) seedlings under low temperature combined with low light. *Plant Physiol. Biochem.* 223, 109843. <https://doi.org/10.1016/j.plaphy.2025.109843>.

Qiao, H., Chen, J., Dong, M., Shen, J., Yan, S (2024). Nanocarrier-Based Eco-Friendly RNA Pesticides for Sustainable Management of Plant Pathogens and Pests. *Nanomaterials* 14, 1874. <https://doi.org/10.3390/nano14231874>.

Qiao, J., Zhang, R., Liu, Y., Liu, Y (2023). Evaluation of the Biocontrol Efficiency of *Bacillus subtilis* Wettable Powder on Pepper Root Rot Caused by *Fusarium solani*. *Pathogens* 12, 225. <https://doi.org/10.3390/pathogens12020225>.

Qin, X., Wang, Z., Lai, J., Liang, Y., Qian, K (2025). The Synthesis of Selenium Nanoparticles and Their Applications in Enhancing Plant Stress Resistance: A Review. *Nanomaterials* 15, 301. <https://doi.org/10.3390/nano15040301>.

Rathmell, W.G., Sequeira, L (1974). Soluble Peroxidase in Fluid from the Intercellular Spaces of Tobacco Leaves. *Plant Physiol.* 53, 317–318. <https://doi.org/10.1104/pp.53.2.317>.

Rivero-Montejo, S.d.J., Rivera-Bustamante, R.F., Saavedra-Trejo, D.L., Vargas-Hernandez, M., Palos-Barba, V., Macias-Bobadilla, I., Guevara-Gonzalez, R.G., Rivera-Muñoz, E.M., Torres-Pacheco, I (2023). Inhibition of Pepper Huasteco Yellow Veins Virus by Foliar Application of ZnO Nanoparticles in *Capsicum annuum* L. *Plant Physiol. Biochem.* 203, 108074. <https://doi.org/10.1016/j.plaphy.2023.108074>.

Rodríguez, V.M., Velasco, P., Cartea, M.E., Poveda, J (2023). Systemic Biochemical Changes in Pepper (*Capsicum annuum* L.) against *Rhizoctonia solani* by Kale (*Brassica oleracea* var. *acephala* L.) Green Manure Application. *BMC Plant Biol.* 23, 515. <https://doi.org/10.1186/s12870-023-04525-z>.

Senapati, M., Tiwari, A., Sharma, N., Chandra, P., Bashyal, B.M., Ellur, R.K., Bhowmick, P.K., Bollinedi, H., Vinod, K.K., Singh, A.K., Krishnan, S.G (2022). *Rhizoctonia solani* Kühn Pathophysiology: Status and Prospects of Sheath Blight Disease Management in Rice. *Front. Plant Sci.* 13, 881116. <https://doi.org/10.3389/fpls.2022.881116>.

Shahbaz, M., Akram, A., Mehak, A.; Haq, E.u., Fatima, N., Wareen, G., Fitriatin, B.N., Sayyed, R.Z., Ilyas, N., Sabullah, M.K (2023). Evaluation of Selenium Nanoparticles in Inducing Disease Resistance against Spot Blotch Disease and Promoting Growth in Wheat under Biotic Stress. *Plants* 12, 761. <https://doi.org/10.3390/plants12040761>.

Shalaby, T.A., Taha, N., El-Beltagi, H.S., El-Ramady, H (2022a). Combined Application of *Trichoderma harzianum* and Paclobutrazol to Control Root Rot Disease Caused by *Rhizoctonia solani* of Tomato Seedlings. *Agronomy* 12, 3186. <https://doi.org/10.3390/agronomy12123186>.

Shalaby, T.A., El-Bialy, S.M., El-Mahrouk, M.E., Omara, A.E.-D., El-Beltagi, H.S., El-Ramady, H (2022b). Acclimatization of In Vitro Banana Seedlings Using Root-Applied Bio-Nanofertilizer of Copper and Selenium. *Agronomy* 12, 539. <https://doi.org/10.3390/agronomy12020539>

Snedecor, G.W., Cochran, W.G (1989). *Statistical Methods*, 8th ed.; Iowa State University Press: Ames, IA, USA.

Song, J., Yu, S., Yang, R., Xiao, J., Liu, J (2023). Opportunities for the Use of Selenium Nanoparticles in Agriculture. *NanoImpact* 31, 100478. <https://doi.org/10.1016/j.impact.2023.100478>.

Song, X., Zhu, L., Wang, D., Liang, L., Xiao, J., Tang, W., Xie, M., Zhao, Z., Lai, Y., Sun, B., Tang, Y., Li, H (2023). Molecular Regulatory Mechanism of Exogenous Hydrogen Sulfide in Alleviating Low-Temperature Stress in Pepper Seedlings. *Int. J. Mol. Sci.* 24, 16337. <https://doi.org/10.3390/ijms242216337>.

Taha, N.A., Hamden, S., Bayoumi, Y.A., Elsakhawy, T., El-Ramady, H., Solberg, S.Ø (2023). Nanofungicides with Selenium and Silicon Can Boost the Growth and Yield of Common Bean (*Phaseolus vulgaris* L.) and Control *Alternaria* Leaf Spot Disease. *Microorganisms* 11, 728. <https://doi.org/10.3390/microorganisms11030728>.

Tan, N., Yuan, W., Xu, Y., Wang, J., Yuan, B., Huo, H., Qiu, W., Zhou, Y (2025). Migrated Silicon Dioxide Nanoparticles Activates the Rice Immunity for Systemic Resistance against Two Pathogens. *Adv. Agrochem* 4, 78–89.

Vijayreddy, D., Kandan, A., Dutta, P., Pathak, J (2025). Dual Role of Zinc Oxide Nanoparticles in Antifungal Action against *Rhizoctonia solani*, Biosafety and Phytotoxicity Concerns in Crop Plants. *Physiol. Mol. Plant Pathol.* 102712. <https://doi.org/10.1016/j.pmpp.2025.102712>.

704 Wu, Z., Huang, Y., Li, Y., Dong, J., Liu, X., Li, C (2019). Biocontrol of *Rhizoctonia solani* via Induction of  
 705 the Defense Mechanism and Antimicrobial Compounds Produced by *Bacillus subtilis* SL-44 on  
 706 Pepper (*Capsicum annuum* L.). *Front. Microbiol.* 10, 2676.  
 707 <https://doi.org/10.3389/fmicb.2019.02676>.

708 Xiao, M., Liu, X., Wang, X., Lu, W., Zhang, T., Yang, Y (2025). Microbial Inoculants Drive Disease Sup-  
 709 pression and Rhizosphere Modulation for Effective Management of Pepper Phytophthora Blight. *Appl.*  
 710 *Soil Ecol.* 208, 105971. <https://doi.org/10.1016/j.apsoil.2025.105971>.

711 Yadav, S.P., Sharma, C., Pathak, P., Kanaujia, A., Saxena, M.J., Kalra, A (2025). Management of Phy-  
 712 to-Parasitic Nematodes Using Bacteria and Fungi and Their Consortia as Biocontrol Agents. *Environ.*  
 713 *Sci. Adv.* 4, 335–354. <https://doi.org/10.1039/d4va00216d>.

714 Yahia, W.A., Elkhawaga, M.A., Mahfouz, A.Y., Attia, M.S (2023). Achieving Green Synthesis of Silver  
 715 Nanoparticles by *Aspergillus ustus* ON076464 for Improving Immune Response and Vegetative  
 716 Growth of Pepper Plant Towards Wilt Disease Caused by *Fusarium oxysporum*. *Egypt. J. Chem.* 66,  
 717 131–151. <https://doi.org/10.21608/ejchem.2022.161685.6947>.

718 Yan, G., Huang, Q., Zhao, S., Xu, Y., He, Y., Nikolic, M., Nikolic, N., Liang, Y., Zhu, Z (2024). Silicon  
 719 Nanoparticles in Sustainable Agriculture: Synthesis, Absorption, and Plant Stress Alleviation. *Front.*  
 720 *Plant Sci.* 15, 1393458. <https://doi.org/10.3389/fpls.2024.1393458>.

721 Yu, Z., Huang, X., Xiang, J., Qin, X., Xiang, L., Zou, X.; Zhu (2025). PSI-Mediated NADPH Production and  
 722 Enzyme Regulation Are Critical for Carbon Assimilation in Pepper Seedlings under Cadmium Stress.  
 723 *Hortic. Plant J.* <https://doi.org/10.1016/j.hpj.2025.01.005>.

724 Yuce, M., Aydin, M., Turan, M., Ilhan, E., Ekinci, M., Agar, G., Yildirim, E (2025). Ameliorative Effects of  
 725 SL on Tolerance to Salt Stress on Pepper (*Capsicum annuum* L.) Plants. *Plant Physiol. Biochem.* 223,  
 726 109798. <https://doi.org/10.1016/j.plaphy.2025.109798>.

727 Zahedi, S.M., Moharrami, F., Sarikhani, S., et al. (2020). Selenium and Silica Nanostructure-Based Recovery  
 728 of Strawberry Plants Subjected to Drought Stress. *Sci. Rep.* 10, 17672.  
 729 <https://doi.org/10.1038/s41598-020-74273-9>.

730 Zainab, R., Hasnain, M., Ali, F., Abideen, Z., Siddiqui, Z.S., Jamil, F., Hussain, M., Park, Y.K (2024).  
 731 Prospects and Challenges of Nanopesticides in Advancing Pest Management for Sustainable Agri-  
 732 cultural and Environmental Service. *Environ. Res.* 261, 119722.  
 733 <https://doi.org/10.1016/j.envres.2024.119722>.

734