

Synergistic Characteristics of PM_{2.5} and O₃ on the Development of Behavioral Guidance

Xiao-jing Hu^{1, #}, Yong-hong Luo^{2, #}, Yuan-yin Jiang³, Xiao-bo Liu⁴, Hai-quan Yan⁵, , Xiao-yu Li⁵, Ren-jiang He^{4 *}

1. Institute of Drug Clinical Trial/GCP Center, Affiliated Hospital of Southwest Medical University, Luzhou, Sichuan, China

2. College of Public Health, University of South Florida, Tampa Bay, America.

3. School of Basic Medicine, Southwest Medical University, Luzhou, Sichuan, China

4. Environmental Health Effects and Risk Assessment Key Laboratory, School of Public Health, Southwest Medical University, Luzhou, Sichuan, China.

5. Luzhou Ecological and Environmental Monitoring Central Station of Sichuan Province, Luzhou, Sichuan, China

These authors contributed equally to this work.

* Corresponding author: herenjiang@163.com.

Abstract : Objective: This study aims to establish time- and scenario-specific behavioral recommendations for vulnerable populations, based on the pollution characteristics and meteorological drivers of PM_{2.5} and O₃ in Luzhou (2016–2021), with the objective of mitigating health risks associated with air pollution. **Methods:** Hourly data on pollutants and meteorological factors were acquired from Luzhou Environmental Monitoring Station of Sichuan Province. After going through strict quality assurance and control procedures, an effective data collection rate of 96.6% was achieved, temporal distribution and influencing factors were analyzed. **Results:** A significant decreasing trend in PM_{2.5} concentrations was observed, whereas O₃ levels remained persistently elevated (194–213 µg/m³) during the observation period. PM_{2.5} levels peaked during winter nights (21:00–2:00), while O₃ concentrations were highest in the summer afternoons (14:00–15:00). which was closely related to the meteorological conditions (temperature and humidity) in this area. **Conclusions:** Although a reduction in PM_{2.5} levels, O₃ pollution continues to pose a significant challenge. It is recommended that outdoor activities be avoided during winter nights and outdoor exercise be minimized during summer afternoons. Vulnerable populations should limit prolonged exposure when O₃ concentrations exceed 160 µg/m³. These targeted behavioral interventions can effectively reduce acute exposure risks and contribute to public health management.

Key word: Activity guidance; Behavioral modification; O₃; PM_{2.5}; Temporal variation

1. Introduction

With the implementation of the Action Plan for Air Pollution Prevention and Control and the Three-Year Action Plan for Winning the Blue Sky Defense, significant improvements in China's air quality have been observed annually, with a marked reduction in the concentrations of PM_{2.5} and O₃ in ambient air. Despite these reductions, air pollution remained severe in each city, with PM_{2.5} and O₃ identified as the primary pollutants. (Li et al., 2025). However, PM_{2.5} and O₃ pollution continued to pose significant challenges in various cities, particularly with frequent occurrences of O₃ pollution. O₃ pollution exhibited an upward trend in key cities and regions,. The nature of air pollution has progressively shifted toward a composite form characterized by both PM_{2.5} and O₃,

drawing increasing attention to health effect and behavioral changes for the public (Shao et al., 2025) .

Long-term exposure to environments where pollutant concentrations exceed safety thresholds has been linked to significantly elevated prevalence rates of respiratory and cardiovascular diseases among residents, as well as substantial impairments in physiological function and cognitive performance (Xue et al., 2021; Yang et al., 2019; Meo et al., 2024). Moreover, such exposure has been shown to negatively affect subjective well-being and mental health (Yuan et al., 2018). These adverse effects are particularly pronounced in vulnerable groups, including the elderly (Zheng et al., 2024), children (Cui et al., 2023), and other sensitive populations (Ma et al., 2023).

In terms of environmental pollution control and Response, in developed countries, the mechanism for public participation was quite mature. However, many developing countries were only gradually realizing the importance of emphasizing public participation and community development strategies. (Ahmed N. Budur et al., 2008). Air pollution exerts a direct influence on environmental health behaviors, including defensive actions (e.g., purchasing masks and air purifiers, reducing outdoor exposure, and opting for private vehicles to minimize pollutant exposure) (Zhang et al., 2018; An et al., 2018), health-related exercise behaviors (An et al., 2015), migration patterns (Qin et al., 2017), tourism behaviors (Cheung et al., 2001), and eco-friendly product choices (Zhao et al., 2019). However, research suggests that adaptive individual health behaviors during severe pollution episodes are not universally adopted. For example, surveys indicated that 85-90% of residents in Poland and Houston did not reduce driving on days with poor air quality (Semenza et al., 2008), suggesting that individuals may not decrease car usage during smog alerts (Noonan et al., 2014) and may even increase reliance on private vehicles, perceiving them as barriers against pollution (Ban et al., 2017; Neidell et al., 2009). Domestic studies in China indicate that residents adjust travel and residential location choices in response to PM_{2.5} pollution, with younger populations prioritizing active “protection” strategies, while older groups tend to avoid exposure risks through “avoidance” measures. Nevertheless, the consistent adoption of individual adaptive health behaviors during high pollution periods remains infrequent (Na et al., 2016).

Although extensive research have indicated that The proposed IoT-CAPM-DL model, the hybrid RNN-PBO model, a transfer learning-based lightweight recurrent network with skip connection (TL2RN-SC) modelIt can accurately predict air quality, enhance the understanding of air pollution dynamics, provide real-time information to avoid damage to the environment and public health. ((Mohandas P, *et al.*, 2025; Periasamy S,*et al.*, 2024; Prajul Mohandas, et al., 2025) . Meanwhile extensive research has focused on the health impacts of air pollution (Yi et al., 2024; Zhuang et al., 2024; Tan, et al., 2025), only a limited number of surveys have addressed the effects of haze and smog on residents’ health, with few examining the current status of adaptive behaviors employed to cope with air pollution (Karina et al., 2024; Susa et al., 2025). These studies generally lack specific investigations into behavioral changes or actionable travel recommendations for the public. Research explicitly exploring behavioral modifications and providing practical guidance for reducing exposure during atmospheric pollution events remains scarce.

This study aims to address this gap. By analyzing the daily, weekly, monthly, and seasonal pollution characteristics of PM_{2.5} and O₃ in Luzhou, China, from 2016 to 2021, evidence-based recommendations for resident travel and behavioral change strategies will be developed. The objective is to reduce direct exposure and mitigate high-concentration environmental exposure,

thereby lowering disease incidence and minimizing health risks, particularly among vulnerable groups such as the elderly, infants, children, and other sensitive populations. This study is intended to contribute to reducing the disease burden in Luzhou and to provide actionable insights for similar studies in other regions.

2. Data and methods

2.1 Study area

Luzhou City is situated in the southeast of Sichuan Province, China, at the transition zone between the southern margin of the Sichuan Basin and the Yunnan-Guizhou Plateau. The average annual wind speed in Luzhou is relatively low (1.0–2.5 m/s), with a high frequency of calm wind conditions (6.1%), which tends to facilitate the accumulation rather than dispersion of air pollutants, resulting in elevated pollution concentrations. The city's topography, which is lower in the north and higher in the south, creates a natural barrier that further traps pollutants. In addition, Luzhou, with a built-up area of 101 km² and a population of 1.64 million, is a densely populated urban area. The subtropical humid climate, characterized by high humidity and moderate temperatures, also influences the dispersion of air pollutants. These factors collectively contribute to the persistent accumulation of pollutants, rendering air pollution a significant and ongoing challenge. Luzhou was identified as one of the cities in the region most severely affected by air pollution, with consistently high concentrations of key air pollutants, particularly PM_{2.5} and O₃, when compared to other cities in Sichuan Province. Therefore, it has been designated as a key city for air pollution prevention and control. This highlights the importance of characterizing the primary air pollutants, particularly PM_{2.5} and O₃, in Luzhou to inform the development of targeted strategies aimed at improving air quality and safeguarding public health. There are a total of 4 national air quality monitoring stations in the urban area, namely the Jiushishan, Environmental Monitoring Station, Lan Tian Xianqiao, Xiaoshi Dock, As shown in Figure 1.




Figure 1: Distribution Map of Monitoring Stations in the Study Area

2.2 Data sources and monitoring methods

2.2.1 Data sources

The PM_{2.5} and O₃ concentration data were obtained from the Sichuan Luzhou Ecological Environment Monitoring Center Station, encompassing hourly measurements from 00:00 on January 1, 2016, to 24:00 on December 31, 2021. Data cleaning and handling of missing values and outliers were carried out in accordance with the "Technical Specifications for the Operation and Quality Control of Continuous Automatic Monitoring Systems for Gaseous Pollutants in Ambient Air (SO₂, NO₂, O₃, CO)" (HJ 818-2018) and the "Technical Specifications for the Operation and Quality Control of Continuous Automatic Monitoring Systems for Particulate Matter (PM₁₀ and PM_{2.5})" (HJ 817-2018). After excluding records with missing values, a total of 210,432 data entries were gathered, of which 203,277 were valid, yielding an effective data completeness rate of 96.6%. For PM_{2.5}, the hourly concentration of the four monitoring stations was averaged to obtain the hourly average concentration..the daily average concentrations were calculated as the 24-hour mean values, with monthly and annual averages derived from these daily means. For O₃, the hourly concentration of the four monitoring stations was averaged to obtain the hourly average concentration, the daily maximum 8-hour moving average (MDA8) concentration was used as the daily value, with monthly averages computed from these daily MDA8 values and annual averages represented by the annual 4th highest daily maximum 8-hour average concentration. Additionally, meteorological parameters, including air temperature (T) and relative humidity (RH), were collected from the Luzhou Environmental Protection Online Monitoring Data Sharing Platform for the period from January 1, 2018, to December 31, 2021. All 1,462 daily records were valid, ensuring 100% data completeness.

2.2.2 Monitoring instruments and monitoring methods

Ambient PM_{2.5} concentrations were measured using an XHPM2000E particulate monitor (Xianhe Co., Ltd., Hebei, China), which operates based on the beta attenuation method. Ozone (O₃) concentrations were measured using an EC9810 ozone analyzer (EC Pty Ltd, Australia), which employs the ultraviolet absorption method.

2.3 Analytical methods

2.3.1 Classification of pollution levels and compliance assessment

The assessment of air quality was performed in accordance with the secondary limit of the Ambient Air Quality Standard (GB 3095-2012), the National Ambient Air Quality Standards (NAAQS), and the World Health Organization (WHO) Air Quality Guidelines (AQG 2021) .The classification of pollution levels was performed in accordance with National Ambient Air Quality Standards (NAAQS) .Utilizing National Ambient Air Quality Standards (NAAQS) 's annual PM_{2.5} standard (35.0 µg/m³) annual O₃ guideline (137 µg/m³), the pollution conditions during the study period were classified into four distinct levels. This classification system enabled a comprehensive evaluation of air quality by considering both particulate matter and photochemical pollution indicators.

2.3.2 Statistical and graphical tools

Data preprocessing, time-series analysis, and box-plot visualization were conducted using Origin 9.0 (OriginLab Corporation, Northampton, MA, USA). Heatmap generation and nonparametric statistical analyses, including the Mann-Kendall trend test, Mann-Whitney-Wilcoxon rank-sum test, and Kruskal-Wallis one-way ANOVA by ranks, were performed using R software (version 4.5.1; R Foundation for Statistical Computing, Vienna, Austria). A significance level of $\alpha = 0.05$ was applied to all statistical tests.

3. Results and discussion

3.1 Characteristics of PM_{2.5} and O₃

The annual average concentrations of PM_{2.5} and O₃ in Luzhou City from 2016 to 2021 are summarized in Table 1. The concentrations of PM_{2.5} demonstrated a pronounced declining trend (Mann-Kendall, $\tau = -0.93$, $P < 0.01$) from 2016 to 2021, decreasing from 65 $\mu\text{g}/\text{m}^3$ in 2016 to 40 $\mu\text{g}/\text{m}^3$ in 2021. These concentrations were 1.1, 3.3, and 8.0 times above the secondary limit of the Ambient Air Quality Standard (GB 3095-2012; 35 $\mu\text{g}/\text{m}^3$), the National Ambient Air Quality Standards (NAAQS; 12 $\mu\text{g}/\text{m}^3$), and the World Health Organization (WHO) Air Quality Guidelines (AQG 2021; 5 $\mu\text{g}/\text{m}^3$), respectively. A significant reduction in PM_{2.5} concentrations was observed from 2018 onward ($P < 0.01$), aligning with the implementation of the Action Plan for Air Pollution Prevention and Control (2017) and the Three-Year Action Plan for Winning the Blue Sky Defense Battle (2020), both of which had a substantial influence on the reduction of PM_{2.5} levels.

In contrast, O₃ concentrations remained elevated, fluctuating between 194 and 213 $\mu\text{g}/\text{m}^3$, with no discernible trend (Mann-Kendall, $\tau = 0.07$, $P = 0.69$). These concentrations consistently exceeded the limits of GB 3095-2012 (100 $\mu\text{g}/\text{m}^3$), NAAQS (137 $\mu\text{g}/\text{m}^3$), and WHO AQG 2021 (60 $\mu\text{g}/\text{m}^3$), with exceedance factors of 2.0–2.1, 1.4–1.6, and 3.2–3.6, respectively, thereby indicating severe O₃ pollution in the region.

Table 1. Characteristics of annual average PM_{2.5} and O₃ from 2016 to 2021 in Luzhou, ($\mu\text{g}/\text{m}^3$)

Year	PM _{2.5}	O ₃	Days (Percentage) of PM _{2.5} > 35 and O ₃ > 137	Days (Percentage) of PM _{2.5} > 35 and O ₃ ≤ 137	Days (Percentage) of PM _{2.5} ≤ 35 and O ₃ > 137	Days (Percentage) of PM _{2.5} ≤ 35 and O ₃ ≤ 137
2016	65	208	52 (14.2%)	243 (66.4%)	9 (2.46%)	62 (16.9%)
2017	53	194	28 (7.67%)	240 (65.7%)	20 (5.48%)	77 (21.1%)
2018	37	209	11 (3.01%)	124 (34.0%)	31 (8.49%)	199 (54.5%)
2019	41	211	24 (6.58%)	153 (41.9%)	18 (4.93%)	169 (46.3%)
2020	38	213	18 (4.92%)	135 (36.9%)	26 (7.10%)	187 (51.1%)
2021	40	204	11 (3.01%)	140 (38.4%)	24 (6.58%)	190 (52.1%)

Note: In 2019, data were collected for 364 days, with December 31 missing. Data for other years were complete. PM_{2.5} refers to annual average concentrations, while O₃ represents the annual 4th highest daily maximum 8-hour average concentration.

Table 1 presents the proportion of days with simultaneous exceedances of PM_{2.5} and O₃ (PM_{2.5} > 35 $\mu\text{g}/\text{m}^3$ & O₃ > 137 $\mu\text{g}/\text{m}^3$), which decreased from 14.2% in 2016 to 3.0% in 2021, reflecting a cumulative reduction of 78.8%. This decline signifies a substantial improvement in the control of both PM_{2.5} and O₃ pollution. The proportion of days with PM_{2.5} exceedance and O₃ compliance (PM_{2.5} > 35 $\mu\text{g}/\text{m}^3$ & O₃ ≤ 137 $\mu\text{g}/\text{m}^3$) also declined from 66.4% in 2016 to 38.4% in 2021, representing a decrease of 42.2%, further substantiating the significant reduction in PM_{2.5} levels. The proportion of days with PM_{2.5} compliance and O₃ exceedance (PM_{2.5} ≤ 35 $\mu\text{g}/\text{m}^3$ & O₃ > 137 $\mu\text{g}/\text{m}^3$) remained relatively stable, fluctuating between 5% and 8%, except for a value of 2.5% in 2016. Simultaneously, the proportion of days with compliance for both pollutants (PM_{2.5} ≤ 35 $\mu\text{g}/\text{m}^3$ & O₃ ≤ 137 $\mu\text{g}/\text{m}^3$) increased from 16.9% in 2016 to 52.1% in 2021, further demonstrating the overall improvement in air quality management. However, notable discrepancies persist when compared to the annual standard limits established by the WHO (5 $\mu\text{g}/\text{m}^3$ for PM_{2.5}, 100 $\mu\text{g}/\text{m}^3$ for peak season O₃) and the United States Environmental Protection Agency (USEPA, 9 $\mu\text{g}/\text{m}^3$ for PM_{2.5}, 137 $\mu\text{g}/\text{m}^3$ for O₃).

3.2 Trends and seasonal effects on the characteristics of PM_{2.5} and O₃

The monthly average PM_{2.5} concentrations in the urban area of Luzhou from 2016 to 2021 exhibited a distinct U-shaped seasonal pattern, with the lowest levels observed in July and August, and peak concentrations recorded in December, January, and February (Figure 2). Winter consistently presented the highest PM_{2.5} levels across all months, with median concentrations remaining elevated and exhibiting greater variability. This suggests more frequent pollution episodes, likely influenced by meteorological factors such as temperature inversions and stagnant air, alongside increased emissions, particularly from heating activities. Spring and fall exhibited moderate PM_{2.5} levels, with reduced variability compared to winter. Summer was characterized by the lowest PM_{2.5} concentrations, indicative of cleaner air, likely resulting from enhanced atmospheric mixing and dispersion.

The monthly average O₃ concentrations in Luzhou's urban area from 2016 to 2021 demonstrated a stable distribution (Mann-Kendall, $\tau = 0.03$, $P > 0.1$), following a distinct “n”-shaped seasonal pattern (Figure 2). Summer and spring recorded the highest O₃ concentrations, with medians exceeding 100 ppb on numerous occasions. This trend aligns with the intensified sunlight and higher temperatures, which facilitate photochemical ozone formation. In contrast, winter exhibited the lowest O₃ concentrations, frequently falling below 50 ppb, consistent with diminished photochemical activity. Fall, exhibiting intermediate O₃ levels, was lower than summer and spring but higher than winter.




Figure 2. Monthly PM_{2.5} and O₃ concentration trends for each year

During ozone pollution episodes ($O_3 > 137 \mu\text{g}/\text{m}^3$), temperatures ranged from 18.5 to 34.0 °C, while RH varied between 57.1% and 75.0%. In contrast, non-pollution periods ($O_3 \leq 137 \mu\text{g}/\text{m}^3$) were characterized by temperatures between 2.9 and 20.9 °C and RH ranging from 66.1% to 99.8%. During non-ozone-polluted periods, lower temperatures were associated with higher $\text{PM}_{2.5}$ concentrations (Figure 3). Elevated temperatures facilitate chemical reactions and enhance fluxes within the atmospheric boundary layer, thus influencing the distribution of gases and particulate matter. This promotes the conversion of pollutants into gaseous forms, accelerates photochemical reaction rates, and facilitates NO_2 photolysis and volatile organic compounds (VOCs) oxidation, thereby creating favorable conditions for O_3 formation. In contrast, lower temperatures inhibit photochemical fluxes, resulting in reduced O_3 concentrations. High-humidity environments increase ultraviolet scattering by aerosols, diminishing photolysis efficiency. Furthermore, elevated humidity fosters the wet deposition of HNO_3 , curtailing the recycling of NO_x and weakening O_3 production (Zou et al., 2024).




Figure 3. Distribution of temperature and relative humidity at the different $\text{PM}_{2.5}$ and O_3

3.3 Diurnal and weekly cycle of PM_{2.5} and O₃

3.3.1 Diurnal variation of PM_{2.5} and O₃




Figure 4. Hourly boxplot of PM_{2.5} and O₃ by different seasons from 2016 to 2021

Figure 4 presents the boxplots of hourly concentrations of PM_{2.5} and O₃ from 2016 to 2021. PM_{2.5} concentrations during winter were significantly higher than those observed in the other three seasons throughout all 24 hours (Mann-Whitney-Wilcoxon, $P < 0.01$), with the peak occurring between 23:00 and 24:00. A distinct diurnal pattern was evident, characterized by elevated levels from 21:00 to 02:00 and lower concentrations between 13:00 and 17:00. A notable inflection point, marked by a rapid increase in PM_{2.5}, occurred between 17:00 and 19:00. The periods from 11:00 to 13:00 and from 18:00 to 20:00 correspond to morning and evening traffic rush hours, respectively, suggesting a significant contribution from traffic-related dust and vehicle emissions to fine particulate matter concentrations. In contrast, the diurnal variation of O₃ followed a consistent “single-valley, single-peak” pattern across all seasons. This pattern can likely be attributed to suppressed photochemical activity during nighttime, due to lower temperatures, reduced sunlight, and weaker solar radiation. Additionally, nitric oxide (NO) actively consumes O₃, and this titration effect is intensified during traffic peak hours by increased emissions of nitrogen oxides (NO_x) from vehicles, further reducing O₃ levels. Therefore, O₃ concentrations reached their daily minimum between 07:00 and 08:00. From 09:00 onward, strengthening sunlight, rising solar radiation, and increasing temperatures promote photochemical production, leading to the gradual accumulation of

O₃. Peak concentrations typically occurred around 14:00–15:00, followed by a decline as solar radiation diminished and temperatures decreased. Furthermore, O₃ concentrations were highest from June to August, consistent with enhanced photochemical formation driven by more intense solar radiation and higher temperatures during the summer.

3.3.2 Weekly cycle of PM_{2.5} and O₃

Figure 5 illustrates that PM_{2.5} concentrations were significantly higher on all days during winter compared to other seasons (Mann-Whitney-Wilcoxon, $P < 0.01$). The day-of-week effect was found to be minimal. While PM_{2.5} levels remained relatively consistent from Monday to Sunday within each season, a statistically significant difference was observed between weekdays and weekends (Kruskal-Wallis, $H = 10.76$, $P = 0.001$, $\epsilon^2 = 0.0002$). Outliers, characterized by extremely high PM_{2.5} events, were most prevalent in winter. Similarly, O₃ concentrations were significantly elevated across all days in summer compared to winter (Mann-Whitney-Wilcoxon, $P < 0.01$). Additionally, although concentrations remained relatively stable across days, a clear weekday-weekend trend was detected (Kruskal-Wallis, $H = 45.70$, $P < 0.001$, $\epsilon^2 = 0.0009$). Outliers, defined by very high ozone events, occurred predominantly in summer, consistent with episodic high-ozone days. The very small effect sizes ($\epsilon^2 < 0.01$) indicate that these differences stem from the large sample size rather than representing practically significant factors. From a public health perspective, PM_{2.5} and O₃ levels show no substantial variation between weekdays and weekends, suggesting a negligible "weekend effect".




Figure 5. Weekly boxplot of PM_{2.5} and O₃ by different seasons from 2016 to 2021

3.4 Potential effects on behavioral change

During periods of high temperature and low humidity ($T > 30\text{ }^{\circ}\text{C}$, $\text{RH} < 50\%$), intense solar radiation accelerates the photolysis of NO_2 and the oxidation of VOCs, resulting in peak ground-level O_3 concentrations typically occurring between 14:00 and 15:00. To minimize exposure, it is recommended to avoid outdoor activities from 12:00 to 17:00, as this can reduce the inhaled O_3 dose by 40–60%. If outdoor exposure is necessary, it is advised to wear a mask with an activated carbon layer (compliant with GB/T 32610 Class A standards—Class A masks are regarded as the most effective in terms of protection, with a protection level of no less than 90%, and are suitable for use in extremely polluted environments.), replacing it every 4 hours. Vehicle occupants should activate the air conditioning recirculation mode and clean the filters biweekly. It should be noted that vigorous exercise can increase the inhaled O_3 dose by 2–3 times (USEPA, 2022). For commuting by cycling or walking, trips should ideally be scheduled during early morning (06:00–08:00) or evening (19:00–21:00) hours, avoiding major roads to minimize exposure to traffic-related emissions. In the cold and humid season ($T < 10\text{ }^{\circ}\text{C}$ & $\text{RH} > 70\%$), emissions from coal combustion for heating, combined with a thickened inversion layer and stable atmospheric conditions, contribute to a reduction in boundary layer height. High humidity fosters the aqueous-phase formation of SPM, resulting in a bimodal $\text{PM}_{2.5}$ concentration pattern, with peaks between 11:00–13:00 (associated with morning traffic and cooking emissions) and 21:00–02:00 (due to nocturnal boundary layer compression). During hazy days, outdoor activities should be minimized, particularly during the morning hours of 07:00–10:00, when the inversion layer is most pronounced. Indoors, the use of air purifiers equipped with a HEPA filter (with a Clean Air Delivery Rate $\geq 200\text{ m}^3/\text{h}$) is effective. If outdoor activity is unavoidable, the period between 10:00 and 14:00 is relatively safer. Enclosed modes of transportation can significantly reduce personal exposure (Li et al., 2015), and wearing a KN95/N95 respirator (with a filtration efficiency $\geq 95\%$) is strongly recommended. Precipitation events have a scavenging effect on $\text{PM}_{2.5}$, making post-rain ventilation by opening windows beneficial.

Under compound pollution conditions ($T\ 15\text{--}25\text{ }^{\circ}\text{C}$ & $\text{RH}\ 50\text{--}70\%$), synergistic increases in $\text{PM}_{2.5}$ and O_3 concentrations are frequently observed. Monitoring real-time Air Quality Index (AQI) and O_3 levels is essential. Outdoor activities should be minimized when the AQI exceeds 100 or when O_3 concentrations surpass $137\text{ }\mu\text{g}/\text{m}^3$. Vulnerable populations, such as children and individuals with chronic obstructive pulmonary disease (COPD), should adopt an “indoor micro-environment strategy”, establishing a clean air space by utilizing air purifiers, air conditioning, and keeping windows closed.

The implementation of a real-time alert system for special populations is recommended. A push notification advising “Reduce Outdoor Activities” should be activated on mobile devices when the AQI exceeds 100 (indicating an exceedance in either $\text{PM}_{2.5}$ or O_3). For vulnerable groups (e.g., children, COPD patients), a “Red Alert” protocol should be initiated. Automated notifications recommending medical consultation should be dispatched when the AQI exceeds 150 or when O_3 concentrations exceed $180\text{ }\mu\text{g}/\text{m}^3$.

4. Conclusion

From 2016 to 2021, the annual average $\text{PM}_{2.5}$ concentration in the urban area of Luzhou ranged from 38.0 to $65.0\text{ }\mu\text{g}/\text{m}^3$, exhibiting a consistent year-on-year decrease, while the O_3 concentration remained between 194 and $213\text{ }\mu\text{g}/\text{m}^3$, without displaying a significant temporal trend. Throughout

the study period, the number of days with concurrent exceedances of both PM_{2.5} (>35 µg/m³) and O₃ (>137 µg/m³) decreased annually, while the proportion of days meeting both standards (PM_{2.5} ≤ 35 µg/m³ and O₃ ≤ 137 µg/m³) increased correspondingly, indicating effective control of PM_{2.5} and O₃ pollution. PM_{2.5} concentrations were higher in winter and lower in summer, while O₃ concentrations peaked in summer and were at their lowest in winter. PM_{2.5} levels were elevated from 21:00 to 02:00 the following day and lower between 13:00 and 17:00. O₃ concentrations reached their daily minimum between 07:00 and 08:00, with peaks occurring between 14:00 and 15:00.

Based on these patterns, several recommendations for exposure mitigation are proposed: avoiding outdoor activities from 21:00 to 02:00 during winter and wearing KN95 or higher-grade masks if outdoor exposure is unavoidable; reducing outdoor exercise between 13:00 and 17:00 in summer, with a preference for well-ventilated indoor spaces; encouraging children and the elderly to use public transportation and minimize waiting time during cold and humid conditions; and consulting real-time O₃ forecasts before outdoor activities in hot and dry weather, postponing prolonged exercise when O₃ concentrations exceed 160 µg/m³. The implementation of precise, time- and meteorology-specific behavioral interventions is recommended to significantly reduce acute exposure risks for sensitive populations. These findings provide a data-driven foundation for individual health protection and public health management.

5. Research Prospects

This research provides a preliminary analysis of PM_{2.5} and O₃ pollution characteristics and proposes mitigation strategies based on the findings. However, the temporal incongruity between pollution exposure metrics and population health data, alongside privacy constraints, precluded any analysis of individual health effects or exposure-disease relationships. Consequently, epidemiological studies and the evaluation of public health interventions are necessary next steps to corroborate the behavioral suggestions put forth herein.

Acknowledgements

The authors would like to gratefully acknowledge the Data monitoring department of Luzhou Ecological and Environmental Monitoring Central Station of Sichuan Province for providing the pollutant data. We would like to thank EditChecks (<https://editchecks.com.cn/>) for providing linguistic assistance during the preparation of this manuscript.

Funding

This work was supported by Luzhou People's government-Southwest Medical University Science and Technology strategic cooperation project (2023LZXNYDJ050), Scientific Research Project of South-west Medical University (2021ZKQN004, 2022QN065), and Luzhou Science and Technology Project (2022-JYJ-155). Sichuan College Students Innovation and Entrepreneurship Training Project (S202510632177)

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Author contributions declaration

Xiao-jing Hu: Responsible for collecting, organizing and analyzing data, drafting the initial version, and revising the thesis. Email: 334387072@qq.com

Yong-hong Luo: Responsible for designing Paper, drawing charts, writing original paper. Email: yonghong@usf.edu

Yuan-ying Jiang: Responsible for collecting, organizing and analyzing data, providing financial support. Email: j3350265161@qq.com

Xiao-bo Liu: Responsible for collecting, organizing and analyzing data, providing financial support. Email: 1812275340@qq.com

Hai-quan Yan: Responsible for collecting and organizing data, providing financial support. Email: 82491687@qq.com

Xiao-yu Li: Responsible for collecting, organizing and analyzing data, drafting the initial version, and revising the thesis. Email: 278636120@qq.com

Ren-jiang He: Responsible for collecting, organizing and analyzing data, revising the thesis, providing financial support. Email: herenjiang@163.com

References

1. Shaorong Li, Jianhui Guo, Yaqi Wang, et al. The Characteristics of PM_{2.5} and O₃ Synergistic Pollution in the Sichuan Basin Urban Agglomeration[J]. Atmosphere, 2025, 16, 329.

2. Ping Shao, Siyuan Wang, Chaoyong Tu, et al. Variation characteristics and impact factors of the co-occurrence of surface ozone and particulate matter over the Southern Sichuan region, China[J]. Nat Hazards, 2025(121):10807-10824.

3. Xue T, Han Y Q, Fan Y F, et al. Association between a Rapid Reduction in Air Particle Pollution and Improved Lung Function in Adults. Annals of the American Thoracic Society. 2021, 18(2):247-256.

4. Yang BY, Guo Y M, Markevych I, et al. Association of Long-term Exposure to Ambient Air Pollutants With Risk Factors for Cardiovascular Disease in China. JAMA Netw Open. 2019 Mar; 2(3):e190318.

5. Meo S A, Salih M A, Al-Hussain F, et al. Environmental pollutants PM_{2.5}, PM₁₀, carbon monoxide (CO), nitrogen dioxide (NO₂), sulfur dioxide (SO₂), and ozone (O₃) impair human cognitive functions. Eur Rev Med Pharmacol Sci. 2024 Jan; 28(2):789-796.

6. Yuan L, Shin K, Managi S. et al. Subjective well-being and environmental quality: The Impact of air pollution and green coverage in China. Ecological Economics, 2018, 153, 124–138.

7. Zheng XY, Tang SL, Liu T, et al. Effects of long-term PM_{2.5} exposure on metabolic syndrome among adults and elderly in Guangdong, China. Environ Health. 2022 Sep 10; 21(1):84.

8. Cui ZM, Ma YY, Yu YY, et al. Short-term exposure to ambient fine particulate pollution aggravates ventilator-associated pneumonia in pediatric intensive care patients undergoing cardiovascular surgeries. Environ Health. 2023 Apr 26; 22(1):39.

9. Ma H, Zhang Q, Liang W, et al. Short-Term Exposure to PM_{2.5} and O₃ Impairs Liver Function in HIV/AIDS Patients: Evidence from a Repeated Measurements Study. Toxics. 2023, 11(9):729.

404 10. Ahmed N. Bdour, Moshrik R. Hamdi, Moayyad S. Shawaqfeh, et al. Enhancing Public
405 Participation in Local Air Pollution Assessment: A Citizen Participation Prototype from Zarqa
406 Governorate, Jordan. *Environmental engineering science* 2008, 25(4): 451-459.

407 11. Zhang J J, Mu Q. Air pollution and defensive expenditures: Evidence from particulate-filtering
408 facemasks. *Journal of Environmental Economics and Management*, 2018(92),517–536.

409 12. An, R., Yu, H. Impact of ambient fine particulate matter air pollution on health behaviors: A
410 longitudinal study of university students in Beijing, China. *Public Health*, 2018,159, 107–115.

411 13. An R, Xiang X. Ambient fine particulate matter air pollution and leisure-time physical inactivity
412 among US adults. *Public Health*, 2015,129(12), 1637–1644.

413 14. Qin Y, Zhu H J, Run away? Air pollution and emigration interests in China. *Journal of Population*
414 *Economics*, 2018,31(1), 235–266.

415 15. Cheung C., Law, R. The impact of air quality on tourism: The case of Hong Kong. *Pacific*
416 *Tourism Review*,2001,5(1-2), 69–74.

417 16. Zhao, C N, Zhang M, Wang W W. et al. Exploring the influence of severe haze pollution on
418 residents' intention to purchase energy-saving appliances. *Journal of Cleaner Production*, 2019,212,
419 1536–1543.

420 17. Semenza J C, Wilson D J, Parra J, et al. Public perception and behavior change in relationship
421 to hot weather and air pollution. *Environmental Research*, 2008,107(3),401–411

422 18. Noonan D S. Smoggy with a chance of altruism: The effects of ozone alerts on outdoor recreation
423 and driving in Atlanta. *Policy Studies Journal*, 2014,42(1), 122–145.

424 19. Ban J, Zhou L, Zhang Y, et al. The health policy implications of individual adaptive behavior
425 responses to smog pollution in urban China. *Environment International*, 2017,106, 144–152.

426 20. Neidell M. Information, avoidance behavior, and health the effect of ozone on asthma
427 hospitalizations. *Journal of Human Resources*, 2009,44(2), 450–478.

428 21. NA Jun, LI Yan-xia, CHEN Kang-jing, et al., Survey on Residents' Individual Adaptive
429 Behaviors in Coping with Air Pollution in Five Cities/Counties of Liaoning Province, *J Environ*
430 *Occup Med* 2016,33 (3) ,263-266.

431 22. Prajul Mohandas, Subramanian P, Surendran R. Optimizing Air Pollution Prediction in Urban
432 Environments using a Hybrid RNN-PBO Model with IoT Data. 2025 International Conference on
433 Visual Analytics and Data Visualization (ICVADV), Tirunelveli, India, 2025, 480-487.

434 23. Mohandas P, Subramanian P, Rajendran S. Air Pollution Monitoring System using Stacked
435 Attentional Vectormap Convolutional Bidirectional Network with Bobcat Optimization and IoT-
436 Cloud, *Global NEST Journal*, 2025, 27(3), 06937

437 24. Periasamy S, Subramanian P, Surendran R. An intelligent air quality monitoring system using
438 quality indicators and Transfer learning based Lightweight recurrent network with skip
439 connection. *Global NEST*, 2024, 26(5), 06096.

25. Zhang Y, Zeng J, Tian X Y, et al. Estimating the mutually adjusted health effects of short- and long-term exposure to PM_{2.5} on respiratory mortality in a population-based study, *Atmospheric Pollution Research*, 2024,15(5), 102091.
26. Shao Z, Zheng X, Jing Zhao, et al. Evaluating the health impact of air pollution control strategies and synergies among PM_{2.5} and O₃ pollution in Beijing-Tianjin-Hebei region, China, *Environmental Research*, 2025, 274, 121348
27. Tan Y X, Xiang, Q Q, Zhang, Y L, et al.,. Impaired physical function in relation to later-life exposure to ambient fine particulate matter and ozone among Chinese middle-aged and older adults. *BMC Public Health* 2025, 25(1):2616.
28. Su B B, Liu, C, Chen, L, et al. Xiaoying Long-term exposure to PM_{2.5} and O₃ with cardiometabolic multimorbidity: Evidence among Chinese elderly population from 462 cities. *Ecotoxicology and Environmental Safety*. 2023, 255, 14790
29. Zheng X Y, Tang S L, Liu T, et al. Effects of long-term PM_{2.5} exposure on metabolic syndrome among adults and elderly in Guangdong, China. *Environ Health*. 2022, Sep 10; 21(1):84.
30. Cui Z M, Ma Y Y, Yu Y Y, et al. Short-term exposure to ambient fine particulate pollution aggravates ventilator-associated pneumonia in pediatric intensive care patients undergoing cardiovascular surgeries. *Environ Health*. 2023 Apr 26, 22(1):39.
31. Ma H F, Zhang Q, Liang W, et al. Short-Term Exposure to PM_{2.5} and O₃ Impairs Liver Function in HIV/AIDS Patients: Evidence from a Repeated Measurements Study. *Toxics*. 2023 Aug 25; 11(9):729.
32. Susa, A, Zekovic, M, Davidovic, D, et al. From Perception to Action: Air Pollution Awareness and Behavioral Adjustments in Pregnant Women in Serbia. *Healthcare*. 2025, 13(12):1475.
33. Landeros-Mugica K, Urbina-Soria J, Angeles-Hernandez DI, et al. Air Pollution and Climate Change Risk Perception among Residents in Three Cities of the Mexico Megalopolis Atmosphere. 2024, 15(1):42.
34. Li Y P, Fan Z Y, Li K, et al. Commuter exposure to PM_{2.5} and its influencing factors in different commuting modes[J]. *Environmental Chemistry*, 2015, 34(8):1408-1416.