

Comparative study of the diversity and distribution of crustacean communities in three freshwater biotopes (Tonga, Birds and El Kennar lakes) located in northeastern Algeria

Imene Bali¹, Hinda Berghiche*¹, Fatiha Bendali-Saoudi¹ and Choukri Barour²

¹Laboratory of Applied Animal Biology, Faculty of Sciences, Department of Biology, Badji Mokhtar - Annaba University12, P.O. Box, 23000 Annaba, Algeria.


²Laboratory of Aquatic and Terrestrial Ecosystems, Faculty of Natural and Life Sciences, University of Mohamed Chérif Messaadia, BP 1553, Annaba Road, Souk Ahras, Algeria.

Received: 24/04/2025, Accepted: 09/12/2025, Available online: 10/12/2025

*to whom all correspondence should be addressed: e-mail: hinda.berghiche@univ-annaba.dz

<https://doi.org/10.30955/gnj.07578>

Graphical abstract



Abstract

The crustacean group is a keystone species that plays a significant role in the aquatic ecosystem by participating as decomposers or consuming detritus. Understanding the ecology and spatiotemporal distribution of crustaceans' communities is crucial for the effective management of the species and the associated structural and functional services that they provide in freshwater ecosystems. This study focused on freshwater crustaceans' communities from northeastern Algeria, located over a distance of 337 km. The objective of this research is to present data on the distribution and composition of the communities, evaluate both α - and β -diversity, and identify underlying potential factors or parameters that can serve as baseline data for future studies pertaining to global changes. From May 2022 to April 2023, the target crustacean populations were sampled every two months from Tonga Lake (El Kala National Park), Birds Lake (El Tarf province), and El Kennar Lake (Jijel province). A total of 8,175 specimens were sampled and inventoried; 14 species were identified belonging to 9 genera: 9 Cladocerans species, 3 Copepods species, one species of Ostracods and another of Decapod. The recorded taxa match 7 families: Daphniidae, Cyclopidae, Cyprididae, Atyidae, Chydoridae, Temoridae and Moinidae. The non-metric multidimensional scaling

(NMDS) ordination analysis showed a clear spatial effect in the crustacean communities and highlight three distinct species assemblages between lakes. The NMDS also showed possible effects of rainfall, temperature, pH, electrical conductivity, and water temperature on community composition and distribution. Furthermore, the most abundant species were *Cypris sp* (40.36%), *Cyclops fuscus* (38.10%), and *Simocephalus expinosus* (56.19%) in Tonga Lake ($S=9$), Birds Lake ($S=9$), and El Kennar Lake ($S=10$), respectively. The Shannon index showed similar levels of diversity, with the highest value in Tonga Lake ($H'=2.48$, moderate level of diversity). The Pielou evenness index also showed higher equitability in both Tonga and Birds lakes than in El Kennar one; this latter recorded the highest species abundance ($A=3,771$). According to Sorenson's coefficient, our communities have quite a bit of overlap (from 63% to 78% of similarity). Our research essentially confirms the hypothesis that global changes and possibly biotic and physical processes may have an effect on the biodiversity and composition of crustacean communities in Algeria. The functional diversity and the interactions with environmental factors, from global to local scales should be studied in our ongoing project concerning our local crustaceans' communities that are still poorly documented.

Keywords: Biodiversity; crustaceans; ecological indexes; freshwater ecosystems; NMDS ordination; physicochemical parameters.

1. Introduction

Biodiversity is particularly noteworthy in freshwater ecosystems. They are the most dynamic, with a high number of species (Rolls *et al.* 2018). Freshwater habitats occupy, 1% of the Earth's surface, yet are hotspots that support, 10% of all known species (Strayer and Dudgeon 2010). Freshwater crustacean communities play an important role in nutrient cycling and the transfer of energy from primary producers to higher trophic levels in

aquatic food systems as herbivores, detritivores, predators and prey (Lomartire *et al.* 2021). They also serve as biological indicators for assessing surface water quality (Krepski *et al.* 2024). However, they are subject to unprecedented levels of human disturbance, with varying degrees of impacts on water, sediments and biota. Biomonitoring of these communities has therefore become a widely accepted and irreplaceable tool for ecological conservation and management of aquatic ecosystems (Xiong *et al.* 2020). Their short reproductive cycles make them responsive to environmental changes, making them effective indicators of climate change (Beaugrand *et al.* 2015).

Crustaceans originated ~ 500 million years ago during the Precambrian period (Zhang *et al.* 2007). Although most crustaceans are aquatic and free-living, some are terrestrial (e.g., woodlice) or parasitic (e.g., fish lice). They are the most species-rich group of marine animals, living from the deepest ocean areas to muddy wetlands (Wacker and Harzsch 2021). There are approximately 150,000 species exist in the worldwide, of which only 52,000 have been described (Mahmood Ghafor 2020). There are four primary clades of crustaceans: Branchiopoda (e.g., Anostraca, Notostraca, and Phyllopoda) (Walossek 1993), Maxillopoda (e.g., Copepoda, Branchiura and Cirripedia) (Suárez-Morales 2015), Malacostraca (e.g., Eumalacostraca, Hoplocarida) and Ostracoda (e.g., Myodocopa, and Podocopa) (Schwentner *et al.* 2017). In addition, crustaceans contain two minor clades, Remipedia and Cephalocarida (e.g., horseshoe shrimp), each of which contains 12–17 species (Schwentner *et al.* 2017).

The concept of biodiversity hotspots pioneered by ecologist Norman Myers as a template for setting global conservation priorities (Myers *et al.* 2000). The map of hotspots overlaps extraordinarily well with the map of the natural sites that are most beneficial to humanity (Pimm and Raven 2020). Hotspots are some of the richest and most important ecosystems in the world, and are home to many vulnerable populations whose survival depends directly on nature (Myers *et al.* 2000). They are particularly sensitive to human disturbance and much of their remaining high-quality habitat persists only within protected areas (Mittermeier *et al.* 2004). According to one estimate, although they represent 2.5% of the Earth's surface, forests, wetlands and other ecosystems (Barrett *et al.* 2016), in hotspots are responsible for 35% of the 'ecosystem services' on which vulnerable human populations depend. This is why Conservation International has adopted the idea of protecting these incredible areas known as 'hotspots'. Today, the mission has expanded beyond protecting hotspots. Because it is not enough to protect species and places; for humanity to survive and flourish, protecting nature must be a fundamental part of every human society (Brooks *et al.* 2006).

Mediterranean wetlands identified in the regional context as 'biodiversity hotspots' (Cañadas *et al.* 2014). They play a key role in supporting an exceptional diversity of plant and animal species, many of which are endemic and rare.

This makes them one of the most diverse ecosystems on the planet (Rifai *et al.* 2018). The freshwaters of the circum-Mediterranean countries are indeed known for their exceptional biodiversity, housing. Thirty-five percent of Palearctic freshwater species and more than 6% of the world's freshwater species with a high rate of endemism (Tierno de Figueroa *et al.* 2013). The exceptional diversity of freshwater fauna of the Mediterranean Basin currently faces a crisis in which climate change combined with overexploitation of freshwaters heavily threatens the local fauna (Ayati *et al.* 2023). Recent studies have shown that physical and chemical alterations can have adverse impacts on biota (Krepski *et al.* 2024). These alterations include sewage discharge (Tripathi *et al.* 2023), elevated water temperatures due to thermal pollution (Poole and Berman 2001), and increased river salinity resulting from mining activities (Slugocki and Czerniawski 2023). Thus, environmental parameters play a pivotal role in influencing the presence and spatial distribution of zooplankton (Umi *et al.* 2024). Responses of crustaceans to climate variability and change are not uniform and vary depending on the species, life-history stage, reproductive status, and geographical distribution (Azra *et al.* 2022). Fluctuations in crustaceans populations are influenced by various factors, such as temperature, phytoplankton and the density of predators, particularly fish species (Gouws *et al.* 2010).

Algeria is characterized by a climate that is extremely favorable to the wide distribution of crustaceans. Lucas conducted previous studies on Ostracods (1849), followed by Moniez. (1891), Gauthier (1928a, b, and c), Gauthier and Brehm. (1928) and De Deckker (1981). No further work was carried out until the study by Samraoui *et al.* (1998), which provided checklists of species of Rotifers, Cladocerans, Copepods and Ostracods from Numidia, in the north-east of Algeria. More recently, the two studies of Ghaouaci *et al.* (2017 and 2018) including non-marine living Ostracods, Cladocerans and Copepods from the continental waters of Algeria. Our study provides the first comparative analysis of crustacean species living in different ecoregions of Algeria. It fills an important gap in regional biodiversity research. We applied useful and powerful ecologic analysis by using many R packages to differentiate species' assemblages: NMDS analysis and DER algorithm. The Numeric Ecology provides more technics to characterize the fauna and the flora structure and composition. While previous studies in Algeria have mostly focused on individual sites (e.g., Lake Tonga, Birds Lake, or isolated branchiopod records), present work integrates data from different habitats to identify broader ecological patterns. On a worldwide scale, this study contributes to freshwater biodiversity research by providing new information on how crustacean assemblages respond to ecological gradients in North Africa, a poorly studied region that can improve global understanding of biogeographic distribution and conservation priorities.

Unfortunately, the diversity and distribution of the crustacean species in Algeria have not been reported in

detail and research on crustacean diversity is rarely carried out (Ayati *et al.* 2019). Given the important role of the crustacean group in the freshwater ecosystems especially in Tonga, Birds and El Kennar lakes, it is important to study crustacean diversity in many Algerian ecoregions. We hypothesized that crustacean community composition could prove fine-scale variation in biotope types and water quality because crustacean species could respond typically to environmental gradients. Thus, the purpose of this study was, firstly, to determine the diversity and distribution of the crustacean taxa along a geographic transect in the north-eastern of Algeria to appraise both α - and β -diversity levels, secondly, to contribute to the conservation and monitoring of the freshwater ecosystems as a precious resource.

2. Material and methods

2.1. Study area

The study took place during May 2022–April 2023 period and was conducted in the north-eastern region of Algeria in order to cover spatial and temporal changes in the crustacean community. Three specific lakes were selected to carry out sampling operations (**Figure 1**).




Figure 1. Crustacean sampling sites in two geographical regions of eastern Algeria: 1- Tonga Lake, 2- Birds Lake (El Tarf) and 3- El Kennar Lake (Jijel), Arcgis 10.4, (2025).

-Tonga Lake (36°53'1.03" N, 8°31'46.60" E) is a 2,600 hectares freshwater wetlands connected to the Mediterranean Sea by the artificial Strait of Messida. Although, it is within the boundaries of the El Kala National Park under the Ramsar Convention (1983). This site was selected due to its recognition as a key Ramsar site with limited existing data on crustacean communities.

Table 1. Mean, standard deviation, minimal and maximal values of the environmental parameter measured in Tonga, Birds and El Kennar lakes from May 2022 to April 2023. For each parameter, the different letters indicate significant differences, according to Dunn's test, between lakes.

Site		Temperature (C°)	pH	Conductivity (µs/cm)
Tonga Lake	mean	23.38 <i>a</i>	7.65 <i>a</i>	1427 <i>b</i>
	sd	2.70	0.68	188.74
	min	19.20	7.05	1100
	max	26.60	8.80	1598
Birds Lake	mean	22.07 <i>a</i>	8.48 <i>a</i>	648.33 <i>a</i>
	sd	3.86	0.39	262.44
	min	17.90	8.00	400
	max	27.20	9.00	980
El Kennar Lake	mean	21.78 <i>a</i>	7.56 <i>a</i>	799.17 <i>ab</i>
	sd	3.90	0.93	452.30
	min	17.70	6.66	414
	max	26.50	8.90	1600

The 2016 study remains one of the few providing quantitative information on the prevalence of these organisms within macroinvertebrate assemblages.

Birds Lake (36°47' N 08°7' E), covering 150 hectares, has been a Ramsar site since 1999, known for its biodiversity. It forms part of a larger wetland complex in eastern Algeria, much of which is designated as a national park. Tonga and Birds lakes are located in El Tarf province. This site was selected because it hosts various crustacean groups, including Cyclopidae (cyclopoid copepods), Daphnidae (cladocerans), and Cyprididae (ostracods), highlighting its ecological importance in Algerian freshwater habitats.

El Kennar Lake (36°48' N; 05°57' E), also known as Ghedir Béni Hamza, is located to the east of Jijel province and covers an area of approximately 36 hectares. This site was chosen because it represents a vulnerable aquatic ecosystem that is heavily impacted by anthropogenic pressures, particularly urban and agricultural contamination, making it valuable for studying human-induced effects on crustacean communities.

This study focuses on the Algerian region, where we worked over a distance of 300 km as the crow flies and respected the distance between the three lakes. In addition, the crustacean biodiversity of Algerian freshwater is underestimated, and further research is needed to supplement and deepen the current conclusions.

2.2. Sample collection

To cover a broad range of taxa, samples of crustacean specimens were collected every two months from May 2022 to April 2023. The freshwater was characterised on the basis of several physicochemical parameters such as temperature (T) in °C, pH and conductivity (EC) in µs/cm, measured by a multi-parameter water analyzer (HI 9829). Temperature and rainfall data were kindly provided by the Agricultural Services of El Tarf and Jijel provinces (**Table 1**). Bimonthly sampling is sufficiently regular and more accurate for monitoring changes in communities, as it allows intermediate transitions between seasons to be captured. It thus offers a better balance between ecological representativeness and field constraints.

Each station was defined as a research area delimited by GPS. Within each area, samples were taken at random using a hand net (300 μm mesh) on the shore for an area of 1 m^2 (Decapoda) and a plankton net (64 μm) in the pelagic zone with a volume of 1 L (microcrustaceans), using a standardized effort (same number of net hauls or net pulls per site). This protocol ensured that the sampling effort remained constant across the three sites in each lake, reduced biases related to environmental variability, and provided comparable data between the different ecosystems studied.

2.3. Laboratory analyses

Samples were collected from 3 stations in each lake for crustacean inventory. Sampling methods should be adapted to the habitat and the group to be studied. According to Mack *et al.* (2012), qualitative samples of freshwater crustaceans were collected using a hand net measuring 64 μm , which effectively captured all zooplankton groups, and they were preserved immediately in 70% ethanol solution in polyethylene bottles. In the laboratory, the specimens of crustaceans were examined using a Olympus microscope CH20 for a dissection, small entomological pins can be mounted. The dissection can be easily performed if a specimen is placed on the slide with glycerin (or other viscous medium).

Ostracods specimens were dissected under a microscope. The valves were opened and the soft body extracted in order to isolate the various appendages (antennae, walking legs, cleaning legs, uropod, maxillipeds, maxillules, mandibles, fork, and male copulatory organs), all of which are necessary for taxonomic identification by Meisch. (2000).

Cladocerans were dissected using entomological needles in a glycerin solution in order to observe the details of the antennules and rostrum, or the details of the spinulation of the dorsal edge of the ephippium or on the edges of the carapace, cephalic shield, and thoracic limbs. Using as a reference the contribution of Błędzki and Rybak (2016).

Copepods were dissected under a microscope using fine needles on semi-permanent slides, using a glycerin solution to separate the different body parts such as the double genital somite, caudal rami, antennules, labrum, mandibles, maxillary palps, maxillae, and maxillipeds (Błędzki and Rybak 2016).

Shrimps identified according to morphological criteria (body shape, size, and coloration; number of rostral teeth; degree of abdominal development and its relative position in relation to the cephalothorax; presence or absence of a keel and tooth on the abdominal segments; and finally, shape and length of the antennae and antennules) according to Fischer *et al.* (1987).

2.4. Statistical analysis

All our statistical analyses were performed using R, version 4.4.1 (R Core Team 2024; Ihaka and Gentleman 1996) and RStudio (Posit Team 2024) for Windows. Shapiro-Wilk test was used to test variables normality. The main aim of this survey is to characterize both α - and

β -diversity levels of the crustacean communities in freshwater ecosystems. Shannon (H') diversity index and Pielou's evenness index (J') were calculated using 'EcolIndR' package (Gonzalez 2023). We also calculated the Sørensen similarity index to assess the quotient of similarity for two communities based on the number of species in each community and the number of species in common. This index varies between 0 (no overlap) and 1 (perfect overlap). In an ecological context; no one diversity index can adequately summarize species diversity and for this reason we used the DER algorithm (Gonzalez 2023) in our study to estimate multiple diversity indexes and for comparing crustacean species diversity between assemblages. All indexes were transformed to a scale range of 0 to 1, and the algorithm calculated the polar coordinates of all samples with all possible combinations for all indexes and other ecological descriptors. This algorithm provides tools for differentiating assemblages on the basis of many diversity indexes. The DER algorithm was applied to maximize dispersion among our three freshwater biotopes. Also, we can determine the differences in terms of rarity, abundance, heterogeneity, richness, evenness and diversity among assemblages. To investigate the crustacean species assemblages; we carried out also a multivariate analysis by applying a Non-metric MultiDimensional Scaling (NMDS) as a distance-based ordination technique. Because it focuses on the distance matrix, it is very flexible and any distance measure can be used. As usual, the data matrix (n sample units \times p species) is converted into an $n \times n$ distance matrix (or a dissimilarity matrix). The NMDS algorithm was performed by using 'vegan' R package edited by Oksanen *et al.* (2025). Finally, we also examined, by an ombrothermic diagram, the relationship between temperature and precipitation to determine the length of dry, wet, and extremely wet periods in the selected geographical regions. Data were expressed as mean \pm standard error (se). Several R packages were also used in our statistical analysis and to plot data results such as 'ggplot2' (Wickham 2016), 'ggcorrplot' (Kassambara 2023a), 'webr' (Moon 2020), 'Hmisc' (Harrell 2025), 'ggpubr' (Kassambara 2023b), 'PMCMRplus' (Pohlert 2024), 'dunn.test' (Dinno 2024) and 'psych' (Revelle 2024). All these R free-packages are also available on CRAN web site (The Comprehensive R Archive Network: <https://cran.r-project.org>).

3. Results

3.1. Crustacean species composition

In general, a total of 8,175 individuals of crustaceans were recorded at the Tonga, birds and El Kennar lakes, which included 14 species of freshwater crustaceans belonging to 9 genera (*Ceriodaphnia*, *Daphnia*, *Simocephalus*, *Chydorus*, *Moina*, *Cyclops*, *Temora*, *Cypris* and *Atyaephyra*) and 7 families (Daphniidae, Temoridae, Cyclopidae, Atyidae, Chydoridae, Cyprididae and Moinidae). Nine species were reported from Cladocera group, 3 species from Copepoda and one species was reported from Ostracoda and Decapoda group (**Figure 2**).




Figure 2. Species abundance stacked barchart showing the centesimal frequencies of different crustacean freshwater species recorded in Birds, El Kennar and Tonga lakes located in northeast of Algeria during the sampling period: May 2022-April 2023. Each color represents a different species and each bar represents a single site ($A = 8175$, $S = 14$).




Figure 3. Pie charts representing the crustacean freshwater taxonomic composition in Tonga lake (A), Birds Lake (B) and El Kennar Lake (C) communities. The percentages and the taxonomic assignment are given at the species level during the sampling period: May 2022-April 2023 from three northeastern regions of Algeria. The recorded richness (S) in each biotope is respectively: 9, 9 and 10 species.

In our study, the Cladocera group was represented by *Chydorus brevilabris* (Frey), *Ceriodaphnia sp.*, *Moina micrura* (Kurz), *Daphnia magna* (Straus), *Daphnia pulex* (Leydig), *Daphnia sp.*, *Daphnia atkinsoni* (Braid), *Simocephalus expinosus* (Koch) and *Simocephalus vetulus* (Müller). Moreover, the Copepoda group was represented by *Cyclops fuscus* (Jurine), *Cyclops sp.* and *Temora affinis* (Poppe), and the Decapoda and Ostracoda groups were represented by the lowest number of taxa, namely *Atyaephyra desmarestii* (Millet) and *Cypris sp.*

Pie graphs illustrate the percentages of taxa and taxonomic composition of the crustacean communities in Tonga, Birds and El Kennar lakes (Figure 3). It can be seen

that the highest richness, in the different sampling sites, of freshwater crustaceans was recorded in El Kennar Lake (10 species with 3,771 specimens); the lowest one was recorded in Birds Lake (9 species with 3,345 specimens) followed by Tonga Lake (9 species with 659 specimens). Furthermore, the species *Cypris sp.* was the most abundant (40.36%) in Tonga Lake, while *Cyclops fuscus* (38.10%) and *Simocephalus expinosus* (56.19%) species dominated in Birds and El Kennar lakes, respectively.

3.2. Ecological indexes

In ecology, the conception of species diversity is used to assess the facility of the communities to maintain itself stable; even though it gets disturbed. In this sense, the diversity level of crustaceans' assemblages in our three freshwater biotopes was estimated using the Shannon index (H'), Pielou's evenness index (J') was also calculated to evaluate how the 14 species are evenly distributed in our communities related to the three lakes.

In Figure 4, it can be seen that the diversity of our crustacean communities in the three distinct biotopes is in the medium category of the biodiversity Shannon's index; this latter varies from $H' = 2.10$ to $H' = 2.48$. We also conclude that Tonga Lake is the most diverse lake despite its low species' abundance (as shown by Figure 4). Regarding to the Pielou's evenness index, the highest value was recorded in Tonga Lake ($J' = 0.78$) and the lowest one was recorded in El Kennar Lake ($J' = 0.63$); these results mean that Tonga and Birds lakes have more equitable distribution of their species; however the El Kennar Lake is characterized by the highest dominance rate of species which is *Simocephalus expinosus* (56.19%), as shown in Figure 3.




Figure 4. Diversity of the crustacean freshwater communities in Tonga, Birds and El Kennar lakes estimated at the α -diversity level during the sampling period: May 2022-April 2023 from three northeastern regions of Algeria. The barplots are expressing two ecological indexes (Shannon's diversity index, Pielou's evenness index) and two other ecological parameters (Abundance and Richness).

As mentioned in the materials and methods section; we used the DER algorithm for differentiating samples on the basis of several ecological indexes that better reflect the dispersion among species assemblages. The DER algorithm is used in ecological surveys because sometimes no one diversity index can adequately summarize species diversity. All indexes were transformed to a scale range of 0 to 1, and the algorithm calculated the polar coordinates of all samples with all possible combinations for all indexes and other ecological descriptors (as abundances). This algorithm provides tools for differentiating assemblages on the basis of many diversity indexes; the DER algorithm result is resumed by the **Figure 5** and show the similarity level between crustacean communities.




Figure 5. Differentiating assemblages of crustacean species on the basis of the DER algorithm (Gonzalez 2023) applied to abundances, Shannon's diversity index, Pielou's evenness index and Margalef's index. Color gradient is the rarity Leroy index in each sample site. Bubble size scale is the species richness (S). The DER algorithm summarizes species diversity among all of freshwater crustacean samples collected from Tonga, Birds and El Kennar lakes in northeast Algeria during the sampling period: May 2022-April 2023.




Figure 6. The non-metric multidimensional scaling (NMDS) of the freshwater crustacean communities' structure. The ordination plot is based on the Bray-Curtis dissimilarity matrix (stress value = 0.00). The ordination plot shows both the lakes (biotopes) and the crustacean species (S=14). The blue arrays present possible effects of measured environmental and climatic variables on species assemblages. The NMDS plot clearly shows a noteworthy level in β -diversity.

For further investigation, in our survey, the idea of the NMDS analysis is to specify the original positions of the crustacean communities in multidimensional space and to define the number of reduced dimensions (typically 2). The ordination plot (**Figure 6**) presents an excellent representation of our samples in reduced dimensions because the recorded stress value is equal to 0 (<0.2). As expected, the ordination study of the crustacean species assemblages was conclusive about the diversity and composition in El Kennar, Tonga and Birds lakes. Obviously, Tonga Lake is more characterized by *A. desmaresti* and *T. affinis* species while, *Cyclops sp.*, *M. micrura* and *S. vetulus* species correspond more to Birds Lake. By contrast, El Kennar Lake is more marked by the presence of the following species: *Daphnia sp.*, *Ceriodaphnia sp.*, *D. atkinsoni* and *C. brevilabris*.

In conclusion, NMDS analysis showed that the three lakes were clearly separated according to their crustacean communities. The 1st axis (NMDS1) separated El Kennar Lake (S= 10, A= 3,771) and Birds Lake (S= 9, A= 3,745) from Tonga Lake (S= 9, A= 659); whereas the 2nd axis (NMDS2) distinguished very well the analyzed lakes. The NMDS ordination revealed an overlap between the three communities; this is justified by the presence of common species. This result is supported by the Sørensen index, which reflects the level of similarity between crustacean communities: 63% between Tonga/El Kennar lakes (6 species in common), 74% between Birds/El Kennar (7 species in common) and 78% between Tonga/Birds lakes (7 species in common). Additionally, the NMDS ordination plot shows, firstly, a significant effect of the rainfall on the species composition in El Kennar Lake, secondly, a notable effect of both water temperature and conductivity on species composition in Tonga Lake, and thirdly, a possible effect of water pH on species composition in Birds Lake.




Figure 7. Ombrothermic diagram showing the monthly mean temperature (red line) and rainfall (bars) in both El Tarf and Jijel provinces, Algeria, from May 2022 to April 2023.

3.3. Climatic data and physicochemical parameters

The temperature and rainfall data obtained from the regional weather stations are presented by the ombrothermic diagrams which are shown in **Figure 7** for both Jijel and El Tarf provinces from May 2022 to April 2023. The two studied geographic regions showed a relative uniform climate characterized as mild Mediterranean. The yearly average rainfall was 62.95 (with a maximum of 228 mm) and 42.00 (with a maximum of 73.00 mm of maximum) mm in Jijel and El Tarf provinces, respectively. This observation states that the geographic region of El Tarf is more dry than in Jijel. Annual average temperature was 20.83°C in the geographic region of El Tarf and 18.83°C in the geographic region of Jijel. The hottest months were 26.00 and 30.00°C in Jijel and El Tarf geographic regions, respectively. These climatic characteristics could impact the distribution and abundances of the crustaceans' communities in their biotopes.

The results related to data analysis of the physicochemical parameters are shown in **Table 1**. The comparisons were carried out by Kruskal-Wallis test followed by Dunn's test for *post hoc* pairwise comparisons. The results showed that there is a significant difference ($p < 0.05$) between the studied lakes only for conductivity parameter. On the other hand, there were no significant differences ($p > 0.05$) between lakes for temperature and pH parameters.

4. Discussion

Biodiversity assessment is a fundamental requirement for the efficient management of species and the structural and functional benefits they provide in freshwater ecosystems, including wetlands (Cantonati *et al.* 2020). As biodiversity loss has become an urgent global issue over the past decade, for this reason, recent research has rekindled interest in biodiversity in these environments (Tickner *et al.* 2020). While the earliest models of biodiversity response often examined a single metric at one spatial scale, most commonly examining species richness within local communities, contemporary biodiversity research typically uses multiple diversity metrics. These include the number of species in a local community (α -diversity), compositional differences between local communities (β -diversity), and total species diversity within landscapes (γ -diversity) (Anderson *et al.* 2011).

North Africa is confronted with extreme conditions of drought and high temperatures that make them even more vulnerable to climate change, with serious consequences on biodiversity and ecosystem stability (Zouggar *et al.* 2023). This is one of the regions overlooking the Mediterranean Sea, which has been considered one of the main hotspots of climate change, suffering from water scarcity, overexploitation, saline intrusion in coastal aquifers due to sea level rise, and reduced resources (Sharan *et al.* 2023). According to the recent Assessment Report on "*Climate and Environmental Change in the Mediterranean Basin*" published by Cramer *et al.* (2022), the Mediterranean is warming 20% faster

than the global average. Consequently, this region is well-defined as a hotspot area based on both current observations and future projections (Urdiales-Flores *et al.* 2023).

Given the high population growth in the northern part of Algeria, combined with industrial development and climate variability, the pressure on water resources is considerable (Nemer *et al.* 2023). Furthermore, as crustacean biodiversity is essential for maintaining ecosystem functions, it is imperative to study the distribution of organisms and their response to climate change and human disturbance. We have therefore chosen to study the abundance and distribution of crustacean communities in wetlands (Tonga Lake and Birds Lake) classified as Ramsar sites and UNESCO World Heritage Biosphere Reserves (1983, 1999) in El Kala and El Tarf provinces, and in geographical areas rich in flora and fauna such as El Kennar Lake in Jijel province. This work enabled us to identify 8,175 specimens belonging to the freshwater crustacean groups Cladocera (9 taxa), Copepoda (3 taxa), Ostracoda (1 taxon) and Decapoda (1 taxon). Six species (*D. magna*, *D. pulex*, *S. expinosus*, *S. vetulus*, *C. fuscus*, and *Cypris sp*) were found to be common to the three lakes. The *C. brevilabris* species collected from Jijel is a new species in Algerian Cladoceran fauna. Thus, by adding these taxa to the 23 species of Calanoides identified in North Africa to date and the 13 species already known in Algeria (Mouelhi *et al.* 2000), it should be noted that the species *T. affinis* was recorded for the first time in this study.

In their natural environment, crustaceans' communities are generally governed by abiotic factors such as salinity, sediment characteristics, substrate, temperature, availability, water flow, flood frequency and pH (Salleh-Mukri and Shuhaida 2021). Pollution by liquid and solid waste is also an abiotic factor that greatly influences the abundance of crustaceans. Biotic factors are linked to the diversity of associated flora and fauna. In addition, human harvesting contributes to the abundance of certain species of crustaceans (Indarjo *et al.* 2020). Although often small, some crustacean species can be hyper-abundant and dominate the biomass of an ecosystem, such as planktonic copepods, which are considered one of the most abundant metazoans on the planet (Behringer and Duermit-Moreau 2021).

The examined ecological indexes indicated that the three biotopes are somewhat different. According to the abundance of crustacean populations, there is a gradient of decreasing abundance: El Kannar Lake has 3,771 individuals, Bird Lake has 3,745 and Tonga Lake has 659 individuals. Given the number of individuals collected, abundance in Tonga Lake is low, probably due to the characteristics of the habitat such as lack of food, competition, vegetation type, the presence of cyanobacteria, waste and pollution (Djabourabi *et al.* 2017). In addition, floods and droughts might influence zooplankton dynamics by affecting both local (food availability) and regional (increasing connection and dispersion) environmental conditions. Also, the levels of nutrient salts recorded as well as the indicator germs of fecal

contamination are quite important and confirm the state of eutrophication in Tonga Lake (Loucif and Chenchouni 2024). Although El Kannar Lake recorded the highest individual abundance. Furthermore, Cladocerans predominate over the other two lakes, suggesting that the water in El Kannar Lake favors the presence of Cladocerans in contrast to Tonga Lake, which favors the presence of Ostracods and Decapods. Birds Lake is characterized by intermediate dominance. The water degradation of Birds Lake is a reality and represents a serious environmental problem due to domestic wastewater discharges into the lake and intensive livestock farming around the lake (cattle and sheep), as well as very significant fecal contamination expressed by high levels of total coliforms, fecal coliforms and fecal streptococci (Mezbour *et al.* 2018). Moreover, observations made during the sampling period showed that lakes are habitats to more aquatic/macrophytic plants, such as *Nelumbo nucifera*, *Pandanus helicopus*, *Eleocharis dulcis* and *Lepironia articulata*, than artificial lakes, which can provide habitat, food and refuge for crustaceans.

In order to enhance and to visualize the complex relationships of crustacean distributions in their biotope, the NMDS ordination technique of is the most reliable and it is known to be an indirect gradient analysis which creates an ordination based on a dissimilarity or distance. This revealed an overlap between the three communities, which is justified by the presence of common species; this result is supported by the Sørensen index, which reflected the level of similarity between crustacean's communities. These results; indicate that, despite the distance (56 km) separating the two Ramsar protected areas, Tonga Lake and Birds Lake, there is a high degree of connectivity at the functional level between the assemblages due to the large number of shared species and their functional traits. Furthermore, this reveals site specialization, as Birds Lake which is dominated by Copepods (*C. fuscus* 1427 and *Cyclops Sp* 790), El Kannar Lake which is dominated by Cladocerans (*S. expinosus* 2119) and Tonga Lake dominated by Ostracods (*Cypris sp* 266) and Decapods (*A. desmaresti* 141).

In our study, we identified 4 orders of crustaceans: Cladocerans, Copepods, Ostracods and Decapods. This means that the knowledge of the ecological characteristics and dynamics of the populations of these orders is essential. In general, Cladocerans are abundant filter feeders in various biotopes, in small and large freshwater, brackish or saltwater ponds, in the open waters of large lakes, in bottom sediments and among vegetation in the littoral zone of flowing waters of different types (Forró *et al.* 2008). Most Cladocerans may be bacterivorous, herbivorous or detritivorous, feeding on a wide range of elements, including organic detritus (Hayashi-Martins *et al.* 2017). The Cladoceran populations reach their abundance peak in the spring and autumn seasons, and when ecological conditions deteriorate, the males appear (Nandini and Sarma 2019). Cladocerans have the capacity to adapt to the environment, i.e. change their shape according to the season, dissolved oxygen, and the presence of fish. Their abundance can be

associated with a close relationship with the dense presence of aquatic vegetation (Bolduc *et al.* 2016), as they are specialized in exploiting microenvironments provided by vegetation (Forró *et al.* 2008). In our work, Cladocerans' populations such as the Daphniidae family: *D. pulex*, *D. magna*, *Simocephalus expinosus* and *S. vetulus* are common in the investigated lakes. *Ceriodaphnia sp* and *Daphnia atkinsoni* are specific in El Kennar Lake. The Chydoridae family is presented by a single species, *Chydorus brevibris* in El Kennar Lake, and the Moinidae family is presented by *M. micrura* in Bird's Lake. Furthermore, Bouzidi *et al.* (2010) revealed the presence of 4 species of *Ceriodaphnia dubia*, *Ceriodaphnia reticulata*, *Chydorus sp* and *Moina micrura*. This is in agreement with our results, except for the *Diaphanosoma brachyurum* species, which were absent during our study period. These species disappear during the period from July to October periods, probably due to the physicochemical parameters of the water, such as conductivity, salinity and habitat depth. Our results are in agreement with those of Touati. (2008) regarding the spatiotemporal distribution of the genera *Daphnia* and *Simocephalus* in the temporary pools of eastern Numidia (Annaba and El Kala). Conductivity, salinity and water depth are the physicochemical factors influencing the spatial distribution of *Daphnia* and *Simocephalus*. In previous studies, 36 species belonging to 18 genera and 7 families (Sididae, Daphniidae, Bosminidae, Chydoridae, Ilyocryptidae, Moinidae and Macrothricidae) have been identified in Algeria. In addition to the 70 species already known from Algeria, we have now updated the list of 81 Cladoceran taxa known from the country. Among the taxa observed, *Simocephalus vetulus* (O.F. Müller, 1776) is the most frequently encountered species, recorded in 35 different sites, followed by *Daphnia magna* Straus, found in 25 sampling sites (Ghaouaci *et al.* 2018). We also report the occurrence of *Tanymastigites ajjeri* in Algeria, providing an update on the geographical distribution of large branchiopods (Chergui *et al.* 2024). Thirty species of Cladocerans were identified and classified in the families of Sididae, Bosminidae, Chydoridae, Daphniidae, Moinidae and Macrothricidae in various habitats in Morocco (Aoujdad *et al.* 2014). Thirteen Cladoceran species have been recorded throughout Turkey, some for the first time in these localities. The genus *Ceriodaphnia* proved to be the most dominant group, followed by the genera *Daphnia*, *Macrothrix*, *Moina* and *Bosmina* *Simocephalus* (Kaya and Altındağ 2007). This proves that these species are able to acclimatize and adapt to their environment.

Copepods, the second order of crustaceans, are found in almost all freshwater habitats, from the largest ancient lakes to subterranean waters, from pools of glacial meltwater to hot springs, and from hypersaline lakes to phytotelmata. They are extremely abundant in freshwater and comprise a major component of most planktonic, benthic and groundwater communities, including semi-terrestrial situations such as damp moss and leaf litter in humid forests. The largest copepod family is the Cyclopidae which comprises over 800 species (Dussart and Defaye 2006), the great majority belonging to two

freshwater subfamilies, Eucyclopinae and Cyclopinae. Based on the results obtained, the taxonomic richness of the Copepods is demonstrated by the presence of three species: *Cyclops fuscus*, *Cyclops sp* and *Temora affinis*. In general, the faunal composition and temporal distribution of Copepods in Tonga, Birds and El Kannar lakes are similar to the results obtained by various researchers in African lakes (Mahmoudi *et al.* 2022) and on the Atlantic coast of Morocco (El Khalki and Moncef 2007). It is important to note that the predominance of this group is explained by their highly varied diet, which allows them to develop in environments where prey such as protozoa, larvae of other crustaceans and rotifers are available. Houmani *et al.* (2023) revealed that the *Cyclops fuscus* species is highly prevalent in El Chatt Lake in the El Tarf province, whereas *Temora affinis*, which belongs to the suborder Calanoides, is only reported in our study in Tonga Lake. Our results revealed that Copepod had low species diversity, with only 3 taxa found in the samples. Hamil *et al.* (2021) also found 3 taxa (*Arctodiaptomus salinus*, *Copidodiaptomus numidicus* and *Cyclops strenuus*) in the Ghrib Dam the northwestern of Algeria. According to Ghaouaci *et al.* (2017), the most abundant Copepod species in the provinces of Tébessa and Souk Ahras in northeastern Algeria is *Eucyclops leschermoutouae* (Alekseev and Defaye 2004). While the study of the diversity and distribution of zooplankton organisms in the Hassi Ben Abdallah lake in Ouargla, south-east Algeria, showed a total of 5 Copepod taxa: (*Eudiaptomus sp.* (Kiefer, 1932) *Calanus sp.* (Leach, 1819) *Acartia sp.* (Dana, 1846), *Cyclops sp.* (O. F. Müller, 1776) and *Bryocamptus sp.* (Claus, 1863) (Kebabsa *et al.* 2019). The analysis of available information on diaptomid copepods in Tunisian inland waters revealed a total of 10 diaptomid species belonging to 7 genera and 2 subfamilies. This biodiversity is therefore strongly linked to the diversity of inland waters that characterize the country (Marrone *et al.* 2023).

As for Ostracods, they are widely distributed in all types of aquatic habitat, fresh, brackish and salt water (Meisch 2000; Altınsağılı *et al.* 2020). The efficient colonization of their habitat can be attributed to several factors, such as their different modes of reproduction (parthenogenetism, amphigonism or both) (Ozawa 2013) or their relatively high ecological tolerances to different environmental conditions (Martens and Savatnalinton 2011). Previous studies of Ostracods in Algeria Samraoui *et al.* (1998), who gave lists of species of Rotifers, Cladocerans, Copepods and 6 Ostracods from Numidia, in the north-east of the country. In the south-western Algerian Sahara (Tindouf), Karanovic and Pesce. (2000) described a new species, *Martenscypridopsis materia*, found in inland waters. The first in-depth research on Ostracods in Algeria was undertaken by Ghaouaci *et al.* (2015), (2017) who provided a recent list of non-marine Ostracod taxa. In our case, Ostracods (*Cypris sp*) are represented in negligible proportions compared to the other orders (5.16%) of the total fauna. It is probable that the main factors limiting the distribution of these species are the low content of ions in the water, as mentioned by Iglowska and

Namiotko (2012). This means that the distribution of ostracod species is closely linked to the physicochemical characteristics of the waters (Perçin-Paçal *et al.* 2017). Furthermore, the research conducted by Saoudi *et al.* (2018) on the aquatic macroinvertebrates of the El Mekhada marsh in northeastern Algeria has identified species of Ostracods belonging to the Cyprididae family. Similarly, this study indicates that the physicochemical parameters of freshwater show a higher abundance of taxa during the wet season compared with the dry season, which is consistent with our results. Seasonal changes in water quality also have a direct influence on Ostracod species diversity. Additionally, previous work by Kulköylüoğlu *et al.* (2021), as well as Yavuzatmaca. (2020), presented evidence supporting the preference of these species for certain habitat conditions, particularly freshwater. Their tolerance is low, and they have optimum values, with relatively high abundance in November and December. Thus, winter offers more stable conditions compared with summer (Kulköylüoğlu *et al.* 2022). Over the 2002–2012 period, one hundred and five sites in Tunisia, covering different climatic zones from the Mediterranean to the desert, were sampled. Indeed, based on previous investigations, 9 families (Candonidae, Cyprididae, Cytherideidae, Darwinulidae, Ilyocyprididae, Leptocytheridae, Limnocytheridae, Loxoconchidae and Paradoxostomatidae), 29 genera and at least 45 species of non-marine ostracods are currently known (Marrone *et al.* 2020). For some Ostracod species, clear distribution gradients associated with different climatic conditions have been observed (Marrone *et al.* 2020). In northeastern Italy (Friuli Venezia Giulia), 74 taxa from 30 genera belonging to 9 different families (Darwinulidae, Candonidae, Ilyocyprididae, Notodromadidae, Cyprididae, Limnocytheridae, Cytheridae, Leptocytheridae and Xestoleberidae) were identified. The most common species is *Cypria ophthalmica* (133 records), followed by *Cyclocypris ovum* (86 records), *C. laevis* (49 records), *Cypridopsis vidua* (40 records), and *Notodromas persica* (28 records) (Pieri *et al.* 2009). Until now, 145 non-marine ostracod species have been known from Turkey (Kulköylüoğlu *et al.* 2015). There were a total of 1,787 individuals belonging to 31 species recorded in all the basins of Konya, Antalya and the western Mediterranean in Turkey. Some of these species are common to all the basins, while others are new, such as *Fabaeformiscandona fragilis*, which is new to all the basins. Nine species were found to be common to all in Konya, Antalya and West Mediterranean basins in Turkey (Yavuzatmaca 2019).

Decapods are represented by the species *A. desmarestii* (Millet, 1831), which belongs to the family Atyidae. It is a freshwater shrimp found in North Africa and Southern Europe. It is generally found in the Mediterranean region. In Algeria, Ameur *et al.* (2022) revealed the presence of this species in Oubeira and Tonga lakes and in the mouth of the Mafragh River on the coastal basin of El Battah in Annaba's Gulf. The works of Mabrouki *et al.* (2018) in Morocco and Dhaouadi *et al.* (2009) in Tunisia provided evidence of the presence of this species in the Mediterranean basin. The same family of freshwater

shrimp is represented by the species *Atyoida serrata* according to the inventory of Valade (2003) in Mauritius. On the other hand, the work of Chevalier and Clavier. (2020) in French Guiana on 4 species of shrimp collected in the Haut Courcibo area shows that all the species belong to the Palaemonidae family, with the absence of the Atyidae family. Another research conducted by Dhaouadi. (2003) identified a new southern limit for this shrimp in the El Haoureb dam in central Tunisia and at three other dams, Sidi Salem, Lebna and Sidi Saâd. The distance between these three reservoirs means that they belong to three distinct bioclimatic stages (Dhaouadi *et al.* 2004). The Decapod from Greece, Holthuis. (1961) reported the species from Lake Koronia as *A. desmarestii*. Recent sampling, covering a dense station network of the Greek freshwaters, revealed the presence of *A. desmarestii* in the western and northern part of the mainland (Anastasiadou *et al.* 2004). The information obtained shows that *A. desmarestii* has been found in many Portuguese watercourses, including reservoirs, rice fields, coastal lagoons and temporary streams. Despite the ecological relevance of this shrimp in aquatic communities, scientific information about its distribution in the country is still fragmentary and scattered (Fidalgo and Gerhardt 2002).

Physicochemical factors are the main drivers of crustacean community composition, diversity and spatiotemporal distribution. Temperature is a fundamental factor in climate; its variability affects evaporation and evapotranspiration and, consequently, the salinity level of the water. This parameter depends on the altitude and distance from the sea and also varies with the seasons (Kumar 2012). Seasonal precipitation is important because it allows us to assess the variation in precipitation and its trend according to the different seasons of the year. It is an ecologically important factor, not only for the functioning and distribution of limnic ecosystems (Ramade 2003). The annual precipitation of the Mediterranean area is expected to show a decrease of about 4% for every 1°C increase in global temperature. This decrease is expected mainly in the summer season for the northern regions of the Mediterranean Basin and across all seasons in the central and southern regions (Ali *et al.* 2022).

Apart from the changes in climate conditions, another important aspect of global warming is the increase in the frequency and intensity of extreme weather events (Lazoglou *et al.* 2024). Physicochemical factors spatially structure communities in aquatic environments (Allan *et al.* 2021) and contribute to the local and regional diversity of aquatic fauna and flora. For instance, at the regional scale, communities are influenced by factors such as dissolved oxygen, temperature and salinity, which usually vary from one aquatic habitat to another (Figuerola and De los Rios-Escalante 2021). Environmental factors also change in time, particularly in temperate regions (e.g., from one season to another), influencing the life cycle of aquatic organisms and determining their abundance and community composition across time (Smeti *et al.* 2019). In

addition, anthropogenic factors tend to change the physicochemical characteristics of water, influencing the structure and diversity of aquatic organisms (Fergani and Arab 2013). Physical (temperature, lake size) and chemical (nutrients, metals, and chloride) parameters of freshwater ecosystems have significant effects on crustacean species composition. In our case, decapod species (*A. desmarestii*) are more abundant during the month with the highest rainfall. Conversely, low rainfall or low water levels would reduce oxygenation and, consequently, the metabolic rate of these organisms. Cladoceran (*Ceriodaphnia*, *Daphnia*, *Simocephalus*, *Chydorus*, *Moina*) populations reach peak abundance under optimal thermal conditions, typically between 18 and 22°C, and in waters characterized by neutral to slightly alkaline pH, as recorded in Lake El Kennar. In contrast, copepods (*Cyclops*, *Temora*) exhibit a broader ecological tolerance, persisting at higher pH levels, up to 8.5. Ostracode (*Cypris sp*) their relatively high ecological tolerances to different environmental conditions, our results indicates that the physicochemical parameters of freshwater show a higher abundance of taxa during the month with the highest rainfall, which is consistent with Kulköylüoğlu *et al.* (2021), as well as Yavuzatmaca. (2020). There is an urgent need for effective conservation and adaptive management strategies, including temperature regulation, control of nutrient pollution, and the preservation of habitat connectivity. Implementing such measures can help mitigate biodiversity loss in freshwater ecosystems and enhance their resilience to the impacts of ongoing climate change (Kasinathan *et al.* 2024). Understanding the seasonal changes in physicochemical factors is crucial to determine the factors that drive the abundance and composition of aquatic communities.

This means that crustaceans such as Ostracods, Cladocera, Copepods and Decapods are very sensitive to any type of climate change or to modifications in the physicochemical parameters of the environment in which they live. Any change in these factors can influence their distribution, abundance and growth process. This is proved by the problem of global warming and the record number of droughts that the world has experienced in recent years, according to the World Meteorological Organisation (WMO April 2021).

5. Conclusion

In this study, our observations underline that freshwater crustacean species diversity and distribution could be significantly correlated with environmental variables and climatic data related to their biotopes. We recorded 14 species which match 7 families: Daphniidae, Cyclopidae, Cyprididae, Atyidae, Chydoridae, Temoridae and Moinidae. Furthermore, NMDS analysis showed a spatial effect in the crustacean communities and highlight three species assemblages. We have also observed several species common to all three lakes, some are more abundant and others are rarely found or newly identified in Algeria. The taxonomic composition of crustacean communities, in Tonga, Birds and El Kennar lakes, react to variation in geographic gradients and are likely to be

endangered due to habitat alterations such as increasing temperatures in Algeria and nutrient pollution.

Altogether, our results also emphasize that crustaceans' biodiversity and species identification are an essential contribution to a better understanding of crustacean communities threatened by climate change and anthropogenic activities. Integrating fauna's biological and ecotoxicological assessments is recommended to conserve wildlife and restore natural habitats, and to upgrade and to enhance public awareness toward the importance of freshwater biotopes. This study provides one of the few quantitative assessments of freshwater crustacean communities. It helps to fill the knowledge gap on aquatic biodiversity in Algeria. Future research should target unexplored regions (Saharan areas, high plateaus, temporary wetlands). Comparing data provides a valuable framework for understanding biogeographical dynamics and the impact of environmental changes on freshwater biodiversity.

6. Author Contributions

Conceptualisation: BERGHICHE Hinda, BAROUR Choukri. Developing methods: BERGHICHE Hinda, BAROUR Choukri. Conducting the research: BALI Imene BERGHICHE Hinda, BAROUR Choukri. Identification of species: BALI Imene, BENDALI-SAUDI Fatiha Data analysis, Data interpretation: BAROUR Choukri, BALI Imene, BERGHICHE Hinda Preparation of figures and tables: BAROUR Choukri, BALI Imene Writing: BERGHICHE Hinda, BALI Imene, BAROUR Choukri.

7. Acknowledgements

This work was supported by the National Fund for Scientific Research to Pr. H. Boudjelida, University of Badji Mokhtar Annaba "Laboratory of Applied Animal Biology", and the Ministry of High Education and Scientific Research of Algeria (University Research-Treaining Project to Prof. S. Chouahda).

8. Conflicts of Interest

The authors declare no conflicts of interest.

References

- Alekseev V.R., & Defaye D. (2004), *Eucyclops leschermoutouae* sp. n. from Majorca, Balearic Islands (Crustacea, Copepoda, Cyclopidae), *Zoosystematica Rossica*, 12 (2): 163–169.
- Ali E., Cramer W., Carnicer J., Georgopoulou E., Hilmi N.J.M., Le Cozannet G. & Lionello P. (2022), Cross-chapter paper 4: Mediterranean region, *Climate change*, pp.2233-2272.
- Allan J. D., Castillo M. M., & Capps K. A. (2021), Stream ecology: structure and function of running waters (3rd ed. Springer Nature), Switzerland. <https://doi.org/10.1007/978-3-030-61286-3>
- Altınışağı S., Perçin-Paçal, F., Altınışağı S. (2020) Ecological interactions and the distribution, abundance, and diversity of ostracod species associated with some coastal brackish karstic springs in southwest Turkey, *Regional Studies in Marine Science*, 35, 2352-4855, <https://doi.org/10.1016/j.rsma.2020.101204>.
- Ameur A., Berghiche H., Barour C., & Soltani N. (2022), Morphometric reproductive and environmental biomarker analysis of a crustacean species *Atyaephyra desmarestii* (Millet, 1831) from north east Algeria, *Applied Ecology and Environmental Research*, 20(6), 4579-4597. http://dx.doi.org/10.15666/aeer/2006_45794597
- Anastasiadou C., Koukouras A., Mavidis M., Chartosia N., Mostakim M., Christodoulou M., & Aslanoglou C. (2004), Morphological variation in *Atyaephyra desmarestii* (Millet, 1831) within and among populations over its geographical range, *Mediterranean Marine Science*, 5(2), 5–14. <https://doi.org/10.12681/mms.198>
- Anderson M. J., Crist T. O., Chase J. M., Vellend M., Inouye B. D., Freestone A. L., & Swenson N. G. (2011), Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist, *Ecology letters*, 14(1), 19–28. <https://doi.org/10.1111/j.1461-0248.2010.01552.x>
- Aoujdad R., Maqboul A., Fadli M., & Fekhaoui M. (2014), Structure and organization of the crustaceans cladoceran populations in Moroccan rice fields, *J Entomol Zool Stud*, 2(6), 39-44.
- Ayati K., Hadjab R., Khammar H., Dhaouadi S., Piscart C., & Mahmoudi E. (2019), Origin, diversity and distribution of freshwater epigeal amphipods in Maghreb, In *Annales de Limnologie - International Journal of Limnology*, 55(13). EDP Sciences.
- Ayati K., Hupało K., Dhaouadi S., *et al.* (2023), Conservation of freshwater biodiversity in Tunisia in a climate change context: combining amphipod distribution data and molecular analyses to improve priorities, *Biodivers Conserv*, 32, 2539–2559. <https://doi.org/10.1007/s10531-023-02617-8>
- Azra M., Mohd Noor M., Eales J., Yik Sung Y., & Abd Ghafar M. (2022), What evidence exists for the impact of climate change on the physiology and behaviour of important aquaculture marine crustacean species in Asia? A systematic map protocol, *Environmental Evidence*, 11:9. <https://doi.org/10.1186/s13750-022-00263-1>
- Barrett K., Guyer C., Samoray S. T., Kanno Y. (2016), Stream and Riparian Habitat Use by Anurans along a Forested Gradient in Western Georgia, USA, *Copeia*, 104 (2): 570–576. <https://doi.org/10.1643/CE-15-318>
- Beaugrand G., Conversi A., Chiba S., Edwards M., Fonda-Umani S., Greene C., ... & Sugisaki H. (2015), Synchronous marine pelagic regime shifts in the Northern Hemisphere, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 370 (1659), 20130272. <https://doi.org/10.1098/rstb.2013.0272>
- Behringer D. C., & Duermit-Moreau E. (2021), Crustaceans, One Health and the changing ocean, *Journal of invertebrate pathology*, 186, 107500. <https://doi.org/10.1016/j.jip.2020.107500>
- Błędzki L. A., Rybak J. I. (2016), Freshwater crustacean zooplankton of Europe: Cladocera & Copepoda (Calanoida, Cyclopoida) key to species identification, with notes on ecology, distribution, methods and introduction to data analysis, Springer, Cham, Switzerland.
- Bolduc P., Bertolo A., & Pinel-Alloul B. (2016), Does submerged aquatic vegetation shape zooplankton community structure and functional diversity? A test with a shallow fluvial lake

- system, *Hydrobiologia*, 778, 151-165. <https://doi.org/10.1007/s10750-016-2663-4>
- Bouzidi M. A., Amar Y., Attaoui I., Latrèche A., Benyahia M., Bouguenaya N., & Meliani H. (2010), Copépodes, Cladoceres et Rotifères du lac Sidi M'hamed Benali (Algérie nord-occidentale), *Physio-Géo. Géographie physique et environnement*, (Volume 4), 69-85.
- Brooks T. M., Mittermeier R. A., Da Fonseca G. A., Gerlach J., Hoffmann M., Lamoreux J. F., Mittermeier C. G., Pilgrim D. & Rodrigues A. S. (2006), Global biodiversity conservation priorities, *Science*, 313(5783), 58-61. <https://doi.org/10.1126/science.1127609>
- Cañadas E.M., Fenu G., Peñas J., Lorite J., Mattana E., & Bacchetta G. (2014), Hotspots within hotspots: endemic plant richness, environmental drivers, and implications for conservation, *Biol Conserv*, 170:282–291. <https://doi.org/10.1016/j.biocon.2013.12.007>
- Cantonati M., Poikane S., Pringle C.M., Stevens L.E., Turak E., Heino J., Richardson J.S., Bolpagni R., Borriani A., Cid N., Čvrtlíková M., Galassi D.M.P., Hájek M., Hawes I., Levkov Z., Naselli-Flores L., Saber A.A., Cicco M.D., Fiasca B., Hamilton P.B., Kubečka J., Segadelli S., & Znachor P. (2020), Characteristics, main impacts, and stewardship of natural and artificial freshwater environments: consequences for biodiversity conservation, *Water*, 12(1): 260. <https://doi.org/10.3390/w12010260>
- Chevalier J., & Clavier S. (2020), Les crevettes de la réserve naturelle régionale Trésor: rapport de mission et perspectives – Rapport d'étude, 32p.
- Chergui I., Koulali K., Bouzid A., & Samraoui B. (2024), First record of Tanyastigites ajjeri Thiéry & Rogers 2022 (Crustacea: Branchiopoda: Anostraca) in Algeria, with an updated checklist of Algerian large branchiopods, *Zoology and Ecology* 34(1), 1–8. <https://doi.org/10.35513/21658005.2024.1.1>
- Cramer W., Guiot J., & Marini K. (2022), Energy transition in the Mediterranean. In: Climate and Environmental Change in the Mediterranean Basin—Current Situation and Risks for the Future, First Mediterranean Assessment Report.
- De Deckker P. (1981), Ostracods of athalassic saline lakes, a review, *Hydrobiologia*, 81, 131–144. <https://doi.org/10.1007/BF00048710>
- Dhaouadi-Hassen S. (2003), Etude écobioécologique d'Atyaephyra desmaresti (Millet, 1834) (Crustacea, Decapoda, Natantia, Atyidae) de trois retenues de barrages de Tunisie: Sidi Salem, Lebna et Sidi Saâd, Doctoral thesis, University of Tunis El Manar, Tunisia.
- Dhaouadi-Hassen S., & Boumaïza M. (2009), Reproduction and population dynamics of Atyaephyra desmarestii (Decapoda, Caridea) from the Sidi Salem dam lake (northern Tunisia), *Crustaceana*, 129-139. <http://www.jstor.org/stable/27743267>
- Dhaouadi-Hassen S., Trilles J. P., Charmantier G., & Boumaïza M. (2004), Ecophysiologie d'une crevette d'eau douce, Atyaephyra desmarestii (Millet, 1831) (Crustacea, Decapoda) dans trois barrages tunisiens: résultats préliminaires, *Zool. baetica*, 15, 175-183.
- Dinno A. (2024), dunn.test: Dunn's test of multiple comparisons using rank sums, R package version 1.3.6. <https://CRAN.R-project.org/package=dunn.test>.
- Djabourabi A., Touati H., Sehili N., Boussadia M. I., & Bensouilah M. (2017), Study of the physicochemical parameters of water and phytoplankton in Lake Tonga (wetland of the national park of El Kala, North East of Algeria), *International Journal of Biosciences*, 11(3), 213-226. <http://dx.doi.org/10.12692/ijb/11.3.213-226>
- Dussart B. H., & Defaye D. (2006), World directory of Crustacea Copepoda of Inland Waters. II-Cyclopiformes, Leiden: Backhuys Publishers. 354 pp.
- El Khalki A., & Moncef M. (2007), Study of the copepods population in the Oum Er Rbia estuary (Atlantic Moroccan coast): tides and reservoir release effects, *Lebanese Science Journal*, 3-18.
- Fergani H., & Arab A. (2013), Utilisation des macroinvertébrés benthiques comme bioindicateurs de pollution d'Oued El-Harrach-FBS 4th International Congress of the Populations et Animal Communities, *Dynamics & Biodiversity of the terrestrial & aquatic Ecosystems CIPCA4. TAGHIT (Bechar), Algeria*.
- Fidalgo M. L., & Gerhardt A. (2002), Distribution of the freshwater shrimp, Atyaephyra desmarestii (Millet, 1831) in Portugal (Decapoda, Natantia), *Crustaceana*, 1375-1385. <https://www.jstor.org/stable/20105526>
- Figuerola D., & Rios-Escalante P. (2021), Macrozoobenthos in an altitudinal gradient in North Patagonian Cautín river (Araucanía region, Chile), *Brazilian Journal of Biology*, 82, e240484. <https://doi.org/10.1590/1519-6984.240484>
- Fischer W., Schneider M., & Bauchot M. L. (1987), Méditerranée et mer noire zone de pêche 37 Révision 1 Volume 1 végétaux et invertébrés. Rome.
- Forro L., Korovchinsky N. M., Kotov A. A., & Petrusek A. (2008), Global diversity of cladocerans (Cladocera; Crustacea) in freshwater, *Freshwater animal diversity assessment*, 177-184. https://doi.org/10.1007/978-1-4020-8259-7_19
- Gauthier H. & Brehm V. (1928), Ostracodes et Cladocères de l'Afrique du Nord (3e note), *Bulletin de la Société de l'Histoire Naturelle de l'Afrique du Nord*, 19, 114–121.
- Gauthier H. (1928a), Recherche sur la faune des eaux continentales de l'Algérie et de la Tunisie, Imprimerie Minerva, Alger, 419 pp.
- Gauthier H. (1928b), Ostracodes et Cladocères de l'Afrique du Nord (Première note), *Bulletin de la Société de l'Histoire Naturelle de l'Afrique du Nord*, 19, 10–19.
- Gauthier H. (1928c), Ostracodes et Cladocères de l'Afrique du Nord (2e note), *Bulletin de la Société de l'Histoire Naturelle de l'Afrique du Nord*, 19, 69–79.
- Ghaouaci S., Amarouayache M., Sinev A. Y., Korovchinsky N. M., & Kotov A. A. (2018), An annotated checklist of the Algerian Cladocera (Crustacea: Branchiopoda), *Zootaxa*, 4377(3), 412-430. <https://doi.org/10.11646/zootaxa.4377.3.5>
- Ghaouaci S., Yavuzatmaca M., Külköylüoğlu O., & Amarouayache M. (2017), An annotated checklist of the non-marine ostracods (Crustacea) of Algeria with some ecological notes, *Zootaxa*, 4290(1), 140-154. <https://doi.org/10.11646/zootaxa.4290.1.8>
- Ghaouaci S., Yavuzatmaca M., Külköylüoğlu O., Amarouayache M. & Gouasmia G. (2015), Checklist of the living non-marine Ostracoda (Crustacea) of Algeria, In: 8th European Ostracod Meeting, Tartu, Estonia, July 22–30, 2015, pp. 27.
- Gonzalez C.G. (2023), EcolndR: Ecological indicators, R package version 2.0, <https://CRAN.R-project.org/package=EcolndR>.
- Gouws G., Stewart B. A., & Daniels S. R. (2010), Phylogeographic structure in the gilgie (Decapoda: Parastacidae: Cherax

- quinquecarinatus): a south-western Australian freshwater crayfish, *Biological Journal of the Linnean Society*, 101(2), 385-402. <https://doi.org/10.1111/j.1095-8312.2010.01485.x>
- Hamil S., Bouchelouche D., Arab S., Aili M., Baha M., & Arab A. (2021), The relationship between zooplankton community and environmental factors of Ghrib Dam in Algeria, *Environmental Science and Pollution Research*, 28, 46592-46602. <https://doi.org/10.1007/s11356-020-10844-7>
- Harrell Jr. F. (2025), Hmisc: Harrell Miscellaneous, R package version 5.2-2, <https://CRAN.R-project.org/package=Hmisc>.
- Hayashi-Martins, L. H., Mansano A. S., Hisatugo K. F., Rocha O., & Selegim M. H. R. (2017), "In vitro" evaluation of the bacterivore potential of three Cladoceran species occurring in tropical and subtropical regions, *Brazilian Journal of Biology*, 77, 840-847. <https://doi.org/10.1590/1519-6984.02716>
- Holthuis L. B. (1961), Report on a collection of Crustacea Decapoda and Stomatopoda from Turkey and Balkans, *Zoologische Verhandelingen Leiden*, 47: 1-67
- Houmani M., Bendali-Saoudi F., & Soltani N. (2023), Impact of physicochemical parameters of water on the biodiversity of the invertebrate fauna from Echatt Lake, Northeast Algeria, *Biodiversitas Journal of Biological Diversity*, 24.1. <https://doi.org/10.13057/biodiv/d240131>
- Iglikowska A., & Namiotko T. (2012), The impact of environment factors on diversity of ostracoda in freshwater habitats of subarctic temperate Europe, *Annales Zoologici Fennici*, 49(4), 193-218. <https://doi.org/10.5735/086.049.0401>
- Ihaka R., & Gentleman R. (1996), R: a language for data analysis and graphics, *Computational and Graphical Statistics*, (5) 299-314.
- Indarjo A., Salim G., Zein M., Septian D., & Bija S. (2020), The population and mortality characteristics of mangrove crab (*Scylla serrata*) in the mangrove ecosystem of Tarakan City, Indonesia, *Biodiversitas Journal of Biological Diversity*, 21(8). <https://doi.org/10.13057/biodiv/d210855>
- Karanovic I., & Pesce G.L. (2000), Martenscypridopsis a new ostracod genus (Crustacea: Ostracoda) from African inland waters, *Annals of Limnology*, 36 (3), 149-155. <https://doi.org/10.1051/limn/2000012>
- Kasinathan P. et al. (2024), Evaluating Water Quality through Macroinvertebrate Diversity and Physicochemical Parameters in the Thamirabarani River Basin, *Global NEST Journal*, 26(10), Available at: <https://doi.org/10.30955/gnj.06044>.
- Kassambara A. (2023a), ggcorrplot: Visualization of a correlation matrix using 'ggplot2'. R package version 0.1.4.1, <https://CRAN.R-project.org/package=ggcorrplot>.
- Kassambara A. (2023b), ggpubr: 'ggplot2' Based publication ready plots, R package version 0.6.0, <https://CRAN.R-project.org/package=ggpubr>.
- Kaya M., & Altındağ A. (2007), Some cladoceran species from Turkish inland waters, *Süleyman Demirel University Faculty of Arts and Science Journal of Science*, 2(1), 60-76.
- Kebabza R., Manamani R., & Djerad H. (2019), Zooplankton du lac de Hassi ben Abdallah (Ouargla, Sud-Est Algérie), *Algerian journal of arid environment (AJAE)*, vol. 9, n°2.
- Krepski T., Sługocki Ł., Goździk I., Humiczewski M., Popko R., & Czerniawski R. (2024), Spatial Distribution Patterns of Zooplankton and Macroinvertebrates in a Small River under Strong Anthropogenic Pressure, *Water*, 16(2), 262. <https://doi.org/10.3390/w16020262>
- Külköylüoğlu O., Akdemir D., Yavuzatmaca M. & Yılmaz O. (2015), A checklist of recent non-marine Ostracoda (Crustacea) of Turkey with three new records, *Review of Hydrobiology*, 8/2, 77-90.
- Külköylüoğlu O., Çelikbaş B., & Ataman A. (2022), Habitat destruction in wetland affects Ostracoda (Crustacea) species occurrence patterns amid different aquatic habitats, *Aquatic Ecology*, 56(3), 605-618. <https://doi.org/10.1007/s10452-021-09934-6>
- Külköylüoğlu O., Yavuzatmaca M., & Müller-Hofstede Akdemir D. (2021), Occurrence patterns, photoperiod and dispersion ability of the non-marine Ostracoda (Crustacea) in shallow waters, *Turkish Journal of Fisheries and Aquatic Sciences*, 21(2). http://doi.org/10.4194/1303-2712-v21_2_03
- Kumar C. P. (2012), Climate change and its impact on groundwater resources, *International Journal of Engineering and Science*, 1(5), 43-60.
- Lazoglou G., & Papadopoulos-Zachos A., Georgiades P. et al. (2024), Identification of climate change hotspots in the Mediterranean, *Sci Rep*, 14, 29817. <https://doi.org/10.1038/s41598-024-80139-1>
- Lomartire S., Marques J-C., & Gonçalves A. M-M. (2021), The key of zooplankton ecosystem services: A perspective of interaction between zooplankton and fish recruitment, *Ecological indicators*, Volume 129, 107867, ISSN 1470-160X, <https://doi.org/10.1016/j.ecolind.2021.107867>
- Loucif K., & Chenchouni H. (2024), Water physicochemical quality as driver of spatial and temporal patterns of microbial community composition in lake ecosystems, *Appl Water Sci*, 14, 115. <https://doi.org/10.1007/s13201-024-02176-5>
- Lucas H. (1849), Histoire naturelle des Animaux Articulés. Exploration scientifique de l'Algérie pendant les années 1840, 1841, 1842, *Sciences Physiques Zoologie*, 1, 1-403.
- Mabrouki Y., Taybi A. F., Berrahou A., Chaabane K., & Legssyer B. (2018), Case study of the freshwater shrimp *Atyaephyra desmarestii* (Millet, 1831) (Crustacea, Decapoda) in the watershed of Moulouya and eastern Morocco, *Vie et milieu-life and environment*, 68(4), 175-183.
- Mack H. R., Conroy J. D., Blocksom K. A., Stein R. A., & Ludsins S. A. (2012), A comparative analysis of zooplankton field collection and sample enumeration methods, *Limnol Oceanogr Methods*, 10:41-53. <https://doi.org/10.4319/lom.2012.10.41>
- Mahmood Ghafor I. (2020), Crustacean. Chapitre 1. In: *Crustacea*. (Eds: G. Diarte-Plata & R. Escamilla-Mentes) IntechOpen. <https://doi.org/10.5772/intechopen.89730>
- Mahmoudi K., Bendali-Saoudi F., & Soltani N. (2022), Potential predation of *Cyclops fuscus* Jurine, 1820 (Crustacea Copepoda) against two mosquito species *Culex pipiens* Linnaeus, 1758 and *Culiseta longiareolata* Macquart, 1838 (Diptera Culicidae), *Jour*, 4, 812. <https://doi.org/10.31396/Biodiv.Jour.2022.13.4.805.812>
- Marrone F., Pieri V., Turki S., & Rossetti G. (2020), The Recent non-marine ostracods of Tunisia: an updated checklist with remarks on their regional distribution patterns and ecological preferences, *Journal of Limnology*, 79(3), 293-307. <https://doi.org/10.4081/jlimnol.2020.1982>

- Marrone F., Stoch F., Turki S., & Naselli-Flores L. (2023), The Diaptomidae (Copepoda, Calanoida) of Tunisia and the role of spatial and environmental factors as drivers of their distribution patterns, *Hydrobiologia*, 850(21), 4815-4829. <https://doi.org/10.1007/s10750-023-05173-y>
- Martens K., & Savatnalinton S. (2011), A subjective checklist of the Recent, free-living, non-marine Ostracoda (Crustacea), *Zootaxa*, 2855(1), 1-79. <https://doi.org/10.11646/zootaxa.2855.1.1>
- Meisch C. (2000), Freshwater Ostracoda of Western and Central Europe. Spektrum Akademischer Verlag GmbH, Heidelberg, Berlin, 522 p.
- Mezbour R., Reggam A., Maazi M. C., & Houhamdi M. (2018), Evaluation of organic pollution index and the bacteriological quality of the water of the Lake of birds (ELTarf East-Algerian), *J Mater Environ Sci*, 9(3), 971-979. <https://doi.org/10.26872/jmes.2018.9.3.108>
- Mittermeier R. A., Robles Gil P., Hoffmann M., Pilgrim J., Brooks T., Mittermeier C. G., Lamoreux J. & Da Fonseca G. A. B. (2004), Hotspots revisited: Earth's biologically richest and most endangered terrestrial ecoregions, Preface by Peter A. Seligmann, Foreword by Harrison Ford, Cemex/Conservation International/Agrupacion, *Sierra Madre/Monterrey/Mexico*, 392.
- Moniez R. (1891), Faune des lacs salés d'Algérie. Mémoires de la société zoologique de France, 1891, 246-257.
- Moon K. (2020), webr: Data and functions for web-based analysis. R package version 0.1.5, <https://CRAN.R-project.org/package=webr>.
- Mouelhi S., Balvay G., & Kraiem M. M. (2000), Branchiopodes (Cténopodes et Anomopodes) et Copépodes des eaux continentales d'Afrique du Nord: inventaire et biodiversité, *ZOOSYSTEMA-PARIS*, 22(4), 731-748.
- Myers N., Mittermeier R. A., Mittermeier C. G., Da Fonseca G. A., & Kent J. (2000), Biodiversity hotspots for conservation priorities, *Nature*, 403(6772), 853-858.
- Nandini S., & Sarma S. S. S. (2019), Reproductive strategies of Moina (Cladocera) in relation to their habitat, *Limnetica*, 38(1), 137-145. <https://doi.org/10.23818/limn.38.15>
- Nemer Z., Khaldouy F., Benaissa Z., Belaroui A., Tebbouche M. Y., & Ydri A. (2023), Hydrogeophysical investigation of aquifer parameters and seawater intrusion: a case study from Eastern Mitidja plain, Algeria, *Geomechanics and Geophysics for Geo-Energy and Geo-Resources*, 9(1), 60. <https://doi.org/10.1007/s40948-023-00610-7>
- Oksanen J., Simpson G., Blanchet F., Kindt R., Legendre P., Minchin P., O'Hara R., Solymos P., Stevens M., Szoecs E., Wagner H., Barbour M., Bedward M., Bolker B., Borcard D., Carvalho G., Chirico M., De Caceres M., Durand S, Evangelista H., FitzJohn R., Friendly M., Furneaux B., Hannigan G., Hill M., Lahti L., McGlinn D., Ouellette M., Ribeiro Cunha E, Smith T., Stier A., Ter Braak C., Weedon J., & Borman T. (2025), vegan: Community Ecology Package, R package version 2.6-10, <https://CRAN.R-project.org/package=vegan>.
- Ozawa H. (2013), The history of sexual dimorphism in Ostracoda (Arthropoda, Crustacea) since the Palaeozoic. Chapitre 4. In: Sexual Dimorphism. (Eds: Hiroshi Moriyama), *Intech, Janeza Trdine 9,51000 Rijeka, Croatia*.
- Perçin-Paçal F., Altınışli S., & Altınışli S. (2017), Distribution, diversity and some ecological characteristics of ostracods (Crustacea: Ostracoda) in Gökceada (Imbros) island (northern Aegean Sea, Turkey), *Biologia*, 72(7), 774-789. <https://doi.org/10.1515/biolog-2017-0088>
- Pieri V., Martens K., Stoch F., & Rossetti G. (2009), Distribution and ecology of non-marine ostracods (Crustacea, Ostracoda) from Friuli Venezia Giulia (NE Italy), *Journal of limnology*, 68(1), 1.
- Pimm S.L., & Raven P.H. (2020), Norman Myers (1934-2019), *Nat Ecol Evol* 4, 177-178. <https://doi.org/10.1038/s41559-020-1095-8>
- Pohlert T. (2024), PMCMRplus: Calculate Pairwise Multiple Comparisons of Mean Rank Sums Extended, R package version 1.9.12, <https://CRAN.R-project.org/package=PMCMRplus>.
- Poole G. C., & Berman C.H. (2001), Pathways of human influence on water temperature dynamics in stream channels, *Environ. Manage.* 27 787-802.
- Posit team (2024), RStudio: Integrated development environment for R. Posit Software, PBC, Boston, MA. URL <http://www.posit.co/>.
- R Core Team (2024), R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>
- Ramade F. (2003), Eléments d'écologie, Ecologie fondamentale. Ed. Dunod, Paris, 690 p.
- Revelle W. (2024), psych: Procedures for psychological, psychometric, and personality research. Northwestern University, Evanston, Illinois, R package version 2.4.12, <https://CRAN.R-project.org/package=psych>.
- Rifai S. W., Girardin C. A., Berenguer E., del Aguila-Pasquel J., Dahlsjö C. A., Doughty C. E., & Malhi Y. (2018), ENSO Drives interannual variation of forest woody growth across the tropics, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1760), 20170410. <https://doi.org/10.1098/rstb.2017.0410>
- Rolls R. J., Heino J., Ryder D. S., Chessman B. C., Growns I. O., Thompson R. M., & Gido K. B. (2018), Scaling biodiversity responses to hydrological regimes, *Biological Reviews*, 93(2), 971-995. <https://doi.org/10.1111/brv.12381>
- Salleh-Mukri M., & Shuhaida S. (2021), The habitat preference of mangrove crabs in different mangrove forests of Penang, North Peninsular Malaysia. In *IOP Conference Series: Earth and Environmental Science* (Vol. 736, No. 1, p. 012062). IOP Publishing. <https://doi.org/10.1088/1755-1315/736/1/012062>
- Samraoui B., Segers H., Maas S., Baribwegure D. & Dumont H.J. (1998), Rotifera, Cladocera, Copepoda, and Ostracoda from coastal wetlands in northeast Algeria, *Hydrobiologia*, 386, 183-193.
- Saoudi M., Tadjine A., Guerfi S., & Necer A. (2018), A preliminary survey of water physico-chemical characteristics and aquatic macroinvertebrate communities in El Mekhada marsh, north-eastern Algeria, *Zoology and Ecology*, 28(2), 109-116. <https://doi.org/10.1080/21658005.2018.1462926>
- Schwentner M., Combosch D. J., Nelson J. P., & Giribet G. (2017), A phylogenomic solution to the origin of insects by resolving crustacean-hexapod relationships, *Current Biology*, 27(12), 1818-1824. <https://doi.org/10.1016/j.cub.2017.05.040>
- Sharan A., Datta B., & Lal A. (2023), Integrating numerical modelling and scenario-based sensitivity analysis for saltwater intrusion management: case study of a complex heterogeneous island aquifer system, *Environ Monit Assess*, 195:553. <https://doi.org/10.1007/s10661-023-11159-z>

- Sługocki Ł., & Czerniawski R. (2023), Water quality of the Odra (Oder) River before and during the ecological disaster in 2022: A warning to water management, *Sustainability*, 15(11), 8594. <https://doi.org/10.3390/su15118594>
- Smeti E., von Schiller D., Karaouzas I., Laschou S., Vardakas L., Sabater S., Tornés E., Monllor-Alcaraz L.S., Guillem-Argiles N., Martinez E., Barceló D., López de Alda M., Kalogianni E., Elozegi A., & Skoulikidis N. (2019), Multiple stressor effects on biodiversity and ecosystem functioning in a Mediterranean temporary river, *Science of The Total Environment*, Volume 647, Pages 1179-1187. <https://doi.org/10.1016/j.scitotenv.2018.08.105>
- Strayer D. L., & Dudgeon D. (2010), Freshwater biodiversity conservation: recent progress and future challenges, *J. North Am. Benthol. Soc.* 29, 344–358. <https://doi.org/10.1899/08-171.1>
- Suárez-Morales E. (2015), Class maxillopoda, In *Thorp and Covich's Freshwater Invertebrates* (pp. 709-755). Academic Press. <https://doi.org/10.1016/B978-0-12-385026-3.00029-2>
- Tickner D., Opperman J. J., Abell R., Acreman M., Arthington A. H., Bunn S. E., ... & Young L. (2020), Bending the curve of global freshwater biodiversity loss: an emergency recovery plan, *BioScience*, 70(4), 330-342. <https://doi.org/10.1093/biosci/biaa002>
- Tierno de Figueroa J.M., López-Rodríguez M.J., Fenoglio S. *et al.* (2013), Freshwater biodiversity in the rivers of the Mediterranean Basin, *Hydrobiologia*, 719, 137–186. <https://doi.org/10.1007/s10750-012-1281-z>
- Touati L. (2008), Distribution spatio-temporelle des Genres Daphnia et Simocephalus dans les mares temporaires de la Numidie, Doctoral dissertation. 8 May 1945 University of Guelma, Algeria.
- Tripathi S., Purchase D., Chandra R., Nadda A. K., & Chaturvedi P. (2023), Emerging pollutants characterization, mitigation and toxicity assessment of sewage wastewater treatment plant-India: a case study, *Journal of Contaminant Hydrology*, 254, 104139. <https://doi.org/10.1016/j.jconhyd.2023.104139>
- Umi W. A. D., Yusoff F. M., Balia Yusof Z. N., Ramli N. M., Sinev A. Y., & Toda T. (2024), Composition, Distribution, and Biodiversity of Zooplanktons in Tropical Lentic Ecosystems with Different Environmental Conditions, *Arthropoda*, 2(1), 33-54. <https://doi.org/10.3390/arthropoda2010003>
- Urdiales Flores D., *et al.* (2023), Drivers of accelerated warming in Mediterranean climate-type regions. *npj Clim. Atmos. Sci.*, 6, 97.
- Wacker A., & Harzsch S. (2021), Crustaceans in a changing world, *Zoology (Jena, Germany)*, 146, 125921. <https://doi.org/10.1016/j.zool.2021.125921>
- Valade P. (2003), Premier inventaire des poissons et des macrocrustacés des principales rivières pérennes de l'île Maurice, ARDA / GIP ECOFOR. <http://dx.doi.org/10.13140/RG.2.2.29030.19525>
- Walossek D. (1993), The upper Cambrian Rehbachiella and the phylogeny of Branchiopoda and Crustacea, In *The Upper Cambrian Rehbachiella and the phylogeny of Branchiopoda and Crustacea* (pp. 1-202). <https://doi.org/10.18261/8200374874-1993-01>
- Wickham H. (2016), ggplot2: Elegant graphics for data analysis. Springer-Verlag New York.
- Xiong X., Zhang K., Chen Y., Qu C., & Wu C. (2020), Arsenic in water, sediment, and fish of lakes from the Central Tibetan Plateau, *Journal of Geochemical Exploration*, 210, 106454. <https://doi.org/10.1016/j.gexplo.2019.106454>
- Yavuzatmaca M. (2019), Comparative analyses of non-marine Ostracods (Crustacea) among water basins in Turkey, *Acta Zoologica Academiae Scientiarum Hungaricae: An international Journal of Animal Taxonomy and Ecology*, 65(3), 269-297. <https://doi.org/10.17109/AZH.65.3.269.2019>
- Yavuzatmaca M. (2020), Species assemblages of Ostracoda (Crustacea) from west-site of Turkey: their indicator potential for lotic and lentic habitats, *Biologia*, 75(12), 2301-2314. <https://doi.org/10.2478/s11756-020-00494-y>
- Zhang X. L., Shu D. G., & Erwin D. H. (2007), Cambrian naraoiids (Arthropoda): morphology, ontogeny, systematics, and evolutionary relationships, *Journal of Paleontology*, 81(sp68), 1-52. <https://doi.org/10.1666/06-082.1>
- Zouggar A. M., Mirleau P., Chenuil A., Denis F., Caruso A., Ayad S. M. E. A. A., & Lamara S. A. C. (2023), Extreme and Heterogeneous Conditions of the Desert Wetland Chott Ech Chergui (Algeria) Allow Isolating Halophilic, Alkalophilic and Thermophilic Bacteria, *Wetlands*, 43(8), 108. <https://doi.org/10.1007/s13157-023-01757-5>