

1 **A versatile *Bacillus* spp. from garlic plant as a promising eco-friendly agent: Multiple abiotic**
2 **stress tolerance, exopolysaccharides production, antioxidant and antifungal activities**

3 Lamia Benhalima^{1,2*}, Sandra Amri^{1,2}, Saber Belhaoues³, Lina Haddam ¹, Rym Maghlout ¹, and
4 Mourad Bensouilah²


5 ¹University 8 Mai 1945, Faculty of the Nature and the Life Sciences and the Earth and the Universe
6 Sciences, Department of Biology, P. B.401, Guelma 24000, Algeria.

7 ²Laboratory of Ecobiology of Marine Environment and Coastlines, University Badji Mokhtar
8 Annaba, Faculty of Sciences, P.B. 12, Annaba 23000, Algeria.

9 ³National Research Centre for Environment (CRE), Annaba 23000, Algeria.

10 *to whom all correspondence should be addressed: e-mail: benhalima.lamia@univ-guelma.dz

11 Graphical abstract



12 Abstract

13 The selection of plant-associated bacteria tolerant to various stressors offers an eco-efficient strategy
 14 for sustainable agriculture. This study evaluates the resilience of four *Bacillus* spp. endophytes
 15 isolated from garlic (*Allium sativum*) roots against key abiotic and biotic challenges, including
 16 drought, salinity, herbicide (florasulam/2,4-D ester), heavy metals (Pb and Cd), and the
 17 phytopathogen *Botrytis cinerea*. Among the isolates, AseB4 exhibited the highest stress tolerance,
 18 achieving maximal growth (0.985 ± 0.27) under drought conditions (-1.0 MPa) and maintaining a
 19 survival rate of 92.62 ± 1.32 under high salinity (400 mM NaCl). All strains tolerated herbicide
 20 concentrations up to $1200 \mu\text{L mL}^{-1}$, and maintained robust growth under heavy metal stress, with a
 21 maximum tolerated concentration of 5 mM for Pb and Cd. EPS production and antioxidant activity
 22 increased dose-dependently with metal concentration, tending to a saturation point at higher levels
 23 (Tukey's test; $p < 0.05$). Additionally, the isolates displayed strong antagonistic activity against
 24 *Botrytis cinerea* ($>96\%$). These findings highlight the potential of garlic-associated *Bacillus*
 25 endophytes as sustainable microbial resources for stress mitigation, supporting the development of
 26 bio-based agents to reduce chemical inputs and enhance environmental resilience.

27 **Keywords:** Endophytic *Bacillus*, garlic roots, drought, salinity, herbicide, heavy metals, biological
 28 activity.

29 1. Introduction

30 In light of climate change and population pressure, numerous abiotic and biotic stresses have toxic
31 effects on ecosystems and can cause serious human, plant, and animal health risks. Currently, the
32 most significant factors that affect agricultural productivity are salinity, drought and toxic chemicals
33 such as pesticides and heavy metals, which can disturb nutritional balance, photosynthesis,
34 physiological functions, and development of plants (Kamran *et al.*, 2020; Sharath *et al.*, 2021; Anand
35 *et al.*, 2023). Exposure time, concentrations, and formulations of herbicides can seriously alter the
36 metabolism and the growth of the soil microbial biodiversity, leading to negative consequences such
37 as the unavailability of nutrients, low soil fertility, a slowdown in plant development, and increased
38 vulnerability to pathogens (Vera *et al.*, 2025). Furthermore, toxic metals like cadmium and lead,
39 which are released in ecosystems via anthropogenic activities (industrial waste, electronics, mining,
40 and agriculture), can lead to several damages including water and soil pollution, cell deformation in
41 plants and animals, and contamination of the human food chain (Kumar *et al.*, 2025; Zhang *et al.*,
42 2024). Many plants are impacted by these hazardous metals, which cause alterations in nutrient
43 uptake and water absorption, leading to reduced plant development and production, and even death
44 (Shahid *et al.*, 2024). In addition to abiotic stresses, biotic stressors, including pathogens, can cause
45 significant economic losses. *Botrytis cinerea* is one of phytopathogens that can infect more than 200
46 plant species (*e.g.*, tomato) and cause severe loss in crop development (Toral *et al.*, 2020).

47 Increased mass concerns about environmental stresses, food safety, chemicals in food, crop losses,
48 and pollution of ecosystems has led to encourage the investigation of biological approaches.
49 Currently, plant growth-promoting bacteria (PGPB), known for their beneficial abilities, represent a
50 crucial solution for minimizing serious agricultural impacts while promoting yields and food security.
51 Among these bacteria, endophytic *Bacillus* species play a key role in the protection of plants against
52 harmful stresses (Rafanomezantsoa *et al.*, 2025). The endospore forming ability allows the members
53 of *Bacillus* genus to cope with harsh environmental conditions and climate-induced vulnerabilities.
54 Furthermore, their capability to solubilize nutrients and produce phytohormones, siderophores,

55 organic acids, hydrolytic enzymes, and several antimicrobial substances makes them valuable
56 candidates for mitigating both abiotic and biotic stresses in diverse cultivated plants and enhancing
57 productivity (Nandana and Anith, 2025). Another noteworthy characteristic of PGPB *Bacillus* is the
58 production of EPS, which are macromolecular constituents, synthesized by planktonic cells within
59 metabolism and growth. EPS supply energy for cells, retain water, generate biofilm, and adsorb ions
60 like toxic heavy metals (Reddy *et al.*, 2024). Recently, numerous researches highlighted the
61 importance of EPS-producing bacteria (more than 70 species have been characterized) in heavy metal
62 bioremediation which contribute to environmental detoxification and plant protection (Zhang *et al.*,
63 2024; Bhardwaj, 2025). Microbial EPS are currently used in several sectors due to their unique
64 qualities and the simplicity of their extraction. Therefore, the selection of potential EPS-producers
65 remains one of the most crucial approaches. Under various stresses, including toxic metals, reactive
66 oxygen species, highly reactive molecules, can be generated and damage cells (Abou-Aly *et al.*,
67 2019). To counteract this oxidative damage, endophytes can produce potent antioxidant compounds
68 that inhibit the activation or multiplication of oxidative chain reactions (Muhtari *et al.*, 2024).
69 Accordingly, endophytes can also be a key source of novel pharmacological products.

70 In recent years, the attention of researchers has been attracted by endophytes from medicinal and
71 aromatic plants, which can have similar abilities to their hosts in producing bioactive compounds
72 such as antioxidants and antimicrobials, playing a role in the protection and the development of plants
73 (Sarjono *et al.*, 2019; Muhtari *et al.*, 2024). To date, Garlic (*Allium sativum* L.) is known for its
74 culinary and medicinal characteristics (*e.g.*, antimicrobial, antioxidant, anti-inflammatory, and anti-
75 cancer effects) thanks to its bioactive molecules, among which saponins, polysaccharides, and
76 organosulfur compounds (Poplawska *et al.*, 2024). So, garlic plant can also constitute a rich source
77 of endophytic bacteria with biotechnological and medical properties. Our previous study
78 (unpublished data) demonstrated that garlic roots of Guelma district (Northeastern Algeria) represents
79 an exceptional source of endophytes that exhibited PGP features, which supported the findings of
80 Srivastava *et al.* (2024), who recovered several endophytes from garlic plant with antibacterial

81 activity. However, few studies have investigated the tolerance of garlic-associated bacteria to abiotic
82 stresses, although *Allium sativum*, rich in reactive organosulfur compounds (e.g., allicin) (Poplawska
83 *et al.*, 2024), represents a chemically selective niche that may select for adapted and functionally
84 valuable endophytes with potential as a multifaceted microbial toolbox for sustainable agriculture
85 and environmental resilience. In the agricultural areas of Guelma, there is no information on the
86 tolerance and the compatibility of endophytes with the common pollutants. In order to discover
87 interesting functional properties of garlic-root endophytic bacteria in this zone, with a view to future
88 environmental applications, the current study aimed to assess the ability of endophytic *Bacillus* spp.
89 from garlic-roots in Guelma district to tolerate various abiotic stresses (drought, salinity, herbicide,
90 and heavy metals), determine their antioxidant activity and EPS production under lead and cadmium
91 stressors, and evaluate their ability to control the phytopathogen *Botrytis cinerea* in vitro.

92 **2. Materials and Methods**

93 *2.1. Selection of Bacterial strains from garlic roots*

94 Four *Bacillus* spp. were selected among 13 endophytic bacteria isolated from the garlic plant roots
95 *Allium sativum* L. The plant material (n = 5) was sampled during the spring of 2024, from Ben Djerrah
96 region (36° 25' 56" N, 7° 22' 7" E), situated in Guelma district, northeastern Algeria. The samples
97 were transported in aseptic conditions to the laboratory within 1 hour of harvest. Healthy roots were
98 initially washed in tap water to remove soil attached, and then surfaces were sterilized twice with
99 70% ethanol for 30 seconds, separated by a wash with 5% sodium hypochlorite for 3 min. In order to
100 select only the endophytic bacteria, the roots were also soaked in a 10% sodium bicarbonate solution
101 for 15 min and then washed at least five times with distilled water. One hundred ml of the final wash
102 were streaked into nutrient agar (NA) (Merck, Germany) and incubated for a day at 28 °C to confirm
103 the sterilization procedure. The sterilized roots were macerated in 90 ml of sterile physiological saline
104 at 180 rpm for one hour at ambient temperature. Aliquots of 50 µl were streaked on the NA and the
105 Luria Bertani Agar (LBA) (Merck, Germany), and incubated at 30 °C for 24 hours (Nagah *et al.*,
106 2024). After phenotypic identification according to microbiological tests described in Bergey's

107 Manual of determinative Bacteriology (Holt, 1994), and the determination of PGP characteristics, the
108 four best promising *Bacillus* spp. strains were selected for this study to assess their tolerance to the
109 most significant abiotic and biotic stresses. Isolates were denominated AsEBx (AsEB1, AsEB2,
110 AsEB3, and AsEB4). To obtain a standard inoculum, each strain was adjusted to 0.5 McFarland scale
111 after incubation in the Luria-Bertani broth (LBB) (Merck, Germany) at 150 rpm and 30 °C for 18
112 hours.

113 2.2. *Tolerance to drought stress*

114 In order to estimate the growth of endophytic bacteria under osmotic potentials of 0 MPa, -0.25 MPa,
115 -0.5 MPa, -0.75 MPa, and -1.0 MPa, screening of drought tolerant AsEBx strains was performed
116 on LBB medium enriched with 0, 2.4, 4.7, 6.5 and 7.95 g/100 ml polyethylene glycol 6000 (PEG),
117 respectively. As PEG can alter the physicochemical properties of the medium, the pH was carefully
118 adjusted to 7.0±0.2 following its addition in all treatments. After 24 h of incubation at 37 °C with
119 shaking at 180 rpm, the optical density (OD) of 10 ml of each isolate was estimated spectroscopically
120 (600nm). The stains were classified into: Highly sensitive (OD<0.3), sensitive-to-tolerant
121 (0.3<OD<0.39), tolerant (0.4<OD<0.5), and highly tolerant (OD>0.5) (Nader *et al.*, 2024).

122 2.3. *Tolerance to salt stress*

123 The standard bacterial inoculums were inoculated into the Tryptone Soy Broth (TSB) contained
124 escalating concentrations of NaCl (0, 50, 100, 200, and 400 mM) (Merck, Germany), and incubated
125 in a shaker at 140 rpm and 30 °C for 5 days. The growth rate of tolerant-bacteria was calculated every
126 day after measurement of the OD at 600 nm and comparison with the control (Soto-Varela *et al.*,
127 2024). The bacterial strain exhibiting the highest tolerance was selected for subsequent germination
128 assays on tomato seeds under salt stress as described by Akbaba and Özden (2023). Briefly, surface-
129 sterilized seeds (1% NaClO, 1 min) were soaked for 30 min in bacterial suspensions (OD₆₀₀ = 0.1).
130 After drying under sterile conditions, the seeds were incubated on moist filter paper in Petri dishes
131 containing either distilled water or 400 mM NaCl. Germination percentage and seedling growth were
132 assessed after 5 days at 25 °C in the dark.

133 2.4. *Bacterial growth in response to florasulam/2,4-D stress*

134 The exploration of the bacterial growth under herbicide stress was assessed in LBB supplemented
135 with different concentrations of florasulam/2,4-D. This herbicide is widely used in Algeria especially
136 in high production cereal fields. It contains two active compounds, 17.4 mM of florasulam and 900
137 mM of 2,4-D ester. According to the doses recommended by the producer to use in cereal areas, five
138 concentrations (60, 150, 300, 600, and 1200 $\mu\text{l/ml}$) were tested. The quantitative tolerance of the four
139 strains was determined using a spectrophotometer (OD_{600 nm}) after 1, 2, 3, 4, 7, 8, 9 and 10 days of
140 incubation at 180 rpm and 30 °C. The dry biomass weight (g L^{-1}) was estimated after cell bacteria
141 recuperation, washing with PBS (Phosphate-Buffered Saline) (Merck, Germany), and drying at 60
142 °C. A calibration curve DO₆₀₀ vs dry biomass weight was used (Briceño *et al.*, 2020).

143 2.5. *Bacterial growth in response to lead and cadmium stress*

144 The tolerance screening of the endophyte isolates was firstly confirmed qualitatively by inoculation
145 of the four AsEBx strains on LBA supplemented with different concentrations (ranging from 0.1 to
146 0.9 mM) of lead ($\text{Pb}(\text{NO}_3)_2$) and cadmium (CdSO_4) (Sigma-Aldrich, Germany). After incubation at
147 30 °C for 48 hours, the tolerant heavy metal strains were selected based on their ability to grow rapidly
148 even at the highest concentration, and the creamy aspect of their colonies which facilitate their
149 handling (Alvarado-Campo *et al.*, 2023). Quantitative heavy metal tolerance was performed using
150 broth diffusion method. Briefly, concentrations of 1, 2, 3, 4 and 5 mM of heavy metals salts were
151 prepared in LB. A volume of 1 ml of each standard inoculum was added to 1 ml of each concentration
152 and incubated at 30 °C and 140 rpm for 10 days. Bacterial growth was measured spectroscopically
153 (OD_{600nm}) at 1, 2, 3, 4, 7, 8, 9 and 10 days. Negative and positive controls (non-inoculated LB and
154 LB without heavy metal salts, respectively) were prepared in the same condition. The experiment was
155 conducted in triplicate. Maximum tolerance concentration (MTC) was determined by spotting of 10
156 μl of each inoculum on LBA. MTC corresponds to the highest concentration that shows visible
157 bacterial colonies (Shylla *et al.*, 2021).

158 2.6. *Estimation of exopolysaccharide production for tolerating heavy metal stress*

159 The concentrations of EPS produced by heavy metal-tolerant strains were deduced using sulfuric acid
160 anthrone colorimetry method (Zhang *et al.*, 2024). For this, broth cultures were centrifuged at 7 000
161 rpm for 10 min at 4°C. The cell pellet was homogenized with three volumes of absolute ethanol and
162 kept overnight at 4 °C to precipitate EPS and remove residual medium sugars. After drying at 37 °C,
163 the pellet was resuspended in 1 mL distilled water and treated with 0.5 mL of Sevage reagent for
164 protein removal, followed by centrifugation after an hour at 37 °C. The purified EPS fraction was
165 then mixed with 5 mL sulfuric acid anthrone reagent, heated at 100 °C for 10 min and OD was
166 determined spectrophotometrically (620 nm). EPS concentrations were expressed as $\mu\text{g mL}^{-1}$ using
167 glucose standard curve.

168 2.7. Antioxidant activities of metal-tolerant strains

169 The non-enzymatic antioxidant activities of the four metal-tolerant AsEBx strains were carried out
170 using the experimental protocol described by Abou-Aly *et al.* (2019). One percent of each bacterial
171 strain was inoculated in Mueller-Hinton broth (Merck, Germany) supplemented with the different
172 concentrations of lead and cadmium previously cited, and incubated for a day (150 rpm/30 °C). The
173 obtained cultures were centrifuged at 10,000×g for 5 min at 4 °C in order to obtain Cell Free Extracts
174 (CFE). A mixture of 500 μL of CFE and 3000 μL of 2-DiPhenyl-2-Picryl hydrazyl hydrate ([DPPH]
175 = 5 mg 100 mL^{-1} ethanol) (Sigma-Aldrich, Germany) was incubated in the dark for 30 min. The
176 ethanol is used to prepare the negative control. The ODs were determined at 517 nm using acid
177 ascorbic as a standard. The rate of antioxidant activities was calculated as:

$$178 \text{ DPPH residue (\%)} = \frac{\text{OD}_{\{517 \text{ control}\}} - \text{OD}_{\{517 \text{ sample}\}}}{\text{OD}_{\{517 \text{ control}\}}} \times 100 \quad (1)$$

$$179 \text{ Inhibition of DPPH (\%)} = 100 - \text{DPPH residue (\%)} \quad (2)$$

180 2.8. Assessment of antifungal ability of AsEBx strains

181 The antifungal activity of AsEBx strains against the phytopathogen *Botrytis cinerea* (isolated from
182 unhealthy tomato and supplied by the Annaba University, Algeria) was carried out on potato dextrose
183 agar (PDA) (Merck, Germany). In the same plate, a 5 mm fragment of a fresh *B. cinerea* and 5 μL of
184 a cultured *Bacillus* sp. strain in TSB were spotted facing each other. After 7 days of incubation at

185 28°C, the growth inhibition rate of the fungal pathogen was calculated in comparison with the control,
186 which was PDA inoculated with *B. cinerea* and TSB without bacteria (Weinand *et al.*, 2023):

187 Inhibition rate (%)

188
$$= \frac{\text{Diameter control fungal colony} - \text{Diameter fungal colony in presence of bacteria}}{\text{Diameter control fungal colony}} \times 100 \text{ (3)}$$

189 2.9. Statistical analysis

190 The results were reported as mean values with standard deviation represented by bars. The analysis
191 of variance (ANOVA) and Tukey's test were performed to estimate statistically significant
192 differences ($p < 0.05$) using SPSS software (IBM, USA).

193 3. Results

194 3.1. Attributes of selected Bacterial strains from garlic roots

195 From 13 endophyte bacteria isolated from garlic roots *Allium sativum* L., 4 *Bacillus* spp. were selected
196 for the current study. In our previous research, it has been showed that these endophytes exhibited
197 several functional traits such as high enzymatic activity, photohormone production, phosphate
198 solubilization, and promoting development of wheat and tomato seedlings (see Supplementary Table
199 S1). Phenotypic profiles demonstrated that our strains belong to the *Bacillus* spp. species. All the
200 interesting characteristics of the AsEBx strains were mentioned in Table 1.

201 3.2. Tolerance to drought and salinity stresses

202 The results in Table 2 demonstrate that all strains are highly tolerant to drought stress. Regarding the
203 OD values under unstressed conditions, there was a decrease in strains' growth under different
204 drought intensities ($p < 0.05$). Nevertheless, the isolates grew well and their OD values remained above
205 0.5. At high osmotic pressure, the strains AsEB1 and AsEB3 showed the lowest OD values. However,
206 the highest growth level was recorded for AsEB4 followed by AsEB2, indicating the best drought
207 tolerance. With the increase in NaCl concentration, a significant progressive reduction in the growth
208 rate of AsEB1, AsEB2, and AsEB3 strains was observed ($p < 0.05$). The AsEB4 isolate revealed the
209 highest tolerance to salinity stress (Table 2). Compared to the control, the reduction in AsEB4 growth

210 was very slight at 200 mM (1.45%) and remained low even at 400 mM (7.38%). In contrast, the
 211 AsEB2 strain was the most affected by NaCl, with growth reductions of 22.17% and 22.68% at 200
 212 mM and 400 mM NaCl, respectively. Nonetheless, it is important to emphasize that the growth rate
 213 of all tested strains remained above 70%.

214 The AsEB4 strain positively influenced the germination of tomato seeds under salinity condition (400
 215 mM NaCl) (Table. 3) (Tukey's test; $p < 0.01$). Although salinity reduced seedling growth compared to
 216 the standard condition (seeds inoculated with AsEB4), the recovered values remained substantial,
 217 suggesting that AsEB4 can play a role in mitigating salt stress during germination (see Supplementary
 218 Figure S1).

219 **Table 1.** Phenotypic profiles of isolated *Bacillus* spp. from garlic roots.

Isolate	AsEB1	AsEB2	AsEB3	AsEB4
<i>Morphological characterization</i>	Gram (+); Endospore forming rods; Mobile; Aerobic.	Gram (+); Endospore forming rods Mobile; Aerobic.	Gram (+); Endospore forming rods Mobile; Aerobic.	Gram (+); Endospore forming rods Mobile; Aerobic.
<i>Biochemistry characterization</i>				
Catalase	+	+	+	+
Oxidase	-	-	-	-
Citrate	+	+	+	+
Amylase	+	+	+	+
Esterase	+	+	+	+
Gelatinase	+	+	+	+
Lipase	+	+	+	+
Pectinase	+	+	+	+
Protease	+	+	+	+
Urease	+	+	+	+
Lactose	+	+	+	+
Glucose	+	+	+	+
Mannitol	-	+	+	+
Rhamnose	-	+	-	-
Saccharose	+	+	+	-
Methyl red test	+	-	-	-
VP test	+	-	-	+
Indole	-	-	-	-
<i>PGP traits</i>				
Phosphate solubilization	+	+	+	+
Hydrogen cyanide production	+	-	-	+
Indole-3-acetic acid production	> 114 µg/ml	> 97 µg/ml	> 96 µg/ml	> 108 µg/ml
Germination rate of seeds	> 94 %	> 82 %	> 80 %	> 93 %

220 *PGP*: Plant Growth Promoting, (+): positive reaction, (-): negative reaction

221 **Table 2.** Drought and salinity tolerance of endophytic *Bacillus* spp. isolated from garlic roots.

Isolate	AsEB1	AsEB2	AsEB3	AsEB4
<i>Drought stress*</i>				
0 MPa	1.887±0.01 ^a	1.721±0.3 ^{ab}	1.717±0.24 ^{ab}	1.907±0.03 ^a
-0.25 MPa	1.652±0.21 ^{ab}	1.551±0.02 ^{abc}	1.641±0.05 ^{ab}	1.703±0.1 ^{ab}
-0.5 MPa	0.972±0.03 ^{cd}	0.911±0.01 ^{cd}	0.981±0.03 ^{cd}	1.314±0.2 ^{bc}
-0.75 MPa	0.725±0.32 ^{de}	0.845±0.23 ^{cd}	0.872±0.1 ^{cd}	1.016±0.1 ^{cd}
-1.0 MPa	0.693±0.18 ^e	0.832±0.27 ^{cd}	0.632±0.13 ^e	0.985±0.27 ^{cd}
<i>Salinity stress**</i>				
0 mM	100±0.00 ^a	100±0.00 ^a	100±0.00 ^a	100±0.00 ^a
50 mM	99.82±15.29 ^a	97.16±5.5 ^{ab}	99.44±6.35 ^a	100±0.00 ^a
100 mM	96.56±7.96 ^{ab}	88.70±12.11 ^{bc}	92.13±7.36 ^{ab}	100±0.00 ^a
200 mM	96.43±7.99 ^{ab}	77.83±15.04 ^c	87.37±6.52 ^{bc}	98.55±8.45 ^a
400 mM	89.75±11.84 ^b	77.32±16.20 ^c	81.82±15.29 ^c	92.62±1.32 ^a

222 *: Results are expressed in OD₆₀₀ /24 h, **: Results are expressed in survival rate (%) /5 days, results are mean±SD, ^{a-e}:

223 indicate significant differences (Tukey's test; *p*<0.05).

224 **Table 3.** Effect of AsEB4 strain on germination and seedling growth of tomato under saline condition
225 after 5 days.

Parameters	Tomato seeds		
	Control*	Standard condition**	Saline condition***
Germination (%)	50.7±2.1 ^a	83.4±1.0 ^c	71.7±1.7 ^b
Shoot length (cm)	1.4±0.3 ^a	7.6±0.3 ^c	6.1±0.6 ^b
Root length (cm)	2.03±0.5 ^a	9.8±0.2 ^c	7.9±0.4 ^b
Shoot fresh weight (mg)	3.3±0.1 ^a	19±0.1 ^c	14.7±0.2 ^b
Root fresh weight (mg)	5.1±0.2 ^a	47.3±0.2 ^c	39±0.2 ^b
Shoot dry weight (mg)	0.5±0.2 ^a	2.7±0.3 ^c	1.9±0.1 ^b
Root dry weight (mg)	2.7±0.3 ^a	5.9±0.2 ^c	3.3±0.2 ^b


226 *: seeds without bacterial strain, **: seeds inoculated with AsEB4, ***: seeds inoculated with AsEB4 in 400 mM NaCl,

227 results are mean±SD, ^{a-c}: indicate significant differences between values within the same line (Tukey's test; *p*<0.05).

228 3.3. Bacterial growth in response to florasulam/2,4-D stress

229 As illustrate in figure 1, all isolates showed high tolerance to florasulam/2,4-D concentrations up to
230 1200 µL mL⁻¹. Compared to the control, strain AsEB2 exhibited non-significant growth reductions
231 (<10%) after 10 days (*p*=0.72). Furthermore, relative increases in growth of AsEB1, AsEB3, and
232 AsEB4 were noted at high herbicide concentrations. For the isolate AsEB1, the most of OD values at
233 1200 µL mL⁻¹ were higher than those measured in the control, indicating a better adaptive ability and
234 a prolonged tolerance to the herbicide stress. The dry biomass determination revealed that the strain

235 AsEB1 showed the highest biomass weight ($1.87 - 2.41 \text{ g L}^{-1}$), followed by AsEB4 ($1.81 - 2.23 \text{ g L}^{-1}$),
 236 1), AsEB3 ($1.8 - 2.22 \text{ g L}^{-1}$), and AsEB2 ($1.73 - 2.19 \text{ g L}^{-1}$). Detailed significance groupings at each
 237 time point are presented in Supplementary Table S2 (Tukey's test; $p < 0.05$).



238 **Figure 1.** Growth of endophytic *Bacillus* spp. isolated from garlic roots under florasulam/2,4-D
 239 herbicide stress.

240 3.4. Bacterial growth in response to lead and cadmium stress

241 The first screening of heavy metal-tolerant bacteria showed that all the AsEBx strains exhibited
 242 appropriate growth with creamy colonies at the maximum concentration 0.9 mM of Pb and Cd. The
 243 quantitative tolerance screening at increased toxic concentrations revealed that the four endophytic
 244 *Bacillus* spp. remained tolerant to tested heavy metals. Overall, all strains showed comparatively
 245 ($p > 0.05$) a decrease of their OD values, especially between the first and the third day. Starting from
 246 the fourth day, the strains demonstrated a slight increase in growth and maintained stability under a

247 potential metal stress until 10 days (Figure 2 and 3). In the case of lead, the isolate AsEB2 reached a
248 higher growth level (OD=0.572) at 5 mM after 8 days. On the other hand, the isolate AsEB3 exhibited
249 a robust growth at 5 mM of Pb, compared to lower concentrations, recording maximum OD values
250 over the entire period of incubation (Figure 2). In the presence of cadmium, the growth of AsEB1 in
251 toxic concentrations showed the highest OD mean values. In the other strains, OD₆₀₀ mean values
252 fluctuate between 0.3 and 0.5 were noted (Figure 3). Detailed significance groupings for bacterial
253 growth under Pb and Cd stresses at each time point, as determined by Tukey's test, are presented in
254 Supplementary Table S3 and S4. Based on the culture on LBA, results showed important colony
255 development at all tested concentrations with a MTC of 5 mM of each heavy metal (Figure 4),
256 confirming a greater tolerance of the AsEBx strains to the top level of Pb and Cd.


257 3.5. *Exopolysaccharide production for tolerating heavy metal stress*

258 All the strains exhibited the same trends of EPS production (Figure 5). The lowest EPS quantities
259 were noted in control (without any metal). However, EPS production was raised with the increasing
260 of heavy metals concentrations, with partial saturation at higher levels (Tukey's test; $p < 0.05$),
261 indicating probably an adaptive response in the AsEBx strains. Among the four heavy metal-tolerant
262 strains, AsEB1 isolate was the most important producer of EPS. The highest EPS amounts were noted
263 in presence of Cd at 4 and 5 mM (108.61 and 129.1 $\mu\text{g mL}^{-1}$, respectively). Under lead exposure, the
264 maximum ability for EPS secretion was recorded in AsEB3 isolate, which it reached 79.25 $\mu\text{g mL}^{-1}$
265 EPS at 5 mM of Pb. Furthermore, isolates AsEB2 and AsEB4 showed an enhanced production of
266 EPS up to $>65 \mu\text{g mL}^{-1}$ under each tested heavy metal stress.


267 3.6. *Antioxidant activities of metal-tolerant strains*

268 Results present in table 4 revealed that the four heavy metal-tolerant *Bacillus* spp. strains were able
269 to inhibit DPPH and exhibited effective antioxidant activities. Important rates ($> 90 \%$) of inhibited
270 DPPH was observed under the highest concentrations of the two heavy metals. Minimum rates of
271 residual DPPH under Pb and Cd were recorded by AsEB1 (9.1-6.8%, respectively), followed by
272 AsEB4 (9.3-7.9%, respectively), AsEB3 (9.5-8.1%, respectively), and AsEB2 (10.4-9.6%,


273 respectively). Whereas, the bottom level of antioxidant activities was occurred at the control (without
274 stress), demonstrating that DPPH inhibition effectiveness was gradually increased under stress
275 conditions.




276 **Figure 2.** Growth of endophytic *Bacillus* spp. isolated from garlic roots under lead stress.



277 **Figure 3.** Growth of endophytic *Bacillus* spp. isolated from garlic roots under cadmium stress.



278 **Figure 4.** Colony development of heavy metal-tolerant *Bacillus* spp. isolated from garlic roots at the
 279 maximum tolerance concentration of lead and cadmium.



280 **Figure 5.** Exopolysaccharides production by endophytic *Bacillus* spp. strains under lead and
 281 cadmium stresses. ^{a-d}: indicate significant differences (Tukey's test; p<0.05).

282 **Table 4.** Non-enzymatic antioxidant activities of endophytic *Bacillus* spp. isolated from garlic roots.


Heavy metal Concentration	AsEB1		AsEB2		AsEB3		AsEB4	
	DPPH residue (%)	Inhibited DPPH (%)	DPPH residue (%)	Inhibited DPPH (%)	DPPH residue (%)	Inhibited DPPH (%)	DPPH residue (%)	Inhibited DPPH (%)
0 mM (Control)	38.5 ^a	61.5	42.7 ^a	57.3	40.3 ^a	59.7	39.7 ^a	60.3
Pb								
1 mM	20.3 ^b	79.7	29.5 ^b	70.5	25.5 ^b	74.5	18.7 ^b	81.3
2 mM	18.7 ^c	81.3	21.8 ^c	78.2	20.2 ^c	79.8	15.6 ^c	84.4
3 mM	10.6 ^d	89.4	18.2 ^{cd}	81.2	19.7 ^{cd}	80.3	12.3 ^d	87.7
4 mM	9.6 ^d	90.4	12.8 ^d	87.2	13.4 ^d	86.6	10.7 ^d	89.3
5 mM	9.1 ^d	90.9	10.4 ^d	89.6	9.5 ^d	90.5	9.3 ^d	90.7
Cd								
1 mM	17.4 ^b	82.6	28.4 ^b	71.6	24.7 ^b	75.3	20.3 ^b	79.7
2 mM	15.3 ^c	84.7	25.7 ^c	74.3	19.8 ^c	80.2	17.8 ^c	82.2
3 mM	10.2 ^{cd}	89.8	20.3 ^{cd}	79.7	17.3 ^{cd}	82.7	15.4 ^{cd}	84.6
4 mM	8.4 ^d	91.6	13.6 ^d	86.4	9.7 ^d	90.3	11.6 ^d	88.4
5 mM	6.8 ^d	93.2	9.6 ^d	90.4	8.1 ^d	91.9	7.9 ^d	92.1

283 ^{a-d}: indicate significant differences between metal concentrations; Residual DPPH % were used for statistical tests

284 (Tukey's test; p<0.05).

285 3.7. Antifungal ability of AsEBx strains

286 The *in vitro* assays indicated that AsEBx strains exhibit antifungal activity against the gray mold
 287 *Botrytis cinerea*. The co-cultivation experiments revealed a strong inhibition of mycelial growth at
 288 an average rate of 97.78±0.21% by AsEB1, 96.97±0.1% by AsEB2, 96.11±0.24% by AsEB3, and
 289 97.22±0.11% by AsEB4. A representative image of the dual-culture assay is provided in Figure 6.



290
 291 **Figure 6.** Effect of endophytic *Bacillus* spp. strains on the mycelial growth of *Botrytis cinerea* after
 292 7 days at 28 ± 0.2 °C. (A) Colony growth of *B. cinerea* on the PDA medium as control; (B) Inhibition
 293 of *B. cinerea* growth by AsEB1; (C) Inhibition of *B. cinerea* growth by AsEB4.

294 4. Discussion

295 4.1. *Bacillus* strains from garlic roots

296 Endophytic *Bacillus* species are well documented for their plant growth-promoting functions and
297 stress mitigation capacity (Abuhena *et al.*, 2024). Our investigation indicates that garlic (*Allium*
298 *sativum* L.) roots constitute a selective ecological niche that favors metabolically versatile and stress-
299 adapted bacterial populations. While previous studies have mainly focused on classical PGP traits,
300 the present work advances current knowledge by evaluating isolates under both abiotic and biotic
301 pressures. To the best of our knowledge, such a diverse multifunctional profile has rarely been
302 reported for garlic root-associated *Bacillus* endophytes. The robust tolerance observed in the AsEBx
303 strains, combined with their EPS-production, antioxidant activity, and biocontrol capacity, reinforces
304 their relevance as biological inputs for resilient, low-input cropping systems (Nagah *et al.*, 2024).
305 Although the AsEBx were assigned to the genus *Bacillus* based on phenotypic and biochemical
306 characteristics, molecular identification using 16S rRNA gene sequencing would provide a more
307 precise taxonomic resolution and will be considered in future studies.

308 4.2. Tolerance to drought and salinity stresses

309 Water deficit and soil salinity are major constraints that reduce crop productivity and threaten global
310 food security. The exploration of endophytic bacteria capable of tolerating such stresses is critical for
311 developing strategies that enhance plant resilience under adverse environmental conditions. In the
312 present investigation, the ability of *Bacillus* spp. strains to maintain functional activity under severe
313 osmotic stress (-1.0 MPa), highlights its potential for sustaining crop germination and nutrient
314 availability under low water conditions. By facilitating nutrient acquisition when water is scarce,
315 these bacteria may act as biological buffers, mitigating the economic risks associated with drought-
316 induced yield losses (Sharath *et al.*, 2021). Regarding salinity, *Bacillus* spp. isolates displayed a
317 broad tolerance range (50 to 400 mM NaCl). These findings corroborate previous researches that
318 indicate the ability of *Bacillus* species to alleviate salinity stress (Kaur and Karnwal, 2023). The
319 inoculation with AsEB4 strain supported robust germination of tomato seeds under salinity (400 mM
320 NaCl). These data support the potential role of *Bacillus* spp. in mitigating salt stress, possibly through
321 mechanisms such as osmolyte synthesis and antioxidant activity. The observed stress resilience may

322 be attributed to enhanced production of exopolysaccharides and proline, which act as osmoprotectants
323 and free radical scavengers (Abuhena *et al.*, 2024; Nadar *et al.*, 2024).

324 4.3. Tolerance to florasulam/2,4-D stress

325 The screening of herbicide-tolerant bacteria represents a strategic approach to reduce the toxic effects
326 of herbicides and sustain crop productivity in intensive agricultural systems. The present study
327 advances current knowledge by documenting the adaptive capacity of garlic-associated *Bacillus* spp.
328 isolates to florasulam/2,4-D stress. These isolates maintained high cell densities, and biomass
329 accumulation progressed over time despite herbicide pressure. Similar to our data, previous studies
330 reported the tolerance of certain *Bacillus* strains exposed to different active molecules, such as
331 atrazine and paraquat, where growth remained unaffected or was only moderately influenced (Farias
332 *et al.*, 2021; Inthama *et al.*, 2021). While other studies, have reported total growth inhibition of all
333 *Bacillus* strains at 500 ppm of paraquat (Pradhan and Jena, 2023), our data suggest a higher level of
334 metabolic resilience. The pronounced tolerance observed in our garlic root-associated isolates may
335 be related to their endospore forming capability, which ensures persistence under chemical stress.
336 While the clarification of the culture medium after incubation could suggest a potential transformation
337 of the herbicide, this hypothesis requires further analytical confirmation. Notably, previous
338 investigations have shown that certain *Bacillus* species can degrade herbicides via specific metabolic
339 pathways, which may provide a plausible explanation for this observation (Ichor *et al.*, 2024; Zameer
340 *et al.*, 2023). The deployment of such herbicide-tolerant endophytes could help preserve beneficial
341 microbiota in treated soil, reduce the need for repeated chemical inputs, and potentially contribute to
342 the in-situ biodegradation of residual compounds.

343 4.4. Tolerance to heavy metals

344 Heavy metal pollution constitutes one of the foremost concerns for the environment and living
345 organisms' health. Currently, irrigation water and agricultural soils in Guelma district are under
346 increasing heavy metal pressure (Benhalima *et al.*, 2020; Sassane and Touati, 2024). Consequently,
347 resistance development in plant-associated bacteria to cope with heavy metal stressors is possible. In

348 this context, our isolates from Guelma agricultural region demonstrate remarkably high tolerance to
349 Pb and Cd. These levels align with the robust defense mechanisms identified in other specialized
350 endophytes (Biswas *et al.*, 2024; Kumar *et al.*, 2025). In addition to the detected PGP traits (direct
351 processes), our data suggest that AsEBx isolates can also improve plant growth by increasing plant
352 tolerance to high heavy metal levels (indirect process). The high metal tolerance observed in our
353 isolates likely reflects their prior exposure to metal-contaminated soils, as persistent heavy metals
354 significantly influence resistance development within endophyte communities (Liu J. *et al.*, 2024).
355 However, the initial decrease in growth observed during the first days, followed by a recovery phase,
356 may correspond to a metal-induced lag phase, indicating that these isolates undergo necessary
357 physiological adjustments to cope with metal toxicity. This behavior is commonly associated with
358 adaptive metabolic responses, including intracellular redox regulation, metal sequestration, and the
359 expression of chelating proteins, before resuming active proliferation (Liu L. *et al.*, 2024). Such
360 resilient microbial resources could support the development of cost-effective and environmentally
361 sustainable strategies for managing metal-affected soils (Liang *et al.*, 2025).

362 4.5. *Exopolysaccharide production for tolerating heavy metal stress*

363 The production of EPS by the studied *Bacillus* spp., which followed a dose-response relationship with
364 respect to the concentration of tested heavy metals, highlights its role as a key tolerance mechanism
365 against lead and cadmium stresses. EPS are known to act as extracellular biosorbents thanks to their
366 anionic functional groups (e.g., carboxyl and phosphate), which can bind cationic metals through
367 electrostatic interactions and complexation. This physiological plasticity allows bacterial strains to
368 construct a physical barrier, effectively immobilizing and sequestering toxic ions before they reach
369 the intracellular environment. The higher EPS production observed under Cd stress compared to Pb
370 may be reflect variations in metal behavior and interaction with EPS matrix. Cd²⁺ can exert strong
371 intracellular stress due to its mobility and bioavailability, inducing a more pronounced EPS synthesis.
372 In contrast, Pb²⁺ is more readily immobilized at the cell surface and forms stable precipitates, which
373 may reduce its intracellular impact and consequently the need for elevated EPS production (Zadeh *et*

374 *al.*, 2023; Zhang *et al.*, 2024). Such differences suggest that EPS may contribute to distinct
375 sequestration dynamics for Cd and Pb in our strains, although these interactions require further
376 investigation to be confirmed. These findings highlight these strains as efficient, eco-friendly agents
377 capable of transforming toxic stress into a manageable biological response. In this context, integrating
378 such metal-tolerant bacteria into contaminated soils could help reduce the bioavailability of Pb and
379 Cd through a self-sustaining biological process. As noted in similar observation, the sequestration of
380 metals by EPS not only ensures bacterial survival but also contributes to the establishment of a
381 protected rhizospheric niche (Bhardwaj, 2025).

382 4.6. *Antioxidant activities of metal-tolerant strains*

383 . The significant activation of antioxidant systems in AsEBx strains under Pb and Cd stress reflects a
384 dynamic metabolic adaptation to metal-induced oxidative stress. Our findings suggest that these
385 strains can reprogram environmental stress into a physiological trigger for the synthesis of high-value
386 secondary metabolites. Such metabolic plasticity opens avenues for the biotechnological production
387 of natural antioxidants and additives. The strong antioxidant potential of endophytic *Bacillus* is driven
388 by the secretion of specialized compounds, notably polyphenols and flavonoids, which effectively
389 neutralize free radicals (Tran *et al.*, 2025). Given the therapeutic richness of garlic, its associated
390 endophytes appear to recapitulate the host's chemical complexity to enhance plant defenses. This
391 symbiotic legacy positions our isolates not only as defensive partners for the host plant but also as
392 versatile biological agents capable of bridging the gap between microbial resilience and the
393 bioeconomy (Muhtari *et al.*, 2024).

394 4.7. *Antifungal ability of AsEBx strains*

395 The selection of beneficial bacterial strains as biocontrol agents represents a feasible and sustainable
396 approach controlling post-harvest deterioration of several crops and reducing the overuse of
397 chemicals (Herpich *et al.*, 2025). In this study, all tested *Bacillus* sp. strains exhibited remarkably
398 strong antifungal activity against *B. cinerea*, with inhibition rates exceeding 96%. This inhibition
399 level surpasses most recent reports and highlights the exceptional antagonistic capacity of garlic-

400 associated endophytes (Abuhena *et al.*, 2024). The strong inhibitory activity observed in AsEBx
401 strains may be associated with the production of hydrolytic enzymes, antimicrobial metabolites,
402 siderophore, as well as competition for nutrients, which contribute to pathogen suppression by
403 limiting fungal growth and interfering with cell integrity (Nagrle *et al.* 2023). The ability of our
404 isolates to almost completely inhibit a major post-harvest pathogen suggests their potential as high-
405 value bio-fungicides.

406 **5. Conclusion**

407 This study reveals the potential of garlic root-associated *Bacillus* endophytes as multifunctional
408 microbial resources capable of enhancing plant tolerance to multiple stresses, with promising
409 relevance for sustainable agricultural practices and bioeconomic strategies. The combination of plant
410 growth-promoting traits, tolerance to abiotic stressors, and strong antifungal activity underscores their
411 capacity to simultaneously mitigate biotic and abiotic constraints. In addition, the production of EPS
412 and antioxidant activity further strengthens their biotechnological relevance, as these traits may
413 contribute to soil stabilization, stress mitigation, and the development of more robust bioinoculant
414 formulations. Further research should focus on validating their performance under field conditions,
415 clarifying the molecular mechanisms underlying their stress-adaptive responses, and developing
416 scalable formulations to support environmentally responsible agricultural practices.

417 **Acknowledgments**

418 The authors are grateful to the Algerian Ministry of Higher Education and Scientific Research and
419 the Directorate General for Scientific Research and Technological Development (DGRSDT) for
420 supporting this work.

421 **References**

422 Abou-Aly H.E., Youssef A.M., El-Meihy R.M., Tawfik T.A. and El-Akshar E.A. (2019). Evaluation
423 of heavy metals tolerant bacterial strains as antioxidant agents and plant growth promoters.
424 *Biocatalysis and Agricultural Biotechnology*, **19**, 101110.

425 Abuhena M., Zhakypbek Y., Yerlan U., Aben A., Kamarkhan Z., Allakhverdiev S.I., Assemgul S.K.,
426 Al Rashid J., Karim M.D., Kuanysh T.T., Belkozhayev A.M. and Kossalbayev B.D. (2024). An
427 Overview of *Bacillus* species in agriculture for growth promotion, biocontrol and dry tolerance.
428 *ES Food Agroforestry*, **18**, 1321.

429 Akbaba M. and Özden E. (2023). Salt tolerance of endophytic root bacteria and their effects on seed
430 germination and viability on tomato plants. *Brazilian Journal of Microbiology*, **54**(4), 3147-
431 3162.

432 Alvarado-Campo K.L., Quintero M., Cuadrado-Cano B., Montoya-Giraldo M., Otero-Tejada E.L.,
433 Blandón L., Sánchez O., Zuleta-Correa A. and Gómez-León J. (2023). Heavy metal tolerance
434 of microorganisms isolated from coastal marine sediments and their lead removal potential.
435 *Microorganisms*, **11**, 2708.

436 Anand U., Pal T., Yadav N., Singh V.K., Tripathi V., Choudary K., Shukla A.K., Sunita K., Kumar
437 A, Bontempi E, Ma Y., Kolton M. and Singh A.K. (2023). Current scenario and future
438 prospects of endophytic microbes: Promising candidates for abiotic and biotic Stress
439 management for agricultural and environmental sustainability. *Microbial Ecology*, **86**, 1455–
440 1486.

441 Benhalima L., Amri S., Bensouilah M. and Ouzrout, R. (2020). Heavy metal resistance and
442 metallothionein induction in bacteria isolated from Seybouse River, Algeria. *Applied Ecology
443 and Environmental Research*, **18**(1), 1721-1737.

444 Bhardwaj A. (2025). Endophytes unleashed: natural allies for heavy metal bioremediation. *Discover
445 Plants*, **2**, 57.

446 Biswas S., Jayaram S., Philip I., Balasubramanian B., Pappuswamy M., Barceló D., Chelliapan S.,
447 Kamyab H., Sarojini S. and Vasseghian Y. (2024). Appraisal of the potential of endophytic
448 bacterium *Bacillus amyloliquefaciens* from *Alternanthera philoxeroides*: A triple approach to
449 heavy metal bioremediation, diesel biodegradation, and biosurfactant production. *Journal of
450 Environmental Chemical Engineering*, **12**(5), 113454.

451 Briceño G., Lamilla C., Leiva B., Levio M., Donoso-Piñol P., Schalchli H., Gallardo F. and Diez
452 M.C. (2020). Pesticide-tolerant bacteria isolated from a biopurification system to remove
453 commonly used pesticides to protect water resources. *PLoS ONE*, **15**(6), e0234865.

454 de Farias D.I.O.A, Leite R da C., Ribeiro E.A., Martins A.L.L. and Chagas Júnior A.F. (2021).
455 Glyphosate and atrazine inhibit growth of *Azospirillum brasilense*, *Bacillus subtilis*, *Bacillus*
456 *thuringiensis*, *Chromobacterium subtsugae* and *Saccharopolyspora spinosa*. *Agronomía*
457 *Colombiana*, **39** (1), 113-120.

458 Herpich C.H., Welter L.J., Botelho G.R., da Silva F.N., Casa R.T. and Bogo A. (2025). Epiphytic and
459 endophytic bacteria for the control of *Botrytis cinerea in vitro* and in grape berries of cv.
460 sauvignon blanc in southern Brazil. *Bioscience Journal*, **41**, e41005.

461 Holt J.G. (1994). Bergey's manual of determinative bacteriology. Ninteth Edition, Lippincott Williams
462 and Wilkins, Baltimore.

463 Ichor T., Yange A.I. and Ebah E.E. (2024). Production of metabolites from bacterial degradation of
464 some selected herbicides in pristine soil. *Journal of Microbes and Research*, **3**(2).

465 Inthama P., Pumas P., Pathom-Aree W., Pekkoh J. and Pumas C. (2021). Plant growth and drought
466 tolerance-promoting bacterium for bioremediation of paraquat pesticide residues in agriculture
467 soils. *Frontiers in Microbiology*, **12**, 446.

468 Kamran M., Parveen A., Ahmar S., Malik Z., Hussain S., Chattha M.S., Saleem M.H., Adil M.,
469 Heidari P. and Chen J.T. (2020). An overview of hazardous impacts of soil salinity in crops,
470 tolerance mechanisms, and amelioration through selenium supplementation. *International*
471 *Journal of Molecular Sciences*, **21**(1),148.

472 Kaur M. and Karnwal A. (2023). Screening of endophytic bacteria from stress-tolerating plants for
473 abiotic stress tolerance and plant growth-promoting properties: Identification of potential
474 strains for bioremediation and crop enhancement. *Journal of Agriculture and Food Research*,
475 **14**, 100723.

- 476 Kumar H., Khan U. and Gond S.K. (2025). Functional characterization and alleviation of cadmium
477 stress by endophytic bacteria isolated from *Cynodon dactylon* (L.) Pers. *Total Environment*
478 *Microbiology*, **1**(2), 100015.
- 479 Liang B., Feng Y., Ji X., Li C., Li Q., Zeng Z. and Wang Y. (2025). Isolation and characterization of
480 cadmium-resistant *Bacillus cereus* strains from Cd-contaminated mining areas for potential
481 bioremediation applications. *Frontiers in Microbiology*, **16**,1550830.
- 482 Liu J., Liu C., Zheng J., Zhang X., Zheng K. and Zhuang J. (2024). Response of plant endophyte
483 communities to heavy metal stress and plant growth promotion by the endophyte *Serratia*
484 *marcescens* (Strain JG1). *Plants*, **13**, 2755.
- 485 Liu L., Quan S., Li L., Lei G., Li S., Gong T., Zhang Z., Hu Y. and Yang,W. (2024). Endophytic
486 bacteria improve bio- and phytoremediation of heavy metals. *Microorganisms*, **12**, 2137.
- 487 Muhtari K., Sailaja I., Shehawat B.K. and Kaura S. (2024). Antimicrobial and antioxidant activities
488 of endophytic bacteria associated with medicinal plants. *Agricultural and Biological Research*,
489 **40**(4),1178-1184.
- 490 Nader A.A., Hauka F.I.A., Afify A.H. and El-Sawah A.M. (2024). Drought-tolerant bacteria and
491 *Arbuscular* mycorrhizal fungi mitigate the detrimental effects of drought stress induced by
492 withholding irrigation at critical growth stages of soybean (*Glycine max*, L.). *Microorganisms*,
493 **12**, 1123.
- 494 Nagah A., El-Sheekh M.M., Arief O.M., Alqahtani M.D., Alharbi B.M. and Dawwam G.E. (2024).
495 Endophytic *Bacillus vallismortis* and *Bacillus tequilensis* bacteria isolated from medicinal
496 plants enhance phosphorus acquisition and fortify *Brassica napus* L. vegetative growth and
497 metabolic content. *Frontiers in Plant Science*, **15**, 1324538.
- 498 Nagrale D.T., Chaurasia A., Kumar S., Gawande S.P., Hiremani N.S., Shankar R., Gokte-
499 Narkhedkar N., Renu Y. and Prasad G. (2023). PGPR: the treasure of multifarious beneficial
500 microorganisms for nutrient mobilization, pest biocontrol and plant growth promotion in field
501 crops. *World Journal of Microbiology and Biotechnology*, **39**, 100.

- 502 Nandana M.S. and Anith K.N. (2025). Endospore-producing endophytes: Structural adaptations,
503 functional roles, and their significance in plant growth and stress management. *International*
504 *Journal of Plant & Soil Science*, **37**(8), 298-312.
- 505 Popławska N.A., Śliz J., Skorupska M., Czeczotka M.J. and Woźniak K. (2024). Garlic (*Allium*
506 *sativum* L.): A review of varied health benefits. *Journal of Education, Health and Sport*, **72**,
507 51512.
- 508 Pradhan S. and Jena M. (2023). Evaluation of compatibility of native plant growth promoting
509 rhizobacterial isolates against chemical pesticides. *Ecology, Environment & Conservation*. **29**,
510 S206-S211.
- 511 Rafanomezantsoa P., El-Hasan A. and Voegelé, R.T. (2025). Potential of *Bacillus halotolerans* in
512 mitigating biotic and abiotic stresses: A comprehensive review. *Stresses*, **5**, 24.
- 513 Reddy K.V., Ranjit P., Priyanka E., Maddela N.R. and Prasad, R. (2024). Bioremediation of heavy
514 metals-contaminated sites by microbial extracellular polymeric substances-A critical view.
515 *Environmental Chemistry and Ecotoxicology*, **6**, 408–421.
- 516 Sarjono P.R., Putri L.D., Budiarti C.E., Mulyani N.S., Ngadiwiyana, Ismiyanto, Kusrini, D. and
517 Adiwibawa Prasetya N. B. (2019). Antioxidant and antibacterial activities of secondary
518 metabolite endophytic bacteria from papaya leaf (*Carica papaya* L.). *IOP Conference Series:*
519 *Materials Science and Engineering*, **509**(1), Article 012112.
- 520 Sassane A. and Touati M. (2024). Metallic groundwater contamination assessment using the heavy
521 metal pollution index: a case study of the Guelma plain alluvial groundwater, Northeast of
522 Algeria. *Sustainable Water Resources Management*, **10**(1), 30.
- 523 Shahid M., Altaf M., Ali S. and Tyagi A. (2024). Isolation and assessment of the beneficial effect of
524 exopolysaccharide-producing PGPR in *Triticum aestivum* (L.) plants grown under NaCl and
525 Cd -stressed conditions. *Plant Physiology and Biochemistry*, **215**, 108973.

526 Shahid M. and Khan M. S. (2021). Tolerance of pesticides and antibiotics among beneficial soil
527 microbes recovered from contaminated rhizosphere of edible crops. *Current Research in*
528 *Microbial Sciences*, **3**, 100091.

529 Sharath S., Triveni S., Nagaraju Y., Latha P.C. and Vidyasagar B. (2021). The Role of phyllosphere
530 bacteria in improving cotton growth and yield under drought conditions. *Frontiers in*
531 *Agronomy*, **3**, 680466.

532 Shylla L., Barik S.K., Joshi S.R. (2021). Characterization and bioremediation potential of native
533 heavy-metal tolerant bacteria isolated from rat-hole coal mine environment. *Archives of*
534 *Microbiology*, **203**, 2379–2392.

535 Soto-Varela Z.E., Orozco-Sánchez C.J., Bolívar-Anillo H.J., Martínez J.M., Rodríguez N.,
536 Consuegra-Padilla N., Robledo-Meza A. and Amils R. (2024). Halotolerant endophytic bacteria
537 *Priestia flexa* 7BS3110 with Hg²⁺ tolerance isolated from *Avicennia germinans* in a Caribbean
538 Mangrove from Colombia. *Microorganisms*, **12**, 1857.

539 Srivastava P., Tiwari S.P., Srivastava A.K. and Sharma R. (2024). Optimization of sterilization
540 parameters for isolation of endophytes from *Allium sativum* and exploring its antibacterial
541 activity. *Journal of Pure and Applied Microbiology*, Article 9099.

542 Toral L., Rodríguez M., Béjar V. and Sampedro I. (2020). Crop protection against *Botrytis cinerea* by
543 rhizosphere biological control agent *Bacillus velezensis* XT1. *Microorganisms*, **8**(7), 992.

544 Tran L.C., Duc C.K.T., Nguyen T.T., Pham D.T., Phan D.K., Dai T.T.X. and Nguyen T.T. (2025).
545 Evaluation of antioxidant and anti-inflammatory activities of *Bacillus* sp. CZ-Rh4, CZ-Rh7,
546 and CZ-L11 extracts. *Egyptian Journal of Botany*, **65**(3), 557-566.

547 Vera J., Sarango Y., Villamar M., Ortiz J., Sevilla-Carrasco J., Duarte J. and Lucas L. (2025). Effect
548 of herbicides on the growth of beneficial microorganisms in rhizospheric soil. *Revista De La*
549 *Facultad De Agronomía De La Universidad Del Zulia*, **42**(2), e254222.

- 550 Weinand T., El-Hasan A. and Asch F. (2023). Role of *Bacillus* spp. plant growth promoting properties
551 in mitigating biotic and abiotic stresses in lowland rice (*Oryza sativa* L.). *Microorganisms*, **11**,
552 2327.
- 553 Zadeh P.H., Feroso F.G., Collins G., Serrano A., Mills S. and Abram, F. (2023). Impacts of metal
554 stress on extracellular microbial products, and potential for selective metal recovery.
555 *Ecotoxicology and Environmental Safety*, **252**, 114604.
- 556 Zameer M., Tahir U., Khalid S., Zahra N., Sarwar A., Aziz T., Saidal A., Alhomrani M., Alamri A.,
557 Dabool A.S., Sameeh M.Y., Mohamed A.A. and Alharbi A. (2023). Isolation and
558 characterization of indigenous bacterial assemblage for biodegradation of persistent herbicides
559 in the soil. *Acta Biochimica Polonica*, **70**(2), 325-334.
- 560 Zhang H., Wang K., Liu X., Yao L., Chen Z. and Han H. (2024). Exopolysaccharide-producing
561 bacteria regulate soil aggregates and bacterial communities to inhibit the uptake of cadmium
562 and lead by Lettuce. *Microorganisms*, **12**, 2112.

563 **Supplementary Material**564 **Table S1.** Effect of endophytic *Bacillus* spp. isolated from garlic roots on germination and seedling growth of wheat and tomato seeds after 14 days.

Parameters	Wheat seeds					Tomato seeds				
	Control	Treatment				Control	Treatment			
		AsEB1	AsEB2	AsEB3	AsEB4		AsEB1	AsEB2	AsEB3	AsEB4
Germination (%)	49.6±4.1 ^a	90.4±2.6 ^b	85.2±1.2 ^b	88.3±4.6 ^b	95.3±1.2 ^c	51.3±2.1 ^a	94.6±1.8 ^c	82.5±1.4 ^b	81.7±1.2 ^b	93.4±2.0 ^c
Shoot length (cm)	9.1±1.5 ^a	19.4±1.1 ^c	19.8±0.7 ^c	20.2±1.2 ^c	21.3±0.5 ^d	9.8±0.6 ^a	16.2±0.3 ^c	17.3±0.2 ^c	20.4±0.6 ^d	20.8±0.9 ^d
Root length (cm)	13.15±0.5 ^a	21.2±1.5 ^d	22.±0.5 ^c	22.8±0.4 ^c	23.7±0.1 ^d	7.2±0.2 ^a	15.8±0.2 ^c	18±0.9 ^c	19.9±0.5 ^d	20.2±0.6 ^d
Shoot fresh weight (mg)	10±0.1 ^a	150±0.1 ^e	203±0.2 ^c	367±0.1 ^d	369±0.1 ^d	8±0.2 ^a	48±0.2 ^c	39±0.3 ^b	68±0.4 ^d	70±0.2 ^d
Root fresh weight (mg)	7±0.03 ^a	220±0.1 ^e	214±0.1 ^c	299±0.2 ^d	301±0.1 ^d	7±0.1 ^a	89±0.2 ^c	79±0.2 ^b	91±0.3 ^c	96±0.1 ^d
Shoot dry weight (mg)	1.9±0.02 ^a	23.6±0.02 ^b	29.2±0.03 ^b	57.1±0.1 ^d	58.3±0.2 ^d	1.7±0.1 ^a	8.1±0.03 ^c	4.9±0.02 ^b	11.8±0.1 ^d	12.1±0.1 ^d
Root dry weight (mg)	1.1±0.1 ^a	32.3±0.2 ^b	32.2±0.02 ^c	43.8±0.04 ^d	44.4±0.3 ^d	1.08±0.1 ^a	13.2±0.2 ^c	12.6±0.2 ^c	14.1±0.1 ^c	14.9±0.2 ^d

565 ^{a-d}: Significant differences between values within the same line (Tukey's test; $p \leq 0.05$).

Table S2. Tukey's post hoc test letters indicating significance of bacterial growth under herbicide stress at each time point.

Strain	Florasulam/2,4-D ($\mu\text{L}/\text{mL}$)	Days							
		1	2	3	4	7	8	9	10
AsEB1	0	b	a	b	b	a	a	a	b
	60	b	a	a	c	a	a	a	b
	150	b	a	b	b	b	b	a	b
	300	a	a	b	a	b	a	a	b
	600	a	a	b	a	a	b	a	a
	1200	a	a	b	a	a	a	a	a
	p-value	0.0002	0.1542	0.0000	0.0000	0.0001	0.0000	0.1248	0.0001
	AsEB2	0	ab	a	ab	a	ab	c	b
60		ab	a	b	b	a	c	b	a
150		b	c	b	b	b	b	a	a
300		a	b	a	ab	ab	c	b	a
600		ab	b	ab	a	ab	a	b	a
1200		ab	b	b	b	ab	a	a	a
p-value		0.0031	0.0000	0.0001	0.0012	0.0002	0.0001	0.0005	0.7245
AsEB3		0	b	b	c	a	c	c	a
	60	b	b	b	a	c	a	c	c
	150	b	b	b	b	c	ab	c	b
	300	a	b	b	c	b	b	b	ab
	600	b	ba	a	a	b	a	c	a
	1200	a		b	b	a	b	b	b
	p-value	0.0001	0.0004	0.0000	0.0000	0.0000	0.0000	0.0000	0.0015
	AsEB4	0	b	a	c	a	b	b	a
60		b	b	b	b	b	ab	a	b
150		b	a	b	b	b	b	a	b
300		a	b	a	b	b	a	a	b
600		a	a	a	b	b	b	a	a
1200		a	b	b	b	a	a	a	b
p-value		0.0004	0.0001	0.0000	0.0001	0.0001	0.0001	0.1142	0.0001

Different letters within the same column indicate significant differences between treatments at the same time point (Tukey's test; $p < 0.05$).

Table S3. Tukey's post hoc test letters indicating significance of bacterial growth under lead stress at each time point.

Strain	Pb (mM)	Days							
		1	2	3	4	7	8	9	10
AsEB1	0	a	a	a	a	a	a	a	a
	1	b	d	c	d	c	c	c	c
	2	c	d	b	cd	c	c	b	b
	3	bc	bc	b	bc	bc	b	bc	bc
	4	b	b	b	b	b	c	bc	bc
	5	b	c	b	b	c	b	bc	b
	p-value	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
AsEB2	0	a	a	a	a	a	a	a	a
	1	c	c	c	a	a	a	a	a
	2	c	c	c	a	a	a	a	a
	3	bc	bc	bc	a	a	a	a	a
	4	b	ab	b	a	a	a	a	a
	5	ab	ab	ab	a	a	a	a	a
	p-value	0.0004	2.33e-05	6.35e-05	0.384	0.234	0.605	0.096	0.197
AsEB3	0	a	a	a	a	a	a	a	a
	1	c	c	bc	c	a	c	bc	c
	2	c	c	c	c	a	c	c	c
	3	bc	bc	c	bc	a	bc	bc	c
	4	b	b	bc	bc	a	bc	c	bc
	5	ab	ab	b	b	a	b	b	b
	p-value	0.0003	0.000	0.000	0.008	0.21	0.0016	0.012	0.000
AsEB4	0	a	a	a	a	a	a	a	a
	1	ab	b	b	ab	a	a	a	b
	2	bc	b	c	b	a	a	a	b
	3	bc	b	c	b	a	a	a	b
	4	ab	b	b	ab	a	a	a	b
	5	c	b	c	b	a	a	a	b
	p-value	0.0003	0.0000	0.0000	0.0382	0.1245	0.1872	0.0912	0.0415


Different letters within the same column indicate significant differences between treatments at the same time point (Tukey's test; $p < 0.05$).

570 **Table S4.** Tukey's post hoc test letters indicating significance of bacterial growth under cadmium stress at each time point.

Strain	Cd (mM)	Days							
		1	2	3	4	7	8	9	10
AsEB1	0	a	a	a	a	a	a	a	a
	1	b	b	a	b	a	a	a	b
	2	b	b	a	b	a	a	a	b
	3	ab	b	a	b	a	a	a	b
	4	b	ab	a	b	a	a	a	b
	5	ab	b	a	b	a	a	a	ab
	p-value	0.0421	0.0156	0.0812	0.0485	0.4125	0.2841	0.0954	0.0388
AsEB2	0	a	a	a	a	a	a	a	a
	1	b	b	b	c	b	b	b	b
	2	ab	b	b	c	b	b	b	b
	3	b	b	b	b	b	b	b	b
	4	ab	b	b	b	b	b	b	b
	5	ab	b	c	c	b	b	b	b
	p-value	0.0042	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
AsEB3	0	a	a	a	a	a	a	a	a
	1	b	b	b	b	b	b	b	b
	2	b	b	b	b	b	b	b	b
	3	b	b	b	b	b	b	b	b
	4	b	b	b	b	b	b	b	b
	5	b	b	b	b	b	b	b	b
	p-value	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
AsEB4	0	a	a	a	a	a	a	a	a
	1	c	b	b	b	b	b	b	b
	2	b	c	c	b	b	b	b	b
	3	bc	b	b	b	b	b	b	b
	4	bc	b	b	b	b	b	b	b
	5	bc	b	c	c	b	b	b	b
	p-value	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

571 Different letters within the same column indicate significant differences between treatments at the same time point (Tukey's test; $p < 0.05$).

572



573

574 **Figure S1.** Effect of AsEB4 strain on germination of tomato seeds after 5 days. (A) Control, (B)
575 Seeds inoculated with AsEB4 under standard condition, (C) Seeds inoculated with AsEB4 under
576 salinity stress (400 mM NaCl).

ACCEPTED MANUSCRIPT

577 **List of tables**

578 **Table 1.** Phenotypic profiles of isolated *Bacillus* spp. from garlic roots.

579 **Table 2.** Drought and salinity tolerance of endophytic *Bacillus* spp. isolated from garlic roots.

580 **Table 3.** Effect of AsEB4 strain on germination and seedling growth of tomato under saline condition.

581 **Table 4.** Non-enzymatic antioxidant activities of endophytic *Bacillus* spp. isolated from garlic roots.

582 **List of figures**

583 **Figure 1.** Growth of endophytic *Bacillus* spp. isolated from garlic roots under florasulam/2,4-D
584 herbicide stress.

585 **Figure 2.** Growth of endophytic *Bacillus* spp. isolated from garlic roots under lead stress.

586 **Figure 3.** Growth of endophytic *Bacillus* spp. isolated from garlic roots under cadmium stress.

587 **Figure 4.** Colony development of heavy metal-tolerant *Bacillus* spp. isolated from garlic roots at the
588 maximum tolerance concentration of lead and cadmium.

589 **Figure 5.** Exopolysaccharides production by endophytic *Bacillus* spp. strains under lead and
590 cadmium stresses. ^{a-d}: indicate significant differences (Tukey's test; p<0.05).

591 **Figure 6.** Effect of endophytic *Bacillus* spp. strains on the mycelial growth of *Botrytis cinerea* after
592 7 days at 28 ± 0.2 °C. (A) Colony growth of *B. cinerea* on the PDA medium as control; (B) Inhibition
593 of *B. cinerea* growth by AsEB1; (C) Inhibition of *B. cinerea* growth by AsEB4.

594 **Supplementary Material**

595 **Table S1.** Effect of endophytic *Bacillus* spp. isolated from garlic roots on germination and seedling
596 growth of wheat and tomato seeds after 14 days.

597 **Table S2.** Tukey's post hoc test letters indicating significance of bacterial growth under herbicide
598 stress at each time point.

599 **Table S3.** Tukey's post hoc test letters indicating significance of bacterial growth under lead stress
600 at each time point.

601 **Table S4.** Tukey's post hoc test letters indicating significance of bacterial growth under cadmium
602 stress at each time point.

603 **Figure S1.** Effect of AsEB4 strain on germination of tomato seeds after 5 days. (A) Control, (B)
604 Seeds inoculated with AsEB4 under standard condition, (C) Seeds inoculated with AsEB4 under
605 salinity stress (400 mM NaCl).

ACCEPTED MANUSCRIPT