

Short-term effects of cadmium exposure on energy reserves and stress markers on *Ophelia bicornis* Savigny, 1822 (Polychaeta: Opheliidae)

Belabed Soumeya^{1*}, Labiod Ryma², Boukrouma Nadhra³, Mhadhbi Lazhar⁴ and Soltani Noureddine¹

¹Laboratory of Applied Animal Biology (LBAA), Department of Biology, Faculty of Sciences, Badji Mokhtar University, 12, P.O. Box, 23000 Annaba, Algeria.

²Laboratory of Organic Chemistry and Interdisciplinary (LOCI), University of Souk-Ahras, P.O. Box 1553, 41000 Souk Ahras, Algeria.

³Laboratory of Aquatic and Terrestrial Ecosystems (LEAT), Department of Biology, Faculty of Natural and Life Sciences, Mohamed Chérif Messaâdia University, BP1553, Annaba Road, 41000 Souk Ahras, Algeria.


⁴Laboratory of Ecology, Biology and Physiology of Aquatic Organisms, Faculty of Sciences of Tunis, University Tunis El Manar, 2092, Tunis, Tunisia.

Received: 29/09/2025, Accepted: 12/03/2026, Available online: 18/03/2026

*to whom all correspondence should be addressed: e-mail: s.belaabed@univ-soukahras.dz

<https://doi.org/10.30955/gnj.08057>

Graphical abstract



Abstract

Ophelia bicornis (Polychaeta: Opheliidae) has been identified as a potential bioindicator for ecotoxicological studies. This study evaluates the short-term physiological and biochemical effects of cadmium (Cd) exposure on *O. bicornis* by examining energy reserves (lipids, carbohydrates, proteins) and oxidative stress biomarkers (hydrogen peroxide, lipid peroxide (LOOH), and Superoxide dismutase (SOD)). Cd treatment was conducted at a sublethal concentration LC₁₀ obtained after 48h of exposure (31.11 mg/L) as previously determined. Statistical analyses revealed significant interactions between Cd treatment and exposure time affecting several biochemical parameters. Amounts of Carbohydrates increased notably after 48h in Cd-exposed individuals, indicating time-dependent energy mobilization. Lipid contents were also affected by

treatment and exposure duration, reflecting energy reserve use under metal stress. In contrast, protein remained stable, with no significant variation across treatment or time. Oxidative stress biomarkers were markedly influenced by Cd exposure. LOOH levels increased significantly with treatment and time, peaking at 48h. Similarly, H₂O₂ increased with Cd and exhibited a significant time-dependent interaction, highlighting a progressive oxidative challenge. SOD activity was primarily modulated by exposure duration, reaching its maximum at 48h. Overall, the present experiment shows that sublethal Cd disrupts energy metabolism and triggers oxidative stress response in *O. bicornis*.

Keywords: *Ophelia bicornis*, Bioindicator, Toxicity, Energy metabolism, Oxidative stress, Biomarkers.

1. Introduction

In recent years, it has become evident that, to fully assess the impact of anthropogenic trace metal inputs on aquatic ecosystems, it is essential to understand the chemical and physiological processes that govern metal accumulation in organisms, as well as how these metals influence energy reserves and antioxidant activities. Many toxicants produce a sublethal effect by binding to or interfering with essential cellular components such as enzymes and metabolites, thereby disrupting fundamental processes within the organism (Holmstrup *et al.* 2010). They can disrupt energy metabolism, deplete energy stores, and induce oxidative stress, ultimately affecting overall health and survival of aquatic organisms (Song *et al.* 2023). These substances are particularly known for their toxic effects, persistence, and tendency to transfer and accumulate

Belabed Soumeya, Labiod Ryma, Boukrouma Nadhra, Mhadhbi Lazhar and Soltani Noureddine (2026), Short-term effects of cadmium exposure on energy reserves and stress markers on *Ophelia bicornis* Savigny, 1822 (Polychaeta: Opheliidae), *Global NEST Journal*, 28(2), 08057.

Copyright: © 2026 Global NEST. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution International (CC BY 4.0) license.

through all the trophic levels (Di Salvatore *et al.* 2013). Trace metals are among the most extensively studied groups of xenobiotics (Pan and Wang 2012). Besides, various chemical substances, including cadmium, can exert multiple mechanisms of toxicity, each contributing to the overall harmful effects (Valavanidis *et al.* 2006; Regoli and Giuliani 2014). Cadmium (Cd) is a heavy metal hazardous to all living organisms, and its effects have been studied on various organisms (Sharma *et al.* 2015). It is a non-essential element that can be toxic even at trace levels for aquatic species. Cadmium is a highly toxic environmental pollutant that poses a significant threat due to its tendency to accumulate in marine ecosystems, its ability to easily cross cell membranes, and its persistence in the environment, which can be found at high concentrations in seawater (Amira *et al.* 2018). Polychaetes play an important role in the functioning of benthic communities (Hutchings 1998). They represent 35 to 50% of benthic macrofauna and are often the numerically dominant macrobenthic taxon. Found in coastal and littoral ecosystems, where the distribution of species in soft-bottom habitats is primarily influenced by sediment particle size (Gambi and Giangrande 1986) and organic matter content, they are frequently exposed to various contaminants. Due to their sensitivity and ecological significance, polychaetes serve as excellent indicators for studying the effects of pollutants and the transfer of xenobiotics through food chains (Dean 2008). Consequently, they have been extensively used in coastal studies for environmental monitoring and are considered a key group in ecotoxicological research (Scaps 2000). Furthermore, Polychaete Annelids are a major link in food webs and are widely consumed by many species, such as crustaceans, fishes, and birds. Among them, the endobenthic species *Ophelia bicornis* (Savigny 1818) plays a crucial role in evaluating the structure and functioning of estuarine ecosystems. This polychaete is particularly notable for its physiological tolerance to various environmental stressors, especially its resistance to sediment-bound contaminants. Accordingly, reduced energy reserves and elevated metabolic rates in animals have associated to metal toxicity, especially in the case of cadmium, Zinc and copper (Pook *et al.* 2009). Recent studies have explored advanced analytical and computational approaches for environmental pollution assessment. Deep learning and optimization-based models have been applied to classify and predict air pollution levels and analyze environmental data (Mohandas *et al.* 2025b, 2025a). Additionally, hybrid recurrent neural network models combined with Internet of Things (IoT) data have been proposed for urban air pollution prediction (Mohandas *et al.* 2025c). Beyond atmospheric pollution, graph-based and remote sensing-driven approaches have been developed for environmental monitoring (Sivasubramanian *et al.* 2025), and early detection of ecological stress, such as pest disease identification in agricultural systems (Maruthai *et al.* 2025; Mohandas *et al.* 2025a). While these studies provide valuable large-scale insights, experimental ecotoxicological investigations remain essential for

understanding species-specific biochemical responses to metal exposure. *Ophelia bicornis*, which commonly inhabits organic-rich environments, is considered a key species for ecological assessments and a reliable bioindicator for monitoring heavy metal pollution (Bat *et al.* 2016). In this context, *Ophelia bicornis* could be effectively used as a bioindicator species in ecotoxicological studies of marine and coastal ecosystems along the eastern Algerian coast to assess metal contamination (Labiod and Belabed 2024). This species is distributed across the North, Mediterranean, Black Sea, and Atlantic coasts of North Africa and Europe, typically inhabiting high-energy intertidal zones and fine to medium sediments (Maltagliati *et al.* 2005).

The presence of a xenobiotic in the environment always poses a risk to living organisms. However, the relationship between toxic concentrations and the toxic response is fairly complex and difficult to predict, as it depends on several factors, including toxicokinetics and genetic variability. One method of quantifying xenobiotic interaction and its potential impact on living organisms is through biomarker monitoring. Therefore, biomarkers measure exposure, toxicity, and individual susceptibility to environmental chemicals, making them valuable tools for assessing and monitoring the risk of long-term effects associated with exposure to xenobiotics (Depledge 2020). Generally, biochemical biomarkers are frequently used to identify early signs of stress or toxicity in benthic fauna (Mayer *et al.* 2018; Depledge 2020). One of the main consequences of metal toxicity is additional energy costs, and the resulting metabolic load can lead to disruption in oxidative metabolism and increased anaerobic activity (Gashkina 2024). Furthermore, metal toxicity mainly involves changes at biological levels of cellular activities, including reactive oxygen species (ROS) generation, which can cause damage associated with the different pathological processes (Makhdoumi *et al.* 2020). Likewise, Cd accumulation has also been reported to alter key biochemical and physiological functions (Wright and Welbourn 1994) and to induce the production of reactive oxygen species (ROS) such as hydrogen peroxide (H₂O₂), and lipid peroxide (LOOH) (Benedetti *et al.* 2015). A significant increase in the tissue concentration appeared 48h after exposure in *Donax trunculus*, following treatment with LC₅₀ (2.59 mg/L) and LC₂₅ (1.32 mg/L) of Cd (Belabed and Soltani 2018). Moreover, the presence of cadmium chloride (CdCl₂) in seawater led to mortality in the marine benthic species *Donax trunculus* at concentrations corresponding to LC₁₀ (0.72 mg/L) and LC₅₀ (2.59 mg/L) after 96h of exposure, and also affected the biomarker responses such as acetylcholinesterase (AChE) and Catalase (CAT) activities (Belabed and Soltani 2022). Thus, some biomarkers such as Superoxide dismutase, catalase, glutathione peroxidase, glutathione, and glutathione reductase are indicative of oxidative stress caused by exposure to pro-oxidants (Demirci-Çekiç *et al.* 2022), whereas metallothionein contents are widely used as biomarkers of metal contamination by binding and removing toxic metals (Cosson and Amiard 2021). These antioxidant defenses aim to protect cells and tissues from

oxidative damage by neutralizing the toxicity of ROS (Singh *et al.* 2024).

In previous studies (Belabed and Soltani 2018, 2022; Merad *et al.* 2024), Cd was tested on a Molluscan species, *Donax trunculus*, widely used as bioindicator of marine pollution. The present study aimed to investigate the short-term effects (24, 48h) of cadmium exposure at a sublethal concentration LC₁₀ (31.11 mg/L) (Labioud and Belabed 2024) in the marine organism *O. bicornis*, to assess the extent of metal contamination. Acute toxicity data on *O. bicornis* and its responses to toxic substances remain scarce in scientific literature. In light of the limited data available, a battery of stress indices was employed, including SOD activity, the generation of free radicals, and the biochemical composition. Characterization of the response of stress indices could determine the possible adverse effects of cadmium on worms and evaluate the potential harmful impacts on aquatic environments.

2. Materials and methods

2.1. Sampling and contamination protocol

The annelid *O. bicornis* was collected at low tide by hand from littoral sediments in the intertidal zone (depth greater than 20 cm) at El Hennaya Beach (36°54'24.00" N, 08°7'43.62" E), that is located in the district of Berrihane, which belongs to the province of El Tarf (Northeast Algeria). This site is not exposed to any source of pollution because of its location, which is quite remote from the various discharges. Once in the laboratory, the samples were transported in plastic buckets with sediment from the collection site, and they were acclimatized for 48 hours. Afterwards, the collected worms were separated from the residue, cleaned from debris with seawater from the site of origin of worms, and they were transferred to the glass Petri dishes containing filtered natural seawater for another period of acclimatization (24h) in the dark at a temperature of 16°C. This acclimatization phase establishes stable experimental conditions and enhances the accuracy of toxicity test results, reflecting the actual effect of the toxic substance on organisms. Sublethal concentrations of Cd (LC₁₀) were administered to lots (n = 30 for each treatment). Thus, every 24 hours throughout the exposure period, worms (0.3 g in weight) were sampled and immediately homogenized (10% w/v) in Tris-HCl buffer (100 mM; pH = 7.4). The homogenates were centrifuged at 9000×g for 20 min at 4 °C. The supernatant was then preserved at -80°C until biochemical analysis. Analyses were performed to assess the total lipid, carbohydrate, and protein contents as well as oxidative stress responses by measuring Superoxide dismutase (SOD) activity and quantifying Hydrogen peroxide (H₂O₂) and Lipid hydroperoxides (LOOHs) levels. Assays were conducted on six individuals in both the treated and control groups, spanning a treatment period of 48 hours.

2.2. Energy reserve analysis

In this experimental procedure, biochemical analyses concerned only the animals that survived the two days of Cd exposure. The extraction of protein, carbohydrate and lipid contents is determined according to Shibko *et al.*

(1966). The method involves treating a homogenate by adding 1 ml of 20% trichloroacetic acid (TCA) to the sample. After grinding with ultrasound (Sonifier B-30), centrifugation at 5000 × g for 10 minutes at 4°C yields a supernatant 1 that is collected use in the total carbohydrate analysis, while the pellet 1 undergoes washing with 1 ml chloroform ether (1V/1V), centrifuged again (5000 g for 10 min). To determine the carbohydrate content of the supernatant fraction, 4 ml of anthrone was added to 100 µl of supernatant, and quantification was performed by measuring absorbance at 620 nm against a standard curve of glucose standard (Sigma, USA) (Duchateau and Florin 1959). The supernatant 2 recovered will be used for the determination of the total lipids based on the vanillin method of (Goldsworthy *et al.* 1972) and absorbance of the samples was measured at 530 nm. The remaining pellet 2 was re-suspended in 1ml NaOH (0,1 N) used for assessing the total protein content (Bradford, 1976a) (Sigma, USA). The absorbance was measured at 595 nm using bovine serum albumin (Sigma, USA) as a standard. The contents of carbohydrates, proteins and lipids were expressed in µg/mg of body weight.

2.3. Superoxide dismutase (SOD) activity

The superoxide dismutase (SOD) activity was determined by monitoring the photochemical reduction of NBT according to the method of Beauchamp and Fridovich (1971). One unit of SOD activity corresponded to the amount of enzyme required to cause 50% inhibition of NBT reduction at 560 nm measured by a spectrophotometer. The SOD activity was expressed as U/mg of protein.

2.4. Hydrogen peroxide (H₂O₂)

Analysis of hydrogen peroxide (H₂O₂) levels were carried out by the method described by Ou and Wolff (1996), using the ferrous ion oxidation xylenol orange assays (FOX1). The amount of H₂O₂ in the supernatant was determined at 560 nm using a spectrophotometer. Values were expressed as mmol/mg of protein.

2.5. Lipid hydroperoxides (LOOHs) content

Lipid hydroperoxides (LOOH) formation was estimated according to the method described by Jiang *et al.* (1992) using the ferrous oxidation in xylenol orange assay. Hydroperoxides content was colorimetrically determined at 560 nm. Results were expressed as mmol/ mg of protein.

2.6. Total Protein quantification

Protein content was done by the method of Bradford (1976), which is based on the binding of Coomassie brilliant blue G -250, present in the Bradford reagent, to the total protein present in the sample, yielding a stable staining compound. The presence of this compound can be spectrophotometrically quantified at 595 nm.

2.7. Statistical analysis

All statistical analyses were conducted using R software (version 4.5.0; R Core Team 2025). Prior to hypothesis testing, data normality and homogeneity of variances

were assessed using the Shapiro–Wilk test and Levene’s test, respectively. **For carbohydrate content**, which met normality and homogeneity assumptions ($p > 0.05$), **two-way ANOVA** was applied to evaluate the effects of Exposure Time, Treatment, and their interaction. Significant effects were followed by **Tukey’s HSD post hoc test** for pairwise comparisons while controlling for multiple testing. **For all other measured biomarkers** (lipids, proteins, LOOH, H_2O_2 , and SOD), which did not meet normality assumptions, the **non-parametric Scheirer–Ray–Hare test** was used as a factorial alternative to ANOVA, with significant effects further analyzed using **Dunn’s test** for multiple comparisons. Exploratory data analysis was performed to visually assess variable distributions across experimental groups defined by Exposure Time and Treatment. **Boxplots** summarizing median, interquartile range, overall dispersion, and potential outliers were generated using the *ggplot2* package (version 3.4.2; Wickham 2016), supporting the evaluation of model assumptions. **Bar plots with error bars (mean \pm SD)** were used to facilitate visual comparison of central tendencies and variability. To explore overall variation patterns and relationships among biomarkers, **Principal Component Analysis (PCA)** was performed using the *FactoMineR* package (version 2.7; Lê et al. 2008), and results were visualized with the *factoextra* package (version 1.0.7; Kassambara and Mundt 2020). **Redundancy Analysis (RDA)** via the *vegan* package (version 2.6-4; Oksanen et al. 2022) quantified the proportion of variance in biomarker responses explained by Exposure Time and Treatment. All results are presented as mean \pm standard deviation (SD), and the number of biological replicates used in each assay is reported in the Results section. Statistical significance was set at $p < 0.05$.

3. Results

3.1. Changes in the energy reserves

All energy reserve components exhibited large variations. Generally, changes in energy reserve responses are detected after prolonged exposure, over several hours or days. Proteins were quantitatively the most important energy fractions in the biochemical composition of *O. bicornis* (28,88 $\mu\text{g}/\text{mg}$ of tissue), whereas carbohydrates and lipids represented respectively only 3,93 and 1,93 $\mu\text{g}/\text{mg}$ of tissue.

3.1.1. Carbohydrates contents

The values ($\mu\text{g}/\text{mg}$ of Tissue) obtained in *Ophelia bicornis* are presented in **Figure 1**. The two-way ANOVA revealed a marginally significant interaction between the factors Time and Treatment ($p = 0.0569$), suggesting that the effect of the treatment on Carbohydrates content depends on the duration of exposure. No significant main effects were detected for Time ($p = 0.0672$) or for Treatment ($p = 0.1534$), indicating that these factors alone do not account for the observed variations (**Figure 1A**). However, the interaction between these two factors highlights a combined influence on the measured response.

Post hoc analysis using Tukey’s test showed a significant increase in Carbohydrates after 48 hours in the Cd treated series compared to the initial time point (0 hours) ($p = 0.0287$), indicating a Time-dependent effect of the treatment (**Figure 1B**). In contrast, no significant differences were observed within the controls group over time, confirming that the variations in concentration over time are primarily attributable to the treatment.




Figure 1. Effect of cadmium on carbohydrate content ($\mu\text{g}/\text{mg}$ of Tissue) in *O. bicornis* ($n = 6$) (A) Boxplot showing mean carbohydrate content after exposure to cadmium (mean \pm SD). (B) Error bar plot representing the standard deviation of carbohydrate content (mean \pm SD).

3.1.2. Lipids contents

The Scheirer–Ray–Hare test, was used to assess the effects of Exposure Time and Treatment on Lipid contents ($\mu\text{g}/\text{mg}$ of Tissue) in 36 observations. The test revealed a statistically significant effect of Treatment ($H = 4.91$, $p = 0.0267$) and a significant Time \times Treatment interaction ($H = 7.26$, $p = 0.0265$), suggesting that the impact of treatment depends on exposure duration (**Figure 3**). However, the main effect of Time alone was not statistically significant ($H = 4.51$, $p = 0.1048$) (**Figure 2C**).

Post-hoc Dunn tests with Bonferroni correction showed a significant difference in lipids between Cd-treated series and Control groups ($Z = -2.22$, $p = 0.0267$), while no significant pairwise differences were found between time points, including between 24 and 48 hours (adjusted $p = 0.1197$) (**Figure 2D**).




Figure 2. Effect of cadmium on lipid content ($\mu\text{g}/\text{mg}$ of Tissue) in *O. bicornis* ($n = 6$). (C) Boxplot showing mean lipid content after exposure to cadmium (mean \pm SD). (D) Error bar plot representing the standard deviation of lipid content (mean \pm SD).

3.1.3. Proteins contents

The Scheirer-Ray-Hare test, was used to examine the effects of Time, Treatment, and their interaction on Proteins contents ($\mu\text{g}/\text{mg}$ of Tissue) across 36 observations. The results showed that none of the factors reached statistical significance at the 0.05 level (**Figure 3E**). The effect of Time on Proteins ($\mu\text{g}/\text{mg}$ of Tissue) was not significant ($H = 1.7643$, $p = 0.41388$), indicating that Protein's expression did not vary significantly across the different Time points.

Similarly, the Treatment factor was also not significant ($H = 2.6043$, $p = 0.10658$), suggesting no strong evidence that the treatment alone influenced protein levels. The interaction between Time and Treatment was likewise non-significant ($H = 3.5013$, $p = 0.17366$), meaning there was no combined effect of Time and Treatment on Proteins ($\mu\text{g}/\text{mg}$ of Tissue) (**Figure 3F**).




Figure 3. Effect of cadmium on protein content ($\mu\text{g}/\text{mg}$ of Tissue) in *O. bicornis* ($n = 6$). (E) Boxplot showing mean protein content after exposure to cadmium (mean \pm SD). (F) Error bar plot representing the standard deviation of protein content (mean \pm SD).

3.1.4. H₂O₂ levels (mmole/mg of Proteins)

The Scheirer-Ray-Hare test revealed a marginally non-significant effect of Time on H₂O₂ Levels (mmole/mg of Proteins) ($H = 5.39$, $p = 0.0676$), a significant effect of Treatment ($H = 8.02$, $p = 0.0046$), and a significant Time \times Treatment interaction ($H = 9.33$, $p = 0.0094$) (**Figure 4G**).

Post-hoc Dunn tests with Bonferroni correction showed that H₂O₂ Levels (mmole/mg of Proteins) were significantly higher in the Cd-treated group compared to controls ($p = 0.0046$). Regarding Time, no pairwise comparisons reached significance after correction, although the difference between 0 and 48 hours approached significance ($p = 0.0945$). These results indicate that treatment and the interaction between Time and Treatment significantly influence H₂O₂ levels (mmole/mg of Proteins), with Cd treatment increasing levels overall and a time-dependent variation depending on treatment (**Figure 4H**).

3.1.5. LOOH levels (mmole/mg of Proteins)

The Scheirer-Ray-Hare test showed significant effects of Time ($H = 14.45$, $p = 0.00073$) and treatment ($H = 5.71$, $p = 0.01690$) on LOOH levels (mmole/mg of Proteins), while the interaction between Time and Treatment was not significant ($H = 1.31$, $p = 0.52$) (**Figure 5I**).

Post-hoc Dunn tests with Bonferroni correction revealed that LOOH levels (mmole/mg of Proteins) at 48 hours were significantly higher than at 0 hours ($p = 0.0049$) and 24 hours ($p = 0.0019$), with no difference between 0 and 24 hours. Additionally, Cd-treated samples showed significantly higher LOOH levels (mmole/mg of Proteins) compared to controls ($p = 0.0169$). These results indicate that both Time and Treatment independently affect LOOH levels (mmole/mg of Proteins), with increases observed at 48 hours and following Cd exposure (**Figure 5, J**).




Figure 4. Effect of cadmium on H₂O₂ levels (mmole/mg of Proteins) in *O. bicornis* ($n = 6$). (G) Boxplot showing mean H₂O₂ levels after exposure to cadmium (mean \pm SD). (H) Error bar plot representing the standard deviation of H₂O₂ levels (mean \pm SD).




Figure 5. Effect of cadmium on LOOH levels (mmole/mg of Proteins) in *O. bicornis* ($n = 6$). (I) Boxplot showing mean LOOH levels after exposure to cadmium (mean \pm SD). (J) Error bar plot representing the standard deviation of LOOH levels (mean \pm SD).

3.1.6. SOD Activity (U/mg of Proteins)

The Scheirer-Ray-Hare test showed a significant effect of Time on SOD Activity (U/mg of Proteins) ($H = 12.71$, $p = 0.00174$), while neither Treatment ($H = 2.26$, $p = 0.13286$) or the Time \times Treatment interaction ($H = 1.99$, $p = 0.36975$) were significant (**Figure 6K**).

Post-hoc Dunn tests with Bonferroni correction indicated that SOD activity at 48 hours was significantly different from 0 hours ($p = 0.0012$), while differences between 0 and 24 hours and between 24 and 48 hours were not statistically significant after adjustment ($p = 0.51$ and $p = 0.09$, respectively). No significant difference was found between Cd-treated series and control groups ($p = 0.13$). These results suggest that SOD Activity (U/mg of Proteins) changes significantly over Time, particularly between 0 and 48 hours, but is not significantly affected by treatment or the interaction of treatment and Time (**Figure 6L**).




Figure 6. Effect of cadmium on SOD activity (U/mg Of Proteins) in *O. bicornis* (n = 6). (K) Boxplot showing mean SOD activity after exposure to cadmium (mean \pm SD). (L) Error bar plot representing the standard deviation of SOD activity (mean \pm SD).



Figure 7. Principal Component Analysis (PCA) Biplot of Biochemical Biomarkers in Relation to Treatment and Exposure Time in *Ophelia bicornis*. Legend: The PCA Biplot illustrates the distribution of biochemical parameters (n = 6) measured in Tissue samples exposed to Cd-Treated series and control conditions over varying Exposure Times. Variables include Carbohydrates contents ($\mu\text{g}/\text{mg}$ of Tissue), Lipids ($\mu\text{g}/\text{mg}$ Of Tissue), Proteins ($\mu\text{g}/\text{mg}$ of Tissue), H₂O₂ levels (mmole/mg Of Protein), LOOH levels (mmole/mg Of Protein), and SOD activity (U/mg Of Protein). These variables are represented as vectors, where their direction and length indicate the magnitude and contribution to the principal components. Sample groups, defined by Treatment (Controls vs. Cd-Treated series) and Exposure Time (0, 24, 48 hours), are displayed as points with colored 95% confidence ellipses delineating group clustering. Points oriented in the direction of a vector suggest higher values of the corresponding biomarker. (PC1: 43.3%; PC2: 19.5%).

3.2. Principal Component Analysis (PCA) and Redundancy Analysis (RDA) of Biochemical Biomarkers in *Ophelia bicornis*: Effects of Exposure Time and Treatment

The combined application of PCA and RDA provided complementary insights into the biochemical responses to treatment and exposure time. PCA, an unconstrained ordination method, revealed that the first two principal components explained 62.82% of the total variance (PC1:

43.3%; PC2: 19.5%). PC1 was showing strong positive correlations with H₂O₂ levels ($r = 0.89$, $p < 0.001$), Carbohydrate ($r = 0.71$, $p < 0.001$), and LOOH levels ($r = 0.70$, $p < 0.001$). This axis was positively associated with the Cd-treated group, particularly at 48 hours. PC2 was mainly correlated with Lipid ($r = 0.74$, $p < 0.001$) and SOD activity ($r = 0.49$, $p = 0.003$) and was significantly associated with the 48-hour Exposure Time (Figure 7).

In parallel, RDA (constraining the ordination by Treatment and Exposure Time explained 47.3% and 6% of the variance on Axis 1 and Axis 2, respectively). Axis 1 was positively associated with H₂O₂ levels ($r = 0.33$, $p < 0.001$) and SOD activity ($r = 0.12$, $p < 0.01$), and negatively with protein ($r = -0.19$) and lipid ($r = -0.08$), indicating increased oxidative stress in exposed samples. Axis 2 was primarily correlated with Lipid ($r = 0.13$, $p < 0.01$) and SOD activity ($r = 0.09$, $p < 0.05$), reflecting Time-dependent metabolic responses (Figure 8).




Figure 8. Redundancy Analysis (RDA) Biplot of Biochemical Biomarkers Constrained by Cadmium Treatment and Exposure Time. Legend: The RDA Biplot shows the relationship between biochemical parameters (Carbohydrates, Lipids, Proteins, H₂O₂ levels, LOOH levels, and SOD activity) and the explanatory variables Treatment (Controls vs. Cd-Treated series) and Exposure Time (0, 24, 48 hours). Biochemical variables are represented as vectors indicating their strength and direction of influence on the ordination axes. Sample points are grouped according to treatment and exposure duration, with 95% confidence ellipses highlighting clustering patterns. The first two RDA axes explain 27% and 5.1% of the variance, respectively. Vectors pointing toward sample clusters indicate higher biomarker expression under those conditions.

4. Discussion

Short-term tests were designed to measure not only mortality rates, but also to determine tolerance ranges of metabolic systems most sensitive to metal stress. Toxicity tests are necessary in water pollution evaluation, as chemical and physical parameters are insufficient to determine potential effects on aquatic organisms.

Studying the responses of organisms to environmental stressors plays a key role in ecotoxicological studies. The established lethal concentration values could be used as key indicators in toxicity tests to demonstrate the effects of toxic pollutants, particularly for ecotoxicological evaluation of marine ecosystems in risk assessments. This work aims to evaluate the toxic effects of cadmium and its capacity to alter several variations in organism function. Namely, the estimation of biochemical contents, and the evaluation of enzymatic activities in *O. bicornis* species. The most toxicants exert their effects at basic level of the organism by reacting with enzymes or metabolites and other functional components of the cell (Ho Yu 2000). Thus, organisms' energy reserves provide a valuable indicator of their ability to cope with stress long before negative effects appear at more complex biological levels (Smolders *et al.* 2003). To examine energy metabolism perturbations, many bioenergetic parameters are available such as levels of energy reserves as carbohydrates, lipids and proteins (Meyer *et al.* 2002), cellular energy allocation (De Coen and Janssen 2003), scope for growth (Widdows *et al.* 2002). This research represents the first attempt to evaluate cadmium toxicity using *Ophelia bicornis* as a test organism. In this context, the energy status of *O. bicornis* was studied to gain insight into Cd-induced lethal effects and the ability of cadmium to induce disturbances on different levels of organism function. Cadmium is a very toxic metal, and even though it is unable to undergo Fenton-type reactions and therefore is considered a non-redox metal, and it has nevertheless been associated to increased production of reactive oxygen species (ROS) to marine organisms (Unsal *et al.* 2020; Liu *et al.* 2022).

Our results revealed that cadmium exposure and duration both influenced the biochemical responses in *O. bicornis*. Energy-related compounds such as carbohydrates and lipids were mobilized over time in the cadmium-treated series compared to the control group, reflecting a physiological adjustment to metal-induced stress. However, protein levels remained relatively constant, suggesting that they were not significantly affected by treatment or exposure time. These findings are consistent with the notion that the earliest toxic effects of chemicals manifest primarily at the cellular and biochemical levels. As highlighted by Zhou *et al.* (2004), recent studies have increasingly focused on measurement at these levels to assess whether exposure to pollutants causes harmful effects. The observed changes in carbohydrate and lipid contents, along with the stability of protein levels, reflect such early biochemical responses to cadmium-induced stress. The biochemical composition (carbohydrates, lipids and proteins) indeed varies between taxonomic groups, but also among species, developmental stages, analyzed tissues, and environmental conditions. Moreover, this study suggests that the main storage materials in *O. bicornis* are proteins, whereas carbohydrates and lipids were present at a relatively low level.

The increase in carbohydrates levels after 48h of exposure likely reflects a stress-induced mobilization of

glycogen or other carbohydrate stores to provide readily available energy for detoxification processes. The observed increase in carbohydrate levels in Cd-exposed *Ophelia bicornis* may reflect a typical stress response in marine invertebrates. However, the utilization of carbohydrates by aquatic animals is relatively low compared to that of terrestrial animals. In marine invertebrates, carbohydrates are often the first source of energy utilized during acute stress, as they can be quickly metabolized through glycolysis to produce ATP (Wang *et al.* 2016). Increases in circulating glucose in the hemolymph have been reported in response to a variety of stressors such as salinity, temperature, hypoxia, or exposure to heavy metals (Wang *et al.* 2012). Notably, exposure to cadmium induced in the crayfish *Procambarus clarkii* a significant increase in hemolymph glucose concentration (Reddy *et al.* 1994). Carbohydrates and lipids play an important role in building up reserves for embryo development in Annelids. At the same time, the reproductive effort is high with up to 80% of the total energy allocated to gamete biomass (Hoeger and Schenk 2024). With increasing levels of stress, the maintenance of physiological integrity becomes more challenging for organisms. As an immediate response to the toxicity, organisms tend to reduce their metabolic rate (Hand and Hardewig 1996). Some studies have suggested that the lipid contents of the exposed animals were significantly decreased compared to the control (Lucia *et al.* 2010; Yang *et al.* 2013), and this was confirmed by our study. In addition to carbohydrates, total lipid content significantly decreased in *Ophelia bicornis* under cadmium exposure, particularly after 48h, suggesting a depletion of lipid reserves in response to metal-induced stress. This finding is consistent with previous studies in crustaceans such as crabs, where lipid levels in the hepatopancreas and ovary decreased after prolonged cadmium exposure compared to controls (Liu *et al.* 2016). Lipid reserves are mobilized or degraded under metal-induced stress, likely to meet elevated energy demands or due to oxidative damage. Luis and Passos (1995) found that lipids are crucial during the life cycle of *N. diversicolor*, particularly for reproduction. Lipids accumulate as an energetic reserve in oocytes and support metabolism during gametogenesis when the animals stop feeding. In aquatic invertebrates, reductions in the sequestration of reserves have been shown to affect the number or quality of gametes produced (Mathieu and Lubet 1993).

In contrast, protein content remained relatively stable. The stability in protein levels suggests that *O. bicornis* reserves proteins, avoiding their breakdown unless under severe or prolonged stress. Proteins were not preferentially mobilized as energy sources under these exposure conditions. Similarly, according to the hypothesis proposed by Ketata *et al.* (2007), proteins are not primarily involved in energy metabolism in the clam *Ruditapes decussatus* exposed to cadmium. This observation aligns with our results, which also revealed no significant change in protein levels across treatment conditions or exposure durations. This contrasts with the findings of Barber and Blake (2006) reported that proteins

can serve as a significant energy source, particularly when other reserves are depleted. Bivalves store energy primarily as carbohydrates (mainly glycogen) and lipids to maintain physiological integrity, during reproductive periods, whereas annelids tend to rely more on proteins to support their fundamental metabolic activities, it has been shown that proteins are the main components of muscle tissue. According to the research of (Tripp-Valdez *et al.* 2019) proteins, complex macromolecules described as biopolymers, are the most abundant of cellular organic molecules. Under severe stress conditions, proteins may serve as an alternative energy source. However, this process compromises their structural and functional roles, since proteins are not primarily synthesized or stored for energy supply (Le Gal *et al.* 1997). On the other hand, Cd exposure has been shown to alter protein metabolism in bivalves, affecting both synthesis and degradation pathways in various tissues. The effects of cadmium on protein levels were tissue-specific and varied with duration or exposure (Ivanina *et al.* 2008). Cadmium is known to affect reproduction and energy metabolism in marine organisms (Baudou *et al.* 2017; Louis *et al.* 2021). Concurrently, oxidative stress-induced protein oxidation can result in structural alterations and loss of function, disrupting cellular processes dependent on properly functioning proteins (Boguszewska-Mańkowska *et al.* 2015). A significant decrease in lipids, proteins and carbohydrates contents in muscles and gills of *Carcinus Aestuarii* as compared to *Portunus segnis* has been observed (Chetoui *et al.* 2021). Thus, seasonal changes can influence the physiology of crabs by affecting their metabolic activity and response, leading to disturbances in biochemical components such as lipids, proteins and carbohydrates. These alterations appear to be linked to the accumulation of trace elements, environmental factors and the reproductive cycles of aquatic organisms (Chetoui *et al.* 2021).

At the cellular level, oxidative stress plays an important role in the development of damage. Reactive oxygen species (ROS) can be free radicals derived from oxygen, such as superoxide anion (O_2^-) and hydroxyl radicals (OH), or non-radical derivatives such as hydrogen peroxide (H_2O_2). These molecules are produced through several mechanisms (Fouzai *et al.* 2020). While ROS play essential roles in cellular functions, their excessive accumulation beyond the antioxidant capacity of the organism causes oxidative damage to biomolecules, leading to disruptions in cellular functions and structures (Sinenko *et al.* 2021). The degradation of these radicals is controlled by antioxidants systems, which adapt to the level of radicals present. The antioxidant system of marine organisms consists of low molecular weight scavengers and antioxidant enzymes which interact in a sophisticated network. Several studies on aquatic organisms have demonstrated the importance of antioxidant enzymes in protecting cellular systems from oxidative damages induced by xenobiotic. The impact of oxidative stress has been shown to extend beyond individual cellular processes, influencing various downstream effects of metal toxicity (Chen *et al.* 2018). The deleterious effects

of metals like cadmium are responsible for oxidative damage in living organisms. Previous investigations indicated that cadmium induces the formation of oxygen free radicals in tissues and inhibits the activity of some enzymes of the antioxidant defense system (Unsal *et al.* 2020). When organisms are exposed to elevated levels of trace metals, oxidative stress can damage essential biomolecules such as proteins, lipids will be happened, impairing physiological function (Singh *et al.* 2019; Haidar *et al.* 2023). In response, organisms have developed a variety of defense systems to mitigate the toxic effects of metals and maintain internal homeostasis. One of these mechanisms is the production of antioxidant enzymes such as superoxide dismutase (SOD), catalase, and glutathione, which work synergistically to neutralize reactive oxygen species (ROS) generated during metal exposure (Jomova *et al.* 2023). Cope with oxidative stress produced by ROS generation, SOD as the most powerful and the primary detoxification enzyme, removes the superoxide radical through the process of dismutation to oxygen and hydrogen peroxide. In the present study, changes in the activity of SOD were observed in *O. bicornis* under cadmium treatment. Similar to findings in other marine invertebrates exposed to Cu (Brown *et al.* 2004), or reported in field studies (Pérez *et al.* 2004), Cd may induce oxidative damage in *O. bicornis*, potentially through the generation of ROS. In response to oxidative stress, organisms exposed to various organic and metal contaminants often exhibit increased activity of antioxidant enzymes. However, these responses vary depending on the species, the enzymes involved, and whether the exposure is to a single or a mixture of contaminants (Regoli and Giuliani 2014). SOD activity in *O. bicornis* exposed to cadmium remained constant at 24h but showed an increase after 48h of exposure, with values still higher than those of the control at day 0. Our results are in full agreement with those reported in literature. Similar results were obtained in previous investigations of several polychaete species. According to (Sun and Zhou 2008), the activity of SOD in *N. diversicolor* exposed to Cd and petroleum hydrocarbons (PHCs) increased after a 6day exposure, and (Moreira *et al.* 2006) also reported a significant induction of this enzyme when the species was exposed to contaminated sediments. The increase in SOD activity suggested that SOD was induced as consequence of generation of O_2^- . This response likely reflects the role of the antioxidant enzyme SOD in removing excess free radicals, mainly O_2^- and H_2O_2 generated to counteract Cd-induced stress and mitigate their toxicity. However, (Chaâbane *et al.* 2020) showed that SOD activity enhanced in the gills and digestive gland of chromium (VI)-exposed *Venus verrucosa*. ROS accumulation is a widely documented effect of cadmium exposure in marine organisms leading to oxidative damage and cellular dysfunction (Liu *et al.* 2022). Cd does not induce ROS production directly; however, it can cause indirect damage and generate free radicals (Patra *et al.* 2011). This indirect damage was elucidated in the present study by a significant increase of H_2O_2 levels confirming the metal potential to boost pro-oxidants' production. The

enhanced H₂O₂ generation in *O. bicornis* was associated with an increase in LOOH levels, as compared to the control group. Hydrogen peroxide is a central redox signaling molecule, capable of serving as messenger to carry a redox signal from the site of its generation to a target site (Sies *et al.* 2017). The levels of H₂O₂ were found to be increased in *D. magna* adults exposed to the metal mixture (Cu/Cd), oxidative stress was found as a common mechanism underlying the toxicity of these metals (Majid 2024). These findings are in line with those of (Chetoui *et al.* 2022), and (Chaâbane *et al.* 2020). Our study found that Cd exposure at a sublethal concentration (LC₁₀) induced significant oxidative stress responses in *Ophelia bicornis*. Lipid hydroperoxides (LOOH) and hydrogen peroxide (H₂O₂) increased significantly over time, indicating heightened ROS production and lipid peroxidation. Concurrently, superoxide dismutase (SOD) activity increased after 48h, suggesting an enzymatic attempt to counteract ROS accumulation. However, the sustained elevation of H₂O₂ and LOOH, despite SOD activation, highlights an oxidative imbalance. This underscores the vulnerability of *O. bicornis* tissues to damage caused by Cd exposure. These findings regarding energy metabolism and stress responses may help predict the effects of Cd on marine organisms, particularly annelids such as *O. bicornis*, which are highly exposed to metal contaminants in their habitats. *O. bicornis* is especially important in ecotoxicological studies, living in direct contact with contaminated sediments.

5. Conclusion

The physiological and biochemical responses of the annelid *Ophelia bicornis* were investigated under laboratory conditions after short exposure to cadmium (Cd), focusing on antioxidant enzyme activity and the content of major biochemical components. These findings provide insight into the molecular responses triggered by Cd exposure and the underlying mechanisms in *O. bicornis*, emphasizing the significant role of oxidative stress in mediating the cellular response to cadmium-induced damage. responses of the antioxidant system and biochemical parameters in this species are not well-documented; thus, our study provides the first evidence addressing these aspects regarding cadmium toxicity. This research highlights the ecotoxicological relevance of *O. bicornis* as a sensitive bioindicator of cadmium exposure. Even at sublethal concentrations, Cd disrupted the energy balance in exposed organisms and induced oxidative stress. this is evidenced by increased carbohydrates mobilization, alterations of lipid contents, elevated levels lipid hydroperoxides (LOOH) and hydrogen peroxide (H₂O₂), and a time dependent modulation of superoxide dismutase (SOD) activity. These physiological and biochemical changes indicate that *O. bicornis* relies on its energy reserves and antioxidant defenses to cope with metal-induced stress. The consistent correlation between oxidative stress biomarkers and cadmium exposure further underscores their value as early warning indicators. Overall, these findings enhance our understanding of how benthic invertebrates respond to

metal-contaminated environments and confirm the sensitivity of *O. bicornis*. This establishes its use as a bioindicator in environmental monitoring and provides a baseline for evaluating its physiological responses to metal exposure and opens a new perspective for its application in future ecotoxicological and biomonitoring studies.

6. Acknowledgements

The authors would like to express their sincere gratitude to the Laboratory of Ecology, Biology and Physiology of Aquatic Organisms, Faculty of Sciences of Tunis, University of Tunis El Manar, Tunisia, for providing access to the necessary equipment and technical resources. The authors are also thankful to the staff of the Laboratory for their valuable technical assistance and support throughout this study, which greatly facilitated the realization of this scientific contribution.

References

- Amira, A., Merad, I., Almeida, C. M. R., Guimaraes, L. and Soltani, N. (2018). Seasonal variation in biomarker responses of *Donax trunculus* from the Gulf of Annaba (Algeria): Implication of metal accumulation in sediments. *Comptes Rendus, Géoscience*, **350**(4), 173-179, <https://doi.org/10.1016/j.crte.2018.02.002>.
- Barber, B. J. and Blake, N. J. (2006). Reproductive physiology, *Developments in aquaculture and fisheries science*, **35**, 357-416.
- Bat, L., Arici, E., Öztekin, A. and Yardim, Ö. (2016). A preliminary study on the heavy metal levels in *Ophelia bicornis* (Savigny, 1820) in the Black Sea, *Pakistan Journal of Marine Sciences*, **25**(1&2), 93-100.
- Baudou, F. G., Ossana, N. A., Castañé, P. M., Mastrángelo, M. M. and Ferrari, L. (2017). Cadmium effects on some energy metabolism variables in *Cnesterodon decemmaculatus* adults, *Ecotoxicology*, **26**(9), 1250-1258, <https://doi.org/10.1007/s10646-017-1850-z>.
- Beauchamp, C. and Fridovich, I. (1971). Superoxide dismutase: Improved assays and an assay applicable to acrylamide gels, *Analytical biochemistry*, **44**(1), 276-287.
- Belabed, S. and Soltani, N. (2018). Effects of cadmium concentrations on bioaccumulation and depuration in the marine bivalve *Donax trunculus*, *Euro-Mediterranean Journal for Environmental Integration*, **3**(1), 19, <https://doi.org/10.1007/s41207-018-0054-0>.
- Belabed, S. and Soltani, N. (2022). Cadmium exposure on *Donax trunculus* (Mollusca; Bivalvia) lethality parameters and biomarkers responses, *Fresenius Environmental Bulletin*, **231**, 11715.
- Benedetti, M., Giuliani, M. E. and Regoli, F. (2015). Oxidative metabolism of chemical pollutants in marine organisms: Molecular and biochemical biomarkers in environmental toxicology, *Annals of the New York Academy of Sciences*, **1340**(1), 8-19, <https://doi.org/10.1111/nyas.12698>.
- Boguszewska-Mańkowska, D., Nykiel, M. and Zagdańska, B. (2015). Protein oxidation and redox regulation of proteolysis, In *Basic Principles and Clinical Significance of Oxidative Stress*, *IntechOpen*, Chapter 2, <http://dx.doi.org/10.5772/61182>.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the

- principle of protein-dye binding, *Analytical biochemistry*, **72**(1-2), 248-254.
- Brown, R. J., Galloway, T. S., Lowe, D., Browne, M. A., Dissanayake, A., Jones, M. B. and Depledge, M. H. (2004). Differential sensitivity of three marine invertebrates to copper assessed using multiple biomarkers, *Aquatic toxicology*, **66**(3), 267-278.
- Chaâbane, M., Bejaoui, S., Trabelsi, W., Telahigue, K., Chetoui, I., Chalhaf, M., Zeghal, N. and Soudani, N. (2020). The potential toxic effects of hexavalent chromium on oxidative stress biomarkers and fatty acids profile in soft tissues of *Venus verrucosa*, *Ecotoxicology and Environmental Safety*, **196**, 110562.
- Chen, P., Bornhorst, J., Neely, M. D. and Avila, D. S. (2018). Mechanisms and disease pathogenesis underlying metal-induced oxidative stress, *Oxidative medicine and cellular longevity*, **2018**, 7612172.
- Chetoui, I., Bejaoui, S., Trabelsi, W., Rabeh, I., Nechi, S., Chelbi, E., Ghalghaf, M., El Cafsi, M. and Soudani, N. (2022). Exposure of *Mactra corallina* to acute doses of lead: Effects on redox status, fatty acid composition and histomorphological aspect, *Drug and Chemical Toxicology*, **45**(1), 311-323, <https://doi.org/10.1080/01480545.2019.1693590>.
- Chetoui, I., Ghribi, F., Fouzai, C., Trabelsi, W., Zereli, S., El Cafsi, M. and Soudani, N. (2021). Differential Fatty Acids and Stress Biomarkers Responses in Native (*Carcinus Aestuarii*) and Invasive (*Portunus Segnis*) Crabs Exposed to Environmental Pollution, <https://doi.org/10.21203/rs.3.rs-1119403/v1>
- Cosson, R. P. and Amiard, J.-C. (2021). Use of metallothioneins as biomarkers of exposure to metals, In *Use of biomarkers for environmental quality assessment*, (p. 79-111). CRC Press.
- De Coen, W. M. and Janssen, C. R. (2003). The missing biomarker link: Relationships between effects on the cellular energy allocation biomarker of toxicant-stressed *Daphnia magna* and corresponding population characteristics, *Environmental toxicology and chemistry*, **22**(7), 1632-1641.
- Dean, H. K. (2008). The use of polychaetes (Annelida) as indicator species of marine pollution: A review, *Revista de Biología Tropical*, **56**(4), 11-38.
- Demirci-Çekiç, S., Özkan, G., Avan, A. N., Uzunboy, S., Çapanoğlu, E. and Apak, R. (2022). Biomarkers of oxidative stress and antioxidant defense, *Journal of pharmaceutical and biomedical analysis*, **209**, 114477.
- Depledge, M. H. (2020). The rational basis for the use of biomarkers as ecotoxicological tools, In *Nondestructive biomarkers in vertebrates*, (p. 271-295). CRC Press.
- Di Salvatore, P., Calcagno, J. A., Ortíz, N., de Molina, M. del C. R. and Sabatini, S. E. (2013). Effect of seasonality on oxidative stress responses and metal accumulation in soft tissues of *Aulacomya atra*, a mussel from the South Atlantic Patagonian coast, *Marine environmental research*, **92**, 244-252.
- Duchateau, G. and Florin, M. (1959). Blood trehalose level in insects and its significance, *Archives internationales de physiologie et de biochimie*, **67**(2), 306-314.
- Fouzai, C., Trabelsi, W., Bejaoui, S., Telahigue, K., Rabeh, I., Nechi, S., Chelbi, E. and Soudani, N. (2020). Cellular toxicity mechanisms of lambda-cyhalothrin in *Venus verrucosa* as revealed by fatty acid composition, redox status and histopathological changes, *Ecological Indicators*, **108**, 105690.
- Gambi, M. C. and Giangrande, A. (1986). Distribution of soft-bottom polychaetes in two coastal areas of the Tyrrhenian Sea (Italy): Structural analysis, *Estuarine, Coastal and Shelf Science*, **23**(6), 847-862.
- Gashkina, N. A. (2024). Metal toxicity: Effects on energy metabolism in fish, *International Journal of Molecular Sciences*, **25**(9), 5015.
- Goldsworthy, G. J., Mordue, W. and Guthkelch, J. (1972). Studies on insect adipokinetic hormones, *General and Comparative Endocrinology*, **18**(3), 545-551.
- Haidar, Z., Fatema, K., Shoily, S. S. and Sajib, A. A. (2023). Disease-associated metabolic pathways affected by heavy metals and metalloid, *Toxicology reports*, **10**, 554-570.
- Hand, S. C. and Hardewig, I. (1996). Downregulation of cellular metabolism during environmental stress: Mechanisms and implications, *Annual review of physiology*, **58**, 539-563.
- Hoeger, U. and Schenk, S. (2024). Reproduction in Nereidid Polychaetes: Physiological and Biochemical Aspects, In *Frontiers in Invertebrate Physiology: A Collection of Reviews* (pp. 33-90), Apple Academic Press.
- Holmstrup, M., Bindesbøl, A.-M., Oostingh, G. J., Duschl, A., Scheil, V., Köhler, H.-R., Loureiro, S., Soares, A. M., Ferreira, A. L. and Kienle, C. (2010). Interactions between effects of environmental chemicals and natural stressors: A review, *Science of the Total Environment*, **408**(18), 3746-3762.
- Hutchings, P. (1998). Biodiversity and functioning of polychaetes in benthic sediments, *Biodiversity and Conservation*, **7**(9), 1133-1145, <https://doi.org/10.1023/A:1008871430178>.
- Ivanina, A. V., Cherkasov, A. S. and Sokolova, I. M. (2008). Effects of cadmium on cellular protein and glutathione synthesis and expression of stress proteins in eastern oysters, *Crassostrea virginica* Gmelin, *Journal of Experimental Biology*, **211**(4), 577-586.
- Jiang, Z.-Y., Hunt, J. V. and Wolff, S. P. (1992). Ferrous ion oxidation in the presence of xylenol orange for detection of lipid hydroperoxide in low density lipoprotein, *Analytical biochemistry*, **202**(2), 384-389.
- Jomova, K., Raptova, R., Alomar, S. Y., Alwasel, S. H., Nepovimova, E., Kuca, K. and Valko, M. (2023). Reactive oxygen species, toxicity, oxidative stress, and antioxidants: Chronic diseases and aging, *Archives of Toxicology*, **97**(10), 2499-2574, <https://doi.org/10.1007/s00204-023-03562-9>.
- Kassambara, A. and Mundt, F. (2020). *factoextra: Extract and visualize the results of multivariate data analyses*. R package.
- Ketata, I., Smaoui-Damak, W., Guerhazi, F., Rebai, T. and Hamza-Chaffai, A. (2007). In situ endocrine disrupting effects of cadmium on the reproduction of *Ruditapes decussatus*, *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, **146**(3), 415-430.
- Labiod, R. and Belabed, S. (2024). Assessment of the acute toxicity of Cadmium on *Ophelia bicornis* (Polychaeta: Opheliidae), *Tobacco Science and Technology*, 2331-2339.
- Le Gal, y., Lagadic, L., Le Bras, S. and Caquet, T. (1997). Energy load in adenylates (CEA) and other biomarkers associated with energy metabolism, *Biomarkers in ecotoxicology*, fundamental aspects, Paris: 1-9.
- Lê, S., Josse, J. and Husson, F. (2008). FactoMineR: An R package for multivariate analysis. *Journal of Statistical Software*, **25**(1), 1–18. <https://doi.org/10.18637/jss.v025.i01>

- Liu, Y., Chen, Q., Li, Y., Bi, L., Jin, L. and Peng, R. (2022). Toxic effects of cadmium on fish, *Toxics*, **10**(10), 622.
- Liu, Z., Lv, W., Huang, Y., Fan, B., Li, Y. and Zhao, Y. (2016). Effects of cadmium on lipid metabolism in female estuarine crab, *Chironomus dehaani*, *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, **188**, 9-16.
- Louis, F., Delahaut, L., Gaillet, V., Bonnard, I., Paris-Palacios, S. and David, E. (2021). Effect of reproduction cycle stage on energy metabolism responses in a sentinel species (*Dreissena polymorpha*) exposed to cadmium: What consequences for biomonitoring? *Aquatic Toxicology*, **230**, 105699.
- Lucia, M., André, J.-M., Gonzalez, P., Baudrimont, M., Bernadet, M.-D., Gontier, K., Maury-Brachet, R., Guy, G. and Davail, S. (2010). Effect of dietary cadmium on lipid metabolism and storage of aquatic bird *Cairina moschata*, *Ecotoxicology*, **19**(1), 163-170, <https://doi.org/10.1007/s10646-009-0401-7>.
- Luis, O. J. and Passos, A. M. (1995). Seasonal changes in lipid content and composition of the polychaete *Nereis (Hediste) diversicolor*, *Comparative biochemistry and physiology part B: biochemistry and molecular biology*, **111**(4), 579-586.
- Majid, S. (2024). Comparative analysis of metal toxicity responses in aquatic invertebrate and vertebrate model organisms, *doctoral thesis*, 263.
- Makhdoumi, P., Karimi, H. and Khazaei, M. (2020). Review on Metal-Based Nanoparticles: Role of Reactive Oxygen Species in Renal Toxicity, *Chemical Research in Toxicology*, **33**(10), 2503-2514, <https://doi.org/10.1021/acs.chemrestox.9b00438>.
- Maltagliati, F., Casu, M., Lai, T., Sareri, D. I., Casu, D., Galletti, M. C., Cantone, G. and Castelli, A. (2005). Taxonomic distinction of *Ophelia barquii* and *O. bicornis* (Annelida : Polychaeta) in the Mediterranean as revealed by ISSR markers and the number of nephridiopores, *Journal of the Marine Biological Association of the United Kingdom*, **85**(4), 835-841.
- Maruthai, S., Selvanarayanan, R., Thanarajan, T., & Rajendran, S. (2025). Hybrid vision GNNs based early detection and protection against pest diseases in coffee plants. *Scientific reports*, **15**(1), 11778.
- Mathieu, M. and Lubet, P. (1993). Storage tissue metabolism and reproduction in marine bivalves—A brief review, *Invertebrate Reproduction & Development*, **23**(2-3), 123-129, <https://doi.org/10.1080/07924259.1993.9672303>.
- Mayer, F. L., Versteeg, D. J., McKee, M. J., Folmar, L. C., Graney, R. L., McCume, D. C. and Rattner, B. A. (2018). Physiological and nonspecific biomarkers, In *Biomarkers* (p. 5-86). CRC Press.
- Merad, I., Rabei, A. and Soltani, N. (2024). Sex-specific differences in metallothionein induction by cadmium in different organs of *Donax trunculus* Linnaeus, 1758, *Molluscan Research*, **44**(1), 11-21, DOI: 10.1080/13235818.2023.2277834.
- Meyer, B., Atkinson, A., Stöbing, D., Oettl, B., Hagen, W. and Bathmann, U. V. (2002). Feeding and energy budgets of Antarctic krill *Euphausia superba* at the onset of winter—I. Furcilia III larvae, *Limnology and Oceanography*, **47**(4), 943-952, <https://doi.org/10.4319/lo.2002.47.4.0943>.
- Mohandas, P., Subramanian, P., & Surendran, R. (2025a). AI-Driven ResNet50 with Channel Attention for Tamil Nadu Air Pollution Level Classification. *2025 4th International Conference on Innovative Mechanisms for Industry Applications (ICIMIA)*, 1508-1513.
- Mohandas, P., Subramanian, P., & Surendran, R. (2025b). Deep Feedforward Air Pollution Classification Empowered by Sophia-G Optimization. *2025 4th International Conference on Innovative Mechanisms for Industry Applications (ICIMIA)*, 1696-1702.
- Mohandas, P., Subramanian, P., & Surendran, R. (2025c). Optimizing Air Pollution Prediction in Urban Environments using a Hybrid RNN-PBO Model with IoT Data. *2025 International Conference on Visual Analytics and Data Visualization (ICVADV)*, 480-487.
- Moreira, S. M., Lima, I., Ribeiro, R. and Guilhermino, L. (2006). Effects of estuarine sediment contamination on feeding and on key physiological functions of the polychaete *Hediste diversicolor*: Laboratory and in situ assays, *Aquatic toxicology*, **78**(2), 186-201.
- Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E. and Wagner, H. (2022). *vegan: Community ecology package*. R package.
- Ou, P. and Wolff, S. P. (1996). A discontinuous method for catalase determination at 'near physiological' concentrations of H₂O₂ and its application to the study of H₂O₂ fluxes within cells, *Journal of biochemical and biophysical methods*, **31**(1-2), 59-67.
- Pan, K. and Wang, W.X. (2012). Trace metal contamination in estuarine and coastal environments in China, *Science of the total environment*, **421**, 3-16.
- Patra, R. C., Rautray, A. K. and Swarup, D. (2011). Oxidative Stress in Lead and Cadmium Toxicity and Its Amelioration, *Veterinary Medicine International*, 1-9, <https://doi.org/10.4061/2011/457327>.
- Pérez, E., Blasco, J. and Solé, M. (2004). Biomarker responses to pollution in two invertebrate species: *Scrobiculariaplana* and *Nereis diversicolor* from the Cádiz Bay (SW Spain), *Marine Environmental Research*, **58**(2-5), 275-279.
- Pook, C., Lewis, C. and Galloway, T. (2009). The metabolic and fitness costs associated with metal resistance in *Nereis diversicolor*, *Marine Pollution Bulletin*, **58**(7), 1063-1071.
- R Core Team (2025). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Reddy, P. S., Devi, M., Sarojini, R., Nagabhusanam, R. and Fingerma, M. (1994). Cadmium chloride induced hyperglycemia in the red swamp crayfish, *Procambarus clarkii*: Possible role of crustacean hyperglycemic hormone, *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology*, **107**(1), 57-61.
- Regoli, F. and Giuliani, M. E. (2014). Oxidative pathways of chemical toxicity and oxidative stress biomarkers in marine organisms, *Marine environmental research*, **93**, 106-117.
- Scaps, P. (2000). The exploitation of polychaete annelids and their relevance in ecotoxicological studies, Habilitation of supervise research, University of Lille 1, pp. 1-63.
- Sharma, H., Rawal, N. and Mathew, B. B. (2015). The characteristics, toxicity and effects of cadmium, *international journal of nanotechnology and nanoscience*, **3**(10), 1-9.
- Shibko, S., Koivistoinen, P., Tratnyneck, C., Newhall, A. and Freidman, L. (1966). A method for the sequential quantitative separation and glycogen from a single rat liver homogenate or from a sub cellular fraction, *Analytical*

- Biochemistry*, **19**, 415–428, [https://doi.org/10.1016/0003-2697\(67\)90242-4](https://doi.org/10.1016/0003-2697(67)90242-4).
- Sies, H., Berndt, C. and Jones, D. P. (2017). Oxidative Stress, *Annual Review of Biochemistry*, **86**(1), 715-748, <https://doi.org/10.1146/annurev-biochem-061516-045037>.
- Sinenko, S. A., Starkova, T. Y., Kuzmin, A. A. and Tomilin, A. N. (2021). Physiological signaling functions of reactive oxygen species in stem cells: From flies to man, *Frontiers in cell and developmental biology*, **9**, 714370.
- Singh, A., Kukreti, R., Saso, L. and Kukreti, S. (2019). Oxidative stress: A key modulator in neurodegenerative diseases, *Molecules*, **24**(8), 1583.
- Singh, C., Singh, R. and Shekhar, A. (2024). Oxidative Stress in Cadmium Toxicity in Animals and Its Amelioration, In *Cadmium Toxicity Mitigation* (p. 391-411). Cham: Springer Nature Switzerland, https://doi.org/10.1007/978-3-031-47390-6_15.
- Sivasubramanian, S., Venkatesan, S. K. G., Thanarajan, T., & Rajendran, S. (2025). Al-Biruni Earth Radius Optimization for enhanced environmental data analysis in remote sensing imagery. *Agrociencia*, 1-18.
- Smolders, R., De Boeck, G. and Blust, R. (2003). Changes in cellular energy budget as a measure of whole effluent toxicity in zebrafish (*Danio rerio*), *Environmental Toxicology and Chemistry*, **22**(4), 890-899, <https://doi.org/10.1002/etc.5620220429>.
- Song, C., Sun, C., Liu, B. and Xu, P. (2023). Oxidative stress in aquatic organisms, *Antioxidants*, **12**(6), 1223.
- Tripp-Valdez, M. A., Bock, C., Lannig, G., Koschnick, N., Pörtner, H. O. and Lucassen, M. (2019). Assessment of muscular energy metabolism and heat shock response of the green abalone *Haliotis fulgens* (Gastropoda: Philipi) at extreme temperatures combined with acute hypoxia and hypercapnia, *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, **227**, 1-11.
- Unsal, V., Dalkıran, T., Çiçek, M. and Kölükcü, E. (2020). The role of natural antioxidants against reactive oxygen species produced by cadmium toxicity: A review, *Advanced pharmaceutical bulletin*, **10**(2), 184.
- Valavanidis, A., Vlahogianni, T., Dassenakis, M. and Scoullos, M. (2006). Molecular biomarkers of oxidative stress in aquatic organisms in relation to toxic environmental pollutants, *Ecotoxicology and environmental safety*, **64**(2), 178-189.
- Wang, X., Li, E. and Chen, L. (2016). A review of carbohydrate nutrition and metabolism in crustaceans, *North American Journal of Aquaculture*, **78**(2), 178-187.
- Wang, Y., Li, E., Yu, N., Wang, X., Cai, C., Tang, B., Chen, L. and Van Wormhoudt, A. (2012). Characterization and expression of glutamate dehydrogenase in response to acute salinity stress in the Chinese mitten crab, *Eriocheir sinensis*, *PLoS One*, **7**(5), e37316.
- Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag, New York.
- Widdows, J., Donkin, P., Staff, F. J., Matthiessen, P., Law, R. J., Allen, Y. T., Thain, J. E., Allchin, C. R. and Jones, B. R. (2002). Measurement of stress effects (scope for growth) and contaminant levels in mussels (*Mytilus edulis*) collected from the Irish Sea, *Marine Environmental Research*, **53**(4), 327-356.
- Wright, D. A. and Welbourn, P. M. (1994). Cadmium in the aquatic environment: A review of ecological, physiological, and toxicological effects on biota, *Environmental Reviews*, **2**(2), 187-214, <https://doi.org/10.1139/a94-012>.
- Yang, J., Liu, D., Jing, W., Dahms, H.-U. and Wang, L. (2013). Effects of cadmium on lipid storage and metabolism in the freshwater crab *Sinopotamon henanense*. *PLoS One*, **8**(10), e77569.
- Zhou, Q. X., Kong, F. X. and Zhu, L. (2004). *Ecotoxicology (in Chinese)* (Beijing, China: Science Press).